

## Isotopic fractionation in open systems: application to organic matter decomposition in ocean and land

R. Ramesh and Arvind Singh\*

Geosciences Division, Physical Research Laboratory, Ahmedabad 380 009, India

**Stable isotopic evolution of a reservoir from which material is continuously removed with isotopic fractionation is generally described by the Rayleigh equation, widely used in geochemical research to describe two-component systems. It has been extended to the case of multi-component sources as well. Here we present an equation describing isotopic fractionation in open systems, wherefrom material is not only removed with isotopic fractionation, but fresh material of a different isotopic composition is added from an external source, at a given rate, and some examples where this equation is applicable.**

**Keywords:** Denitrification, isotope fractionation, Rayleigh equation, soil organic matter.

LET  $R$  be the (non-radiogenic) stable isotopic ratio under consideration (e.g.  $^{18}\text{O}/^{16}\text{O}$ , D/H,  $^{13}\text{C}/^{12}\text{C}$ ,  $^{15}\text{N}/^{14}\text{N}$ , etc.), denoted by  $N^*/N$ , where  $N^*$  and  $N$  are respectively the number of molecules containing the heavier and lighter isotopes of a given mass of the same element (e.g. oxygen;  $\text{H}_2^{18}\text{O}$  and  $\text{H}_2^{16}\text{O}$ ;  $N^* \ll N$ ; typical values in the ocean are 2000 and  $10^6$  respectively, in  $\sim 3 \times 10^{-17}$  g of water). Taking the logarithm and differentiating, we obtain

$$dR/R = dN^*/N^* - dN/N = dN^*/(RN) - dN/N. \quad (1)$$

At any instant  $dN_e^*$  and  $dN_e$  of the isotopically heavier and lighter molecules, respectively, are being removed by some physical process (e.g. evaporation from a lake) during an infinitesimal time interval  $\Delta t$ . Isotopic fractionation implies that  $dN_e^*$  is not equal to  $dN_e$ . In such a case, we define a fractionation factor between the material leaving (e.g. vapour) and the material left behind (e.g. water) as  $\alpha = [dN_e^*/dN_e]/R$ . In per mil units (‰), this fractionation factor is represented by  $\varepsilon = (\alpha - 1) \times 10^3$ . We restrict our consideration to isothermal processes where  $\alpha$  and  $\varepsilon$  are constants. As in the case of Rayleigh fractionation<sup>1-11</sup>, we assume that the reservoir remains isotopically homogeneous and has no isotopic gradients within, at any time. Let  $dN_r^*$  and  $dN_r$  of the heavier and lighter molecules be added to the reservoir during the interval  $\Delta t$ , but without fractionation (i.e.  $dN_r^*/dN_r$  is the same as  $N_r^*/N_r$ , the isotopic composition of the contributing source, which remains a constant  $R_r \neq \alpha R$ ). Therefore, the net changes in the numbers of heavier and lighter isotopic

molecules during  $\Delta t$  are  $dN^* = dN_r^* - dN_e^*$  and  $dN = dN_r - dN_e$  respectively, and their ratio,

$$dN^*/dN = (\beta R_r - \alpha R)/(\beta - 1), \quad (2)$$

where  $\beta$  is  $dN_r/dN_e$ , the ratio of the amount of material added to that lost (the case  $\beta = 1$  causes a singularity in eq. (2) and is discussed as a special case later). Integrating eq. (2) using eq. (1), with initial value of  $R$  as  $R_0$ , we obtain (see Appendix 1 for details):

$$R = R_0 f^\rho + [\beta R_r/(\alpha + \beta - 1)][1 - f^\rho], \quad \text{when } \beta \neq 1. \quad (3)$$

Here  $f$  is the fraction of the material left in the reservoir relative to its initial amount, i.e.  $(N^* + N)/(N_0^* + N_0)$ , approximated as  $(N/N_0)$ , and  $\rho = \alpha/(1 - \beta) - 1$ . When  $\beta = 0$ , i.e., when there is only loss of material and no addition, this reduces to  $R = R_0 f^{\alpha-1}$ , the classical Rayleigh fractionation equation. Using the  $\delta$  notation for isotope ratios, taking the initial composition  $R_0$  of the reservoir to be the reference value relative to which all other  $\delta$  values are expressed, we have  $\delta = (R/R_0 - 1)10^3\%$  and  $\delta_r = (R_r/R_0 - 1)10^3\%$ . With this standard notation eq. (3) becomes

$$\delta = [1 - f^\rho](\beta\delta_r - \varepsilon)/(\alpha + \beta - 1), \quad \text{when } \beta \neq 1. \quad (4)$$

For the case where  $\beta = 1$ , we cannot express the isotopic ratio of the reservoir as a function of  $f$  because it remains constant at unity (i.e.  $N$  remains as  $N_0$ ), as the amount of material lost is compensated exactly by the incoming material from the source. A different approach is therefore necessary. The change in the number of heavier molecules is given, as before, by  $dN^* = dN_r^* - dN_e^*$ , while the change in the number of lighter molecules is  $dN = dN_r - dN_e = 0$ , as  $\beta = dN_r/dN_e = 1$ . Therefore, the change in the isotopic ratio of the reservoir is  $dR = R_r(dN_r/N) - \alpha R(dN_e/N)$  (mass balance is used:  $RdN = 0$  as  $dN = 0$ ). This can be rewritten as

$$dR/(R_r - \alpha R) = dN_r/N_0. \quad (5)$$

Integrating this with initial value of  $R$  as  $R_0$ , we get:

$$R = R_0 \exp[-(\alpha N_r/N_0)] + (R_r/\alpha) \{1 - \exp[-(\alpha N_r/N_0)]\}. \quad (6)$$

Here  $N_r$  indicates the total number of lighter isotopic molecules added thus far. In the  $\delta$  notation, eq. (6) becomes:


$$\delta = \{(\delta_r - \varepsilon)/\alpha\} \{1 - \exp[-(\alpha N_r/N_0)]\}. \quad (7)$$

It is clear that for large mixing (i.e.  $N_r \gg N_0$ ), the reservoir isotopic composition ( $\delta$ ) saturates to  $(\delta_r - \varepsilon)/\alpha$ , rather than  $\delta_r$  expected in the case of simple mixing, as


\*For correspondence. (e-mail: arvinds@prl.res.in)

here both mixing and removal by fractionation occur simultaneously. However, when  $\alpha$  is small (e.g. close to unity), thus  $\delta$  reaches  $\delta_r - \epsilon$ . We note that eq. (4) was derived in another form by Mook<sup>4</sup>. The case of alteration of radiogenic stable isotopes undergoing isotopic exchange with meteoric water was treated similarly by Albarede<sup>12</sup>. On the other extreme, when  $N_r \ll N_0$ ,  $\delta \approx (\delta_r - \epsilon)(N_r/N_0)$ .

Figure 1 shows an example of the isotopic ( $\delta$ ) evolution of a reservoir as a function of the fraction ( $f$ ) left of the initial amount. The fractionation factor  $\alpha$  has been taken as 1.01 (i.e.  $\epsilon = 10\%$ ) and isotopic composition of the source contributing to the reservoir, as  $-10\%$ . As the material lost from the reservoir is isotopically heavier in this example (i.e.  $\alpha > 1$ ), the isotopic composition of the remaining material rapidly decreases and reaches close to  $-46\%$  (from a value of 0), when  $f$  becomes small ( $\sim 0.01$ ); this is the Rayleigh case where no material is added to the reservoir ( $\beta = 0$  in Figure 1). As  $\beta$  values increase, the reservoir  $\delta$  value is brought up higher and higher, as the source steadily contributes material with a constant isotopic composition ( $\delta_r$ ) of  $-10\%$ . The saturation value is  $\delta_r - \epsilon = -10 - 10 = -20\%$ , as expected from eq. (7).



**Figure 1.** An example of the evolution of the isotopic composition ( $\delta$ ) of a reservoir as a function of the fraction ( $f$ ) of the original material left, for different values of  $\beta$ , the ratio of instantaneous amount added to that lost ( $\beta \neq 1$ ). The isotopic composition of the external source is taken as  $-10\%$  and the fractionation factor, 1.01.




**Figure 2.** An example of the evolution of the isotopic composition ( $\delta$ ) of a reservoir as a function of the fraction ( $f$ ) of the original material left, for different values of  $\beta$ , the ratio of instantaneous amount added to that lost ( $\beta \neq 1$ ). The isotopic composition of the external source is taken as  $-10\%$  and the fractionation factor, 0.99.

Figure 2 shows another example where fractionation factor has been taken as 0.99 ( $\epsilon = -10\%$ ) and isotopic composition of the source contributing to the reservoir, as  $-10\%$ . As the material lost from the reservoir is isotopically lighter in this example (i.e.  $\alpha < 1$ ), the isotopic composition of the remaining material rapidly increases and reaches close to  $+47\%$  (from a value of 0), when  $f$  becomes small ( $\sim 0.01$ ); this is the Rayleigh case where no material is added to the reservoir (closed system,  $\beta = 0$  in Figure 2). As  $\beta$  increases, the reservoir  $\delta$  value is lowered further and further, as the source steadily contributes material with a constant isotopic composition ( $\delta_r$ ) of  $-10\%$ . The saturation value is  $\delta_r - \epsilon = -10 - (-10) = 0\%$ , as expected from eq. (7).

We further discuss a general example for a special case of  $\beta = 1$ , i.e. the rates of addition and removal of material remain same at any instant. Figure 3 shows the isotopic ( $\delta$ ) evolution of a reservoir as a function of  $N_r/N_0$  (ratio of lighter isotopic molecules being added by some source to the number of initial lighter isotopic molecules of the reservoir) for different combinations of enrichment factor ( $\epsilon$ ) and isotopic composition of material being added ( $\delta_r$ ). Initial isotopic composition of the reservoir is taken  $5\%$ . The enrichment factor  $\epsilon$  has been taken as  $-25\%$  and  $-22\%$  and  $\delta_r$  as  $10\%$  and  $5\%$ . Keeping  $\delta_r$  constant,  $\delta$  in the case of more negative  $\epsilon$  increases faster than in the case of less negative  $\epsilon$ . While keeping  $\epsilon$  constant,  $\delta$  in the case of higher  $\delta_r$  increases faster than that in the case of lower  $\delta_r$ .

We now discuss some examples where the given equations are applicable: ocean denitrification and soil organic matter decomposition. Data on both processes are limited. We extracted the data on isotopic composition ( $\delta$ ) and  $f$  values from the literature using origin package.

Primary productivity in surface ocean is generally limited due to unavailability of nutrients. However, some parts of the world oceans are highly productive due to nutrient inputs by upwelling, riverine inputs, etc. Decom-




**Figure 3.** An example (for  $\beta = 1$ , i.e. the rate at which material is being added is equal to the removal rate) of the evolution of the isotopic composition ( $\delta$ ) of a reservoir as a function of  $N_r/N_0$  (ratio of the total number of lighter isotopic molecules being added from an external source to the number of initial lighter isotopic molecules of the reservoir) for different combinations of  $\epsilon$  and  $\delta_r$ .

position of organic matter in these regions at intermediate ocean depths (~500 m) leads to oxygen deficiency at these depths. In such regions microorganisms use nitrate as an oxidant to decompose organic matter and produce  $\text{N}_2\text{O}$  and  $\text{N}_2$  gases ( $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO} \rightarrow \text{N}_2\text{O} \rightarrow \text{N}_2$ ). This microbially catalysed multi-step process is known as denitrification<sup>13,14</sup>. Ocean loses bioavailable nitrogen ( $\text{NO}_3^-$ ) through this process. Denitrification occurs in oxygen minimum zones (OMZ) of the world oceans, e.g. the eastern tropical North Pacific and the Arabian Sea<sup>15,16</sup>. During denitrification, the nitrate concentration,  $[\text{NO}_3^-]$ , in OMZ (generally 150–1000 m depth) decreases, i.e.  $f$  decreases. Denitrifying bacteria reduce lighter nitrate ( $^{14}\text{NO}_3^-$ ) preferentially; hence the remaining  $\text{NO}_3^-$  gets enriched in heavier isotopes ( $^{15}\text{N}$ ). At the same time,  $\text{NO}_3^-$  is being added by settling organic matter from above, by decay. If this is unaccounted for, the isotopic composition of the  $\text{NO}_3^-$  reservoir can be significantly different from that expected from a simple Rayleigh model. Unlike earlier studies, we apply eq. (4), which takes this effect into account, to understand the isotopic evolution of nitrate in the OMZ of the Arabian Sea using literature data<sup>17</sup>.

Brandes *et al.*<sup>17</sup> obtained values for the enrichment factor ( $\epsilon$ ) and initial isotopic composition of nitrate ( $\delta_0$ ) of  $-22\text{‰}$  and  $6\text{‰}$ , respectively, in the Arabian Sea using a simple Rayleigh model and the measured  $\delta^{15}\text{N}$  values of dissolved  $\text{NO}_3^-$ . We know that organic matter steadily rains down to the deeper ocean and therefore treating this system with a simple Rayleigh equation may lead to an incorrect estimation of the fractionation factor. This is borne out by laboratory derived  $\epsilon$  values, which show large variations  $-17\text{‰}$  to  $-29\text{‰}$  (refs 18–20). We use different combinations of  $\epsilon$  ( $-22\text{‰}$  and  $-10\text{‰}$ ) and  $\delta_r$  values ( $5\text{‰}$  and  $10\text{‰}$ ) to understand the isotopic evolution of nitrate and estimate the fraction of material added.  $\beta$  values are taken as 0, 0.2, 0.5 and 0.8. As the data are from Brandes *et al.*<sup>17</sup>, for  $\epsilon = -22\text{‰}$ , it shows no addition from an external source (Figure 4a), even by changing the  $\delta_r$  value  $10\text{‰}$  to  $5\text{‰}$  as  $f$  decreases from 1 to 0.75 (Figure 4a and c). But when  $f$  decreases further (0.75–0.70), ~20% contribution from external source is observed. In Figure 4b and d, when  $\epsilon$  value is  $-10\text{‰}$ , results show that ~50% of the nitrate that is lost during denitrification is compensated by organic matter raining from above, for a small initial decrease in  $f$ ; if  $f$  decreases further, this contribution rises to ~80%. Thus by fixing  $\beta$ , the ratio of the rates of nitrate release due to fresh organic matter getting oxidized and the *in situ* denitrification, we can constrain the fractionation factor better with the eq. (4). In this special case  $\delta_r = \delta_0$ .

Now we discuss another example where soil organic matter (SOM) is decomposed by bacteria up to 100 cm depth. Similar to oceanic denitrification, these decomposing organisms also preferentially release lighter isotopic molecules ( $^{12}\text{C}$ ) for respiration and hence remaining SOM


enriched in heavier isotopes ( $^{13}\text{C}$ ) (refs 21 and 22). Decrease in the total carbon content and increase in  $\delta^{13}\text{C}$  are signatures of fractionation during decomposition of SOM. There is a supply of organic matter from above by diffusion/leaching, so eq. (4) is suitable to characterize this process<sup>23</sup>. Surface isotopic composition of SOM is taken as initial isotopic composition ( $\delta_0$ ; freshly deposited organic matter). The remaining fraction  $f$  of SOM is cal-



**Figure 4.** An example of denitrification in the sea: open squares in all the plots show the variation of  $\delta^{15}\text{N}$  of dissolved  $\text{NO}_3^-$  with  $f$  (fraction of  $\text{NO}_3^-$  remaining) from experimental data<sup>17</sup>. Initial isotopic value of nitrate ( $\delta_0$ ) is  $5.7\text{‰}$  and  $\beta$  values are taken as 0, 0.2, 0.5 and 0.8. Dotted and solid lines are plotted using eq. (4) for different  $\beta$  values. **a**,  $\epsilon = -22\text{‰}$  and  $\delta_r = 10\text{‰}$ ; **b**,  $\epsilon = -10\text{‰}$  and  $\delta_r = 10\text{‰}$ ; **c**,  $\epsilon = -22\text{‰}$  and  $\delta_r = 5\text{‰}$ ; **d**,  $\epsilon = -10\text{‰}$  and  $\delta_r = 5\text{‰}$ .

culated as the ratio of carbon content at a particular depth and the surface<sup>23</sup>. Accoe *et al.*<sup>23</sup> have estimated an  $\epsilon$  value  $-2.3\text{‰}$  using the simple Rayleigh model. We use four different combinations of  $\epsilon$  ( $-2.3\text{‰}$  and  $-1\text{‰}$ ) and  $\delta_r$  ( $-25\text{‰}$  and  $-30\text{‰}$ ) values in eq. (4) to analyse the same data. In Figure 5 *a* and *b*, when  $\delta_r$  remains constant ( $-25\text{‰}$ ), an

increase in enrichment factor shows that contribution of SOM from above increases (i.e.  $\beta$  value increases). In Figure 5 *c*, when  $f$  decreases from 1 to 0.7, no external contribution ( $\beta = 0$ ) is seen. Further decrease in  $f$  shows  $\beta = 0.2$ . When  $\epsilon = -1\text{‰}$  and  $\delta_r = -30\text{‰}$  (Figure 4 *d*), data points are difficult to explain especially when  $f$  decreases from 0.7 to 0.0. This happens because unlike all other cases, here  $\delta$  value of the external source is less than the initial reservoir value. While increasing  $\beta$ , isotopic composition of the resultant SOM cannot be increased, therefore it saturates. This can be used as an indication of absence of carbon percolation from above.



**Figure 5.** Example where soil organic matter (SOM) is decomposed by microorganisms: open squares in all the plots show the variation of  $\delta^{13}\text{C}$  of SOM variation with  $f$  (fraction of SOM remaining) from experimental data<sup>23</sup>. Initial isotopic value of SOM ( $\delta_i$ ) is  $-29.8\text{‰}$  and  $\beta$  values are taken as 0, 0.2, 0.5 and 0.8. Dotted and solid lines are derived using eq. (4) for different  $\beta$  values. *a*,  $\epsilon = -2.3\text{‰}$  and  $\delta_r = -25\text{‰}$ ; *b*,  $\epsilon = -1\text{‰}$  and  $\delta_r = -25\text{‰}$ ; *c*,  $\epsilon = -2.3\text{‰}$  and  $\delta_r = -30\text{‰}$ ; *d*,  $\epsilon = -1\text{‰}$  and  $\delta_r = -30\text{‰}$ .

**Appendix 1**

We consider a reservoir that loses material due to any physicochemical process with isotopic fractionation at a constant temperature. At any instant the stable isotope ratio is given by

$$R = \frac{N^*}{N} \tag{1A}$$

Taking logarithm of both sides of eq. (1A)

$$\ln R = \ln N^* - \ln N \tag{2A}$$

After differentiating, eq. (2A) becomes

$$\frac{dR}{R} = \frac{dN^*}{N^*} - \frac{dN}{N} = \frac{dN^*}{NR} - \frac{dN}{N} \tag{3A}$$

At any instant, let  $dN_e^*$  and  $dN_e$  are being lost and  $dN_r^*$  and  $dN_r$  are being added to the reservoir. So the net change in  $N^*$  and  $N$  is

$$dN^* = dN_r^* - dN_e^* \tag{4A}$$

$$dN = dN_r - dN_e \tag{5A}$$

Dividing eq. (4A) by eq. (5A)

$$\begin{aligned} \frac{dN^*}{dN} &= \frac{dN_r^* - dN_e^*}{dN_r - dN_e} = \frac{\frac{dN_r^*}{dN_r} \times \frac{dN_r}{dN_e} - \frac{dN_e^*}{dN_e}}{\frac{dN_r}{dN_e} - 1} \\ &= \frac{\beta R_r - \alpha R}{\beta - 1} \end{aligned} \tag{6A}$$

where  $\alpha$  is defined as  $(dN_e^*/dN_e)/(N^*/N)$ .

Substituting the value of  $dN^*$  from eq. (6A) to eq. (3A)

$$\frac{dR}{R} = \left( \frac{\beta R_r - \alpha R}{\beta - 1} \right) \frac{dN}{NR} - \frac{dN}{N} = \frac{dN}{N} \left( \frac{\beta \frac{R_r}{R}}{\beta - 1} - \frac{\alpha}{\beta - 1} - 1 \right)$$

or

$$\frac{dR}{\left( \frac{\beta R_r}{\beta - 1} - \left( \frac{\alpha}{\beta - 1} + 1 \right) R \right)} = \frac{dN}{N} \quad (7A)$$

Take integration on both sides

$$\frac{1}{-\left( \frac{\alpha}{\beta - 1} + 1 \right)} \int \frac{d \left\{ \frac{\beta R_r}{\beta - 1} - \left( \frac{\alpha}{\beta - 1} + 1 \right) R \right\}}{\left\{ \frac{\beta R_r}{\beta - 1} - \left( \frac{\alpha}{\beta - 1} + 1 \right) R \right\}} = \int \frac{dN}{N} \quad (8A)$$

Condition for integrating this equation is

$$\frac{\beta R_r}{\beta - 1} \neq \left( \frac{\alpha}{\beta - 1} + 1 \right) R, \text{ it gives } \frac{dN_r}{dN_e} \neq \left( \frac{\alpha - 1}{\frac{R_r}{R} - 1} \right) \quad (9A)$$

At  $\beta = 1$ , this equation is not solvable, hence first we take the case where  $\beta \neq 1$ .

Case I: On integrating eq. (8A), we get

$$-\frac{1}{\left( \frac{\alpha}{\beta - 1} + 1 \right)} \ln \left\{ \frac{\beta R_r}{\beta - 1} - \left( \frac{\alpha}{\beta - 1} + 1 \right) R \right\} = \ln N + k \quad (10A)$$

After applying boundary conditions, at  $t = 0$ ,  $N = N_0$ ,  $R = R_0$ , we evaluate  $k$  and substitute in eq. (10A); we get

$$-\frac{1}{\left( \frac{\alpha}{\beta - 1} + 1 \right)} \ln \left\{ \frac{\frac{\beta R_r}{\beta - 1} - \left( \frac{\alpha}{\beta - 1} + 1 \right) R}{\frac{\beta R_r}{\beta - 1} - \left( \frac{\alpha}{\beta - 1} + 1 \right) R_0} \right\} = \ln \frac{N}{N_0} = \ln f \quad (11A)$$

Further simplifying eq. (11A) becomes

$$R = R_0 f^{-\left( \frac{\alpha}{\beta - 1} + 1 \right)} + \frac{\beta}{\alpha + \beta - 1} R_r \left[ 1 - f^{-\left( \frac{\alpha}{\beta - 1} + 1 \right)} \right] \quad (12A)$$

From the standard definition of  $\delta$ , we write  $R = (1 + \delta \times 10^{-3})R_0$ , we get the final equation

$$\delta = \frac{\beta \delta_r - \varepsilon}{\alpha + \beta - 1} \left( 1 - f^{-\left( \frac{\alpha}{\beta - 1} + 1 \right)} \right) \quad (13A)$$

Substituting  $\beta = 0$ , eq. (13A) converts to the simple Rayleigh equation, which cross-checks the derivation.

Case II: When  $\beta = 1$ ,  $dN = 0$  hence from eq. (5A),  $dN_r = dN_e$ . Equation (3A) becomes

$$\begin{aligned} dR &= \frac{dN_r^* - dN_e^*}{N} = R_r \frac{dN_r}{N} - \alpha R \frac{dN_e}{N} \\ &= \left( \frac{R_r - \alpha R}{N} \right) dN_r = \left( \frac{R_r - \alpha R}{N_0} \right) dN_r \end{aligned} \quad (14A)$$

After rearranging and integrating, this becomes

$$-\frac{1}{\alpha} \int \frac{d(R_r - \alpha R)}{R_r - \alpha R} = \frac{1}{N_0} \int dN_r \quad (15A)$$

After solving fully and applying boundary conditions as in the previous case, we get

$$\delta = \frac{(\delta_r - \varepsilon)}{\alpha} \left( 1 - e^{-\alpha \frac{N_r}{N_0}} \right) \quad (16A)$$

Equations (13A) and (16A) express the resultant isotopic composition after addition and removal of isotopic molecules for  $\beta \neq 1$  (when input and output fluxes are different) and  $\beta = 1$  (when these fluxes are the same), respectively.

1. Dansgaard, W., Stable isotopes in precipitation. *Tellus*, 1964, **16**, 436-468.
2. Lord Rayleigh, On the distillation of binary mixtures. *Phil. Mag.* 1902, **4**, 521-537.
3. Criss, R. E., *Principles of Stable Isotope Distribution*, Oxford University Press, Oxford, UK, 1999, p. 254.
4. Mook, W. G., *Introduction to Isotope Hydrology: Stable and Radioactive Isotopes of Hydrogen, Oxygen and Carbon*, Taylor & Francis Group, London, Great Britain, 2006.
5. Sharp, Z., *Principles of Stable Isotope Geochemistry*, Pearson Prentice Hall, New Jersey, USA, 2007, p. 344.
6. Clark, I. D. and Fritz, P., *Environmental Isotopes in Hydrogeology*, Lewis Publisher, 1997, p. 328.
7. Faure, G., *Principles of Isotope Geology*, John Wiley, Singapore, 1986.
8. Gat, J. R. and Gouffanti, R., *Stable Isotope Hydrology, Deuterium and Oxygen-18 in the Water Cycle*, Technical Report Series No. 210, IAEA, Vienna, 1981, pp. 103-142.
9. Broecker, W. S. and Oversby, V. M., *Chemical Equilibria in the Earth*, McGraw-Hill, New York, London, 1971.
10. Hoefs, J., *Stable Isotope Geochemistry*, Springer-Verlag, Berlin, 1980, 2nd edn.

11. Ray, J. S. and Ramesh, R., Rayleigh fractionation of stable isotopes from a multicomponent source. *Geochim. Cosmochim. Acta*, 2000, **64**, 299–306.
12. Albarede, F., *Introduction to Geochemical Modeling*, Cambridge University Press, UK, 1995, pp. 47–50.
13. Richards, F. A., Anoxic basins and fjords. In *Chemical Oceanography* (eds Riley, J. P. and Skirrow, G.), Academic Press, 1965, vol. 1, pp. 611–645.
14. Payne, W. J., Reduction of nitrogenous oxide by micro-organisms. *Bacteriol. Rev.*, 1973, **37**, 409–452.
15. Cline, J. D. and Richards, F. A., Oxygen deficient conditions and nitrate reduction in the eastern tropical North Pacific Ocean. *Limnol. Oceanogr.*, 1972, **17**, 885–900.
16. Naqvi, S. W. A., Some aspects of the oxygen deficient conditions and denitrification in the Arabian Sea. *J. Mar. Res.*, 1987, **45**, 1049–1072.
17. Brandes, J. A., Devol, A. H., Yoshinari, T., Jayakumar, D. A. and Naqvi, S. W. A., Isotopic composition of nitrate in the central Arabian Sea and eastern tropical North Pacific: A tracer for mixing and nitrogen cycles. *Limnol. Oceanogr.*, 1998, **43**, 1680–1689.
18. Delwiche, C. C. and Steyn, P. L., Nitrogen isotope fractionation in soils and microbial reactions. *Environ. Sci. Technol.*, 1970, **4**, 929–935.
19. Mariotti, A., Germon, J. C., Hubert, P., Kaiser, P., Letolle, R., Tardieux, A. and Tardieux, P., Experimental determination of nitrogen kinetic isotope fractionation: some principles; illustration for the denitrification and nitrification processes. *Plant Soil*, 1981, **62**, 413–430.
20. Barford, C. C., Montoya, J. P., Altabet, M. A. and Mitchell, R., Steady-state nitrogen isotope effects of N<sub>2</sub> and N<sub>2</sub>O production in *Paracoccus denitrificans*. *Appl. Environ. Microbiol.*, 1999, **65**, 989–994.
21. Blair, N., Leu, A., Munos, E., Olsen, J., Kwong, E. and Des Marais, D., Carbon isotopic fractionation in heterotrophic microbial metabolism. *Appl. Environ. Biol.*, 1985, **50**, 996–1001.
22. Cleixner, C., Danier, H. J., Werner, R. A. and Schmidt, H. L., Correlations between the <sup>13</sup>C content of primary and secondary plant products in different cell compartments and that in decomposing basidiomycetes. *Plant Physiol.*, 1993, **102**, 1287–1290.
23. Accoe, F., Boeckx, P., Cleemput, O. V., Hofman, G., Zhang, Y., Li, R. H. and Guanxiong, C., Evolution of the  $\delta^{13}\text{C}$  signature related to total carbon contents and carbon decomposition rate constants in a soil profile under grassland. *Rapid Commun. Mass Spectr.*, 2002, **16**, 2184–2189.

ACKNOWLEDGEMENT. We are grateful to the anonymous reviewers for their critical comments.

Received 10 July 2009; revised accepted 7 January 2010

## Thermal conductivity estimates in the Niger Delta using lithologic data and geophysical well logs

Idara Akpabio<sup>1,\*</sup> and Joe Ejedawe

Exploration Department, Shell Petroleum Development Company, Port Harcourt, Nigeria

<sup>1</sup>Permanent address: Department of Physics, University of Uyo, Nigeria

**Thermal rock properties and heat flow were determined from 260 wells in the Niger Delta. The thermal conductivity data provides inputs for the determination of heat flow and for thermal evaluation of the Niger Delta basin. A map has been constructed using lithologic data and geophysical well logs to give an overview of its distribution. The thermal conductivity for sand and shale, the predominant lithology in the Niger Delta, shows wide variations from one well to another. In the Benin Formation, thermal conductivity has an average value of 8 W/mK. The lowest values are found offshore westward, while highest values occur northward. The conductivity values, however, decrease towards the marine paralic section, with an average value of 5 W/mK, the region of highest interest. The thermal conductivity values have been used in calculating heat flow. A significant regional trend of relatively low (20–30 mW/m<sup>2</sup>) heat flow at the central part of the delta, increases both seaward and northward (40–55 mW/m<sup>2</sup>). The lowest values of heat flow as low as 20 mW/m<sup>2</sup> are recorded in the central part of the delta while the highest values exceeding 50 mW/m<sup>2</sup> are recorded in the northern part of the delta. Knowledge of thermal properties has direct relevance for hydrocarbon exploration. It has been established that the bulk of hydrocarbon accumulation in the Niger Delta is of thermal origin, hence the importance of this findings.**

**Keywords:** Heat flow, temperature, thermal conductivity, sand percentage.

THE thermal conductivity of rocks is one of the major factors that affect temperature in sedimentary basins and therefore, should be addressed in basin analysis; its effect on the temperature distribution is significant, up to the order of 50–80%<sup>1</sup>. As a result of thermal conductivity, thermal structure of a basin may change laterally and vertically even if the heat flow into the basin is regionally the same<sup>2</sup>.

The variability of heat flow in most basins must arise from some combination of at least the following four principal influences: heat redistribution by migration of formation fluids (hydrodynamic effect); variations in conductivity and heat generation in the sedimentary

\*For correspondence. (e-mail: idara\_akpabio@yahoo.com)