

frequency voltage. As the high-frequency voltage gradually increases, the anode current also gradually increases until a stage comes when there is no further change in the current with the increase of high-frequency voltage. The nature of variation is shown in the graph given in Fig. 2.




FIG. 2

Variation of Thermionic current with H. F. Voltage

Thus it seems likely that there is no dependence of dielectric constant on the high-frequency voltage, but the effect observed is of a secondary nature. The high-frequency voltage alters the magnitude of the anode current, but in our experiments and those of previous workers, the anode current was maintained constant by altering only the filament resistance. This procedure certainly altered the effective value of the electronic concentration between the condenser plates. This fact is further strengthened by the observation that the thermionic current begins to assume constant value at that high-frequency voltage where the change of capacity tends to become constant (compare Fig. 1 and Fig. 2).




FIG. 3

Change in Thermionic current (in  $5 \times 10^{-6}$  amp.) due to application of H. F. Voltage

To bring out the relationship more clearly Fig. 3 has been drawn between the change of thermionic current due to impressed High Frequency Voltage and the change of capacity observed. The figure reveals that at least partially the observed change in capacity is due to change in electronic concentration consequent upon the introduction of High Frequency Voltage.

In conclusion the author thanks the Government of Bihar for the kind award of a scholarship during the tenure of which this piece of work was undertaken. The author is also thankful to Prof. S. P. Prasad and Dr. B. N. Singh for constant help and guidance throughout the investigation.

Physics Laboratory,  
Science College, Patna,  
January 1, 1943.

SNEHMOY GHOSH.

1. Prasad and Verma, *Zeits. f. Physik.*, Band 99, 7 and 8 heft, 107. 2. Khastgir and Choudhury, *Indian Journal of Physics*, 1940, 14.

### THE RARE OCCURRENCE OF MELILITE-DIOPSIDE-NEPHELINE ASSOCIATION IN A CALCIPHYRE, NEAR NANJANGUD, MYSORE

In my present detailed studies of the Charnockite rocks in Mysore, while re-examining the micro-sections of some specimens of re-crystallised calc-granulites (Caliphyres) I had collected near Nanjangud in 1923, I recently noticed in one of them a mineral—which, from its optical characters and micro-chemical tests—I recognise as Melilite. A careful examination of the several specimens I had previously collected in the area discloses that Melilites, of varying optical characters, occur in a narrow band of garnetiferous-hornblende diopside granulite,—a component of a thin composite series of calc-granulites which are injected and veined by a later set of pegmatite. Melilites—presumably of varying composition—where found, are intimately associated with diopside, or zoisite and epidote; and are seen either as separate stout laths or in intergrowth with the one or the other of its associated minerals. One of the specimens of these granulites discloses in addition, large plates of nepheline enclosing, poikilitically, several coarse grains of diopside and granular sphene. Scapolite, lime, alumina, garnet, sphene, epidote, and some unidentified rare types of lime silicate minerals are also found in some of the specimens of highly calciferous types.

The occurrence of Melilite and Nepheline in these Calciphyres, is not only of great interest as the first recorded discovery of these minerals in Mysore, but their close association with diopside would throw some light on their paragenesis and on the mode of origin of the melilite rocks which, at least so far as this area is concerned, seem to have been formed from re-actions between an older impure dolomitic limestone and the later injected alkalic liquids connected with the granitic intrusions of the region. A full description of these melilite-bearing rocks, which are still under a detailed study, will be given in Volume XLI, *Records of the Mysore Geological Department*, which will be published in the course of a few months.

Mysore Geological Dept.,  
Bangalore,  
January 27, 1943.

B. RAMA RAO.