

cemented sides, the gritty conglomeratic stuff in the grooves suggesting as if the drains are filled with cement concrete (see the photo). The senior author was struck by its resemblance to the surface structures observed at places along the great Moine thrust plane in Scotland.



It may be mentioned in this connection that there runs a great thrust-fault along the foot of this spur which has brought the Kamlials in bottom of the valley against the Purple Sandstone of the spur, and that the Purple Sandstone is overlain by the Neobolus shales and other stratigraphical formations characteristic of the eastern part of the Salt Range.

According to Dr. Gee, pebbly bed at the top of the Purple Sandstone series is widespread and is a normal horizon of the series. We met Dr. Gee at Khewra at that time; and during two of his excursions, he showed us this pebbly bed in the Dandot gorge as well as in the Khewra gorge. The typical thrust plane structure seen by us at the above locality can, therefore, be only regarded as local. The Salt Range is now politically out of bounds to Indian geologists, but the Pakistan geologists can investigate this thrust plane in detail.

Dhanbad,  
September 17, 1948.

N. L. SHARMA.  
K. C. LAHIRI.

#### THE HOME OF MONAZITE IN THE VIZAGAPATAM AREA

In a recent paper,<sup>1</sup> the occurrence of monazite, zircon and other minerals in the beach sands of Vizagapatam was described. Subsequently, on a study of the origin of the Waltair Highlands by the authors of this note (under publication), the monazites were traced to the alluvium in the streams flowing down the hill ranges of this area. In order to fix definitely the home of this monazite, a very careful examination of the pegmatite veins which cut through the khondalites underlying the red loam of the Waltair Highlands was made. These pegmatites consist of grey, pink, and white feldspars, white and bluish quartz, small books of biotite mica, opaque iron ore minerals, and some dark minerals with greenish tinge and sub metallic lustre. The pegmatites occur as

lit-par-lit injections, veins, or sometimes as lenses and veinlets in the khondalites. Representative samples from four different pegmatites were pulverised and passed through 60-mesh sieve, and panned. In all the samples, the tailings show considerable quantities of monazite and zircon. The monazite is easily distinguished in the concentrates by its greenish yellow colour. The dark mineral with submetallic lustre occurring in the pegmatites was tested by physical and optical methods and identified as monazite.

Though granites occur in the vicinity of the khondalites as intrusives, the pegmatites do not show intrusive relationship with the granites. It is of interest to recall that Masillamani and Chacko<sup>2</sup> and also Masillamani<sup>3</sup> recorded the occurrence of monazite in pegmatites which are intrusive into the khondalites and charnockites in some areas in Travancore. Tipper<sup>4</sup> is of the opinion that the bulk of the monazite in the shore sands is derived from the gneisses of Travancore. Detailed investigations are in progress extending the enquiry to other areas and formations. The results of the investigations carried out so far show that in the Vizagapatam area, the pegmatites cutting through the khondalites carry the bulk of the monazite.

Department of Geology,  
Andhra University,  
Waltair,

C. MAHADEVAN.  
N. SATHAPATHI.

September 28, 1948.

1. Mahadevan, C., and Sreeramadas, A., "Monazite in the beach sands of Vizagapatam District." *Proceedings of the Indian Academy of Sciences*, 1948, 27, No. 4, Sec. A, 275-78. 2. Chacko, I. C., and Masillamani, E., "Report of the State Geologists, Travancore," 1907-1910 p. 7. 1910. 3. Masillamani, E., "A preliminary report on the geology of Eraniel, etc., Travancore," 1911, p. 6. 4. Tipper G. H., "The monazite sands of Travancore," *Rec. G.S.I.*, 1914, 44, 94.

#### INFLUENCE OF DIETARY FACTORS ON THE ENDOGENOUS CALCIUM EXCRETION IN THE ALBINO RAT

EVEN on a calcium-free diet, there is a small but distinct negative calcium balance as evidenced by the results of Jones.<sup>1</sup> The measurement of this endogenous calcium excretion is necessary while studying the availability of calcium. It is known that protein, fat and phosphorus in the diet influence the absorption and utilisation of calcium.<sup>2, 3, 4</sup> The effect of these dietary constituents on the endogenous calcium excretion in the white rat forms the subject-matter of this note. These data would be of value in studying the physiological interrelationships that exist in the metabolism of calcium on the one hand and these dietary factors on the other.

Six healthy adult rats were taken and their normal endogenous calcium excretion was measured for a period of one week. During this period they were fed on a diet adequate in all respects but free from calcium.\* After keeping them on a normal stock diet for one week they were given three other diets which apart from