

the half-arc angle for the same value of  $OH$ . The true explanation is, however, not so simple; for, the situation is complicated by the fact that perception of depth is to a great extent subjective.<sup>3</sup> The thicker and darker a cloud, the more convex does the sky look to the eye. The influence of this subjective impression so far outweighs the geometrical fall of the half-arc angle due to decrease in the height of the cloud-base during day-time that the net effect is one of its actual increase. Such an explanation for this anomaly is justified because, during day-time, the value of  $OH$  is unaffected by the elevation of the cloud-base and is mainly determined by the existing conditions of visibility. During night-time, on the other hand, when the skies are overcast, due to the prevalence of darkness all round, the apparent value of  $OH$  would vary subjectively with the illumination that obtains on the furthest fringes of the cloud canopy. The value of  $OZ$  would similarly depend upon the zenithal luminosity of the overcast. With increasing cloud-height on moon-lit nights,  $OH$  and  $OZ$  would, therefore, vary both objectively and subjectively.

The thicker the cloud during the moon-lit night, the darker it looks both at the zenith as well as near the horizon. This would make the distances of  $OH$  as well as  $OZ$  appear equally enhanced. The influence of subjective impression on  $OH$  may thus almost neutralise that on  $OZ$ . The variation of the half-arc angle with elevation of cloud-base may, therefore, become effectively objective.

A matter of practical interest that emerges from this study is the usefulness of the results reported here for the determination of the height of base of cloud during moon-lit nights. When the sky is overcast during the night, it is difficult for an observer on the ground, particularly in poor moon-light, to distinguish between different types of clouds and much more so to estimate their heights of base. Experience shows that searchlight observations are not quite possible in the presence of moon-light, as the beam of light fails to sufficiently illuminate the base of the cloud to enable observation from the ground, except when the cloud base is very low and the moon is very feeble. A mere measurement of the half-arc angle would, on such occasions, enable a fairly accurate judgment of the cloud-base. If a graph be drawn between the half-arc angle and the height of cloud-base using the data reported here, the height corresponding to any angle can be easily read off therefrom. Such a graph shows that the relationship between the half-arc angle and the height of the cloud-base is almost linear up to about 15,000 feet.

The usefulness of the above graph is not limited to the overcast full-moon skies only. For some of the measurements made by the author, during the course of his work on the effect of illumination on the apparent shape of the sky, show that the half-arc angle for any overcast sky, when the moon is half is practically the same as when the moon is full. The graph can, therefore, be used for all states of the moon from half to full. Nevertheless, when the sky is thickly overcast with a deep layer of low cloud, the moon-light may so

completely be cut off as to render the clouding imperceptible. On such occasions it may not be possible to determine the half-arc angle. But under such conditions, cloud-height can be easily measured by the searchlight method.

The author wishes to thank Mr. B. N. Sreenivasiah, Meteorologist, Madras, for his kind interest in the work.

Meteorological Office,  
St. Thomas' Mount,  
Madras,

D. VENKATESWARA RAO.

November 17, 1947.

1. Venkateswara Rao, D., *Curr. Sci.*, 1947, **16**, 55.
2. —, *Ibid*, 1946, **15**, 40. 3. —, *Proc. Ind. Acad. Sci.*, 1947, **25**, 34. 4. Miller, A., and Neuberger, H., *Bull. Amer. Met. Soc.*, 1945, **26**, 212.

### A PROPOSED NEW COMPONENT OF SOLAR RADIATION

IN a previous communication<sup>1</sup> the author examined variations in the selective absorption of light by molecules of the earth's lower atmosphere in the region of the oxygen absorption bands, and suggested that these were due, in part, to action on the molecules of a high-powered component within the earth's fair-weather electric field.<sup>2</sup> A comparative study of fundamental atmospheric electric phenomena has, however, indicated an additional, possible contributory factor.

In the accompanying comparison diagram are reproduced, with Greenwich mean time as abscissa and a variable ordinate (for which reference should be made to the original papers) the following curves:

(a) Curve 'e' in Fig. 2 of Price and Chapman's<sup>3</sup> estimate of diurnal variations in the non-vanishing, terrestrial magnetic, line-integral current flowing across a representative region of the earth's surface;

(b) Curve 7 in Fig. 3 of Hogg's<sup>4</sup> variations in the rate of production of small ions in the lower atmosphere in fair weather;

(c) the mean of the electrogram and curve shown in Figs. 3 and 4 respectively of the diurnal variations in the apparent vertical component of the earth current observed by Forbush<sup>5</sup>;


(d) curve of variations in the potential gradient of the earth's electric field from a typical fair-weather electrogram of Colaba Observatory, set to G.M.T.;

(e) curve of Medi's<sup>6</sup> observations of day-time variations in atmospheric radiation, placed adjacent to Curve 1 in Fig. 1 of the author's own<sup>7</sup> qualitative observations of night-time variations in atmospheric radiation intensity, particularly in the region of the oxygen absorption bands (the red), the resultant indicating a marked correlation with the atmospheric potential gradient; and

(f) a curve constructed from Perot's<sup>8</sup> remarks on daily variations in the frequency of lines in the "B" band of the atmospheric oxygen absorption spectrum.

It will be seen that in every case there is, broadly speaking, a primary minimum in intensity at about noon, and a primary maximum at about 17.00 hours, both by G.M.T. The only

exception is the minimum, in the ionisation curve (b) of the diagram, which appears earlier in the day. But it must be said here that the Nolans<sup>9</sup> in their estimates of atmospheric ionisation do indeed indicate a primary minimum at noon (see Curve 3 in Fig. 3 of



Comparison diagram accompanying note entitled A proposed New Solar Radiation by A. B. Arlick, Bombay, 20th November 1947.

their paper), besides the main maximum at about 17.00 G.M.T.

Since there is reason to believe that the above widely separated geophysical phenomena are truly representative of local terrestrial conditions, such a parallelism in them may mean that they all have a common origin.

The curve (d) also being a manifestation of the earth-to-air supply current, or compensation current, the suggestion of Bauer and others,<sup>10</sup> to the effect that there exists a non-

ionizing but highly penetrating corpuscular radiation of extra-terrestrial origin which supplies this compensation current, may be adopted.

On a qualitative survey, it seems possible that a radiation of this kind, in penetrating the lower atmosphere, the earth's surface and upper crust, would cause (a) the magnetic line-integral currents by electric convection, (b) variations by irradiation or nuclear impact in the  $\gamma$ -ray activity of the atmospheric radon molecule which is the primary cause of ion-formation in the lower atmosphere, (c) vertical earth currents of the appropriate sign on penetrating the earth's surface, also by electrical convection, (d) changes in the potential gradient of the earth's electric field by progressive replenishment of the earth's escaping negative surface charge, (e) variation, by collision processes, in emissive and absorptive atmospheric radiation from the ordinary air molecules, and (f) variations, by excitation, in the frequency of lines in the absorption bands of atmospheric oxygen during night. It is, therefore, likely that this radiation provides the common origin and causes the fundamental, fair-weather atmospheric electric phenomena noted above.

Further, since it is well known that the major phenomenon—the earth's potential gradient—varies according to solar diurnal time, all the other phenomena can be expected to vary likewise. It is, therefore, clear that the suggested new extra-terrestrial radiation, if it exists, should, directly or indirectly, have its origin on the sun.<sup>11</sup>

While work must first be done on an atmospheric molecule by a strong, applied electrostatic field component (to excite or modify the spectral radiation which it emits or absorbs), according to the electromagnetic theory it would also seem permissible to assume action by collision with particles of a proposed, new radiation of solar origin, in order to account fully for anomalies in the spectrum of the earth's lower atmosphere.

It is hoped to undertake a detailed mathematical analysis of this theory elsewhere, and I wish to express indebtedness to Professor N. R. Tawde for his encouragement and continued interest in this field of geophysics.

Department of Physics,  
Royal Institute of Science, ALFRED B. ARLICK.  
Mayo Road, Bombay,  
November 20, 1947.

1. Arlick, A. B., *Curr. Sci.*, 1946, **9**, 105.
2. Frenkel, J., *J. Phys. U.S.S.R.*, 1944, **8**, 285.
3. Price, A. T. and Chapman, S., *Proc. Roy. Soc.*, 1928, **119 A**, 188.
4. Hogg, A. R., *Beitr. z. Geoph.*, 1935, **43**, 373.
5. Forbush, S. E., *Terr. Mag.*, 1933, **38**, 4.
6. Medi, E., *Ricer. Sci.*, 1938, **2**, 139.
7. Arlick, A. B., *Sci. and Cult.*, 1939, **5**, 62.
8. Perot, A., *Compt. Rend.*, 1915, **160**, 549.
9. Nolan, P. J., and J. J., *Proc. Roy. Irish Acad.*, 1931, **60**, 30.
10. Schweidler, E., *SitzB. Akad. Wiss. Wien*, 1946, **149**, 2a, 133-44.
11. Alven, H., *Phys. Rev.* 1947, **72**, 89.