

Opacity as a General Measure of Coagulation.

IN the course of a work to be published shortly in the *Indian Chemical Journal*, it has been observed that the opacity of manganese dioxide sol increased 'zonally,' that is, with marked discontinuities, during coagulation, and that this feature became more pronounced the slower the coagulation. The curves in Fig. 1 represent but three typical cases observed during subsequent work on the




Fig. 1.

kinetics of the same sol, but using a higher colloid content and with solutions of mercurous sulphate as coagulants. These results not only confirm the 'zonal' change of opacity during coagulation, but show, contrary to expectation from the current theories and general experience, that the opacity has *decreased* during the change. As is usual in these kinetic studies, measurement of the opacity during coagulation was discontinued, as soon as the coagulating system became heterogeneous by flocculation, that is, produced discrete particles of the coagulum sensibly subject to local variations. Curves *a* and *b* refer to experiments made with the white light from a glowing filament; these show that the opacity diminishes with coagulation. Partly different results were noticed when a narrow band, almost monochromatic near H α , was

employed. Curve *c* is one of the results. It shows that there is an initial diminution of opacity followed by a rise, and that both the stages of the change are marked by the 'zonal effect'. It has been almost a tacit assumption with colloid chemists that coagulation entails an increase of opacity; the adoption of the last quantity as a measure of the degree of coagulation has had a wide and long usage in the field of coagulation kinetics. It is of considerable interest, therefore, to observe, it would appear for the first time in this line, not only a limitation of the general validity of the above assumption, but of the possibly general discontinuous character of the change, independent of whether opacity and coagulation vary in the same sense or otherwise, as was also noticed previously in respect of changes of viscosity and refractive index, during coagulation.¹

S. S. JOSHI.

A. PURUSHOTHAM.

Department of Chemistry,
Benares Hindu University,
Benares,
May 14, 1936.

¹ Joshi and Rao, *Curr. Sci.*, 1936, 4, 481; also *J. Indian Chem. Soc.*, 1936, 13, 141.

Fractionation of Starch.

A SURVEY of the literature on amylolysis in relation to the structure of starch reveals the existence of two contradictory views supporting either the homo- or heteromolecular nature of starch. A study of the mutarotation phenomena accompanying the hydrolysis of starch with different amylases led Kuhn¹ to the view that the starch molecule consisted of both α - and β -glycosidic linkages which were specifically attacked by α - and β -amylases; the sense of the mutarotation was independent of the substrate but was characteristic of the enzyme. van Klinkenberg² from his work on the action of integrally pure β -amylase on starch, which hydrolysed a definite fraction (64 per cent.) of the starch substance, advanced the view that the liberation of the α - and β -maltose was not due to the alternative types of hydrolysis of a single substrate, but rather, to the specific hydrolysis of different components of starch which he designated α - and β -starches. A critical examination of the question by Hanes³ has revealed that, while the main conclusions of van Klinkenberg can be confirmed, no