
***Agrobacterium*-mediated transformation of chickpea with α -amylase inhibitor gene for insect resistance**

S IGNACIMUTHU* and S PRAKASH

Entomology Research Institute, Loyola College, Chennai 600 034, India

*Corresponding author (Fax, 91-44-2817 4644; Email, eri_lc@hotmail.com)

Chickpea is the world's third most important pulse crop and India produces 75% of the world's supply. Chickpea seeds are attacked by *Callosobruchus maculatus* and *C. chinensis* which cause extensive damage. The α -amylase inhibitor gene isolated from *Phaseolus vulgaris* seeds was introduced into chickpea cultivar K850 through *Agrobacterium*-mediated transformation. A total of 288 kanamycin resistant plants were regenerated. Only 0.3% of these were true transformants. Polymerase chain reaction (PCR) analysis and Southern hybridization confirmed the presence of 4.9 kb α -amylase inhibitor gene in the transformed plants. Western blot confirmed the presence of α -amylase inhibitor protein. The results of bioassay study revealed a significant reduction in the survival rate of bruchid weevil *C. maculatus* reared on transgenic chickpea seeds. All the transgenic plants exhibited a segregation ratio of 3:1.

[Ignacimuthu S and Prakash S 2006 *Agrobacterium*-mediated transformation of chickpea with α -amylase inhibitor gene for insect resistance; *J. Biosci.* 31 339–345]

1. Introduction

Chickpea is the world's third most important pulse crop and India produces 75% of the world's supply. Chickpea is a good source of carbohydrate (48.2–67.6%), protein (12.4–31.5%), starch (41–50%), fat (6%) and nutritionally important minerals (Geervani and Umadevi 1989). Young leaves, tender shoots and green pods are eaten as a green vegetable. Among the legumes chickpea is the best hypocholesteremic agent, followed by black gram and green gram (Soni *et al* 1982).

Production of chickpea has remained constantly low because of its susceptibility to several pathogens and insect pests. Among the insect pests, bruchids cause substantial loss during storage (Sing *et al* 1994). The cowpea weevil (*Callosobruchus maculatus*) and azuki bean weevil (*C. chinensis*) infest chickpea seeds heavily. Since seeds are used for consumption, use of chemicals to protect the seeds is not recommended. Hence newer approaches such as genetic engineering and molecular breeding have been encouraged (Jaiwal *et al* 2001; Dayal *et al* 2003; Grant *et al* 2003; Saini *et al* 2003).

Many insecticidal proteins and molecules of plant origin such as lectins, α -amylase inhibitor and protease inhibitor can retard insect growth and development when ingested (Boulter 1993; Ussuf *et al* 2001). When α -amylase inhibitor gene from common bean was expressed in transgenic pea, seeds became resistant to the infestation of bruchid weevils (Shade *et al* 1994; Shroeder *et al* 1995). The α -amylase inhibitor markedly suppressed the α -amylase activity in the larval midgut of the weevils (Ryan 1990; Ishimoto and Chrispeels 1996). Since α -amylase inhibitor is easily inactivated by cooking, introducing this gene into host plants can be regarded as a safe strategy.

In this study, α -amylase inhibitor gene (α AII) isolated from the seeds of *Phaseolus vulgaris* L. (common bean) has been introduced into chickpea through *Agrobacterium*-mediated transformation to analyse the ability of the gene to inhibit growth of bruchid weevil *C. maculatus*, which causes severe damage to chickpea seeds during storage. The gene has been introduced under a strong seed-specific promoter, phytohemagglutinin (PHA), which directs the expression of the gene in chickpea seeds.

Keywords. *Agrobacterium*; α -amylase inhibitor; chickpea; insect resistance.

2. Materials and methods

The plasmid pTA3 (7.5 kb) containing α -amylase inhibitor gene flanked by phytohem-agglutinin promoter and polyA sequences (a gift from T J V Higgins, CSIRO, Australia) was digested with *Hind*III restriction enzyme; the digested fragment (pAPSK α AI1, figure 1) was subcloned into the corresponding site of pCAMBIA 2301 vector (11.5 kb) and the plasmid was mobilized into *Agrobacterium* vir helper strain LBA 4404 by triparental mating (Ditta *et al* 1980) in which LBA 4404 was used as recipient and *Escherichia coli* was used as conjugal helper strain. Southern hybridization of transconjugants using DIG labelling was carried out to confirm the site of integration using *Bam*HI restriction enzyme.

Seeds of chickpea cultivar K850 were obtained from the National Germplasm Centre, Indian Agricultural Research Institute (IARI), Pusa, New Delhi. This cultivar was chosen because of its desirable characters such as high yield and short duration. The seeds were washed in running tap water several times. Then the seeds were treated with labolene detergent solution for 5 min and rinsed in double distilled water. The seeds were further sterilized with 70% ethanol for 1 min followed by a treatment with 0.1% mercuric chloride for 4 min and washed six times with sterile double distilled water. Sterilized seeds were germinated on seed germination medium [MS salts, MS iron, B₅ vitamins, MES (3mM), sucrose 3% and agar 0.8% at 5.6 pH].

Embryogenic axis explants obtained from aseptically germinated seedlings were preincubated for 72 h; they were wounded and infected with LBA 4404 (pAPSK α AI1) *Agrobacterium* strain at 0.8=OD₆₀₀ for 30 min and after removal of excess bacterium by blotting, were cocultivated for 72 h in cocultivation medium with acetosyringone, sodium thiosulphate, L-cysteine and dithiothreitol at 25°C in the dark. Cocultivated embryogenic axis explants were rinsed in liquid MS basal medium fortified with 500 mg/l

cefotaxime with a few drops of Tween 20. Then the explants were blotted on sterile paper to remove excess bacteria and moisture and transformed to shoot induction medium (MS salt + B₅ vitamins + BAP 1.0 mg/l) with kanamycin (100 mg/l) for nine weeks. After nine weeks on selection medium the shoots were transferred to regeneration medium [MS salts, MS iron, B₅ vitamins, MES (3 mM), BAP 0.1–5.0 mg/l, sucrose (3%), agar (0.8%), pH 5.6]. Randomly selected transformed shoots derived from embryonic axis were assayed for GUS activity (Jefferson 1987). The shoots of 2–3 cm length were transferred to 0.5 strength MS medium [MS salts, B₅ vitamins, IBA (1.0 mg/l) + Kanamycin (100 mg/l), sucrose (3%), agar (0.8%), pH 5.6] for root induction. The plants were hardened and transferred to controlled green-house conditions and allowed to set seeds which were used for insect bioassay tests. Segregation analysis of the transgenes was also carried out using the R₀ seeds. Total DNA from the transformed plants was extracted using the method of Roger and Bendich (1994). *Hind*III was used as the restriction enzyme. Southern hybridization using DIG labelling was done to confirm the presence of α -amylase inhibitor gene in transformed chickpea plants. Western blot was performed using α AI1 anti-serum raised in rabbit (a gift from T J V Higgins) to confirm the presence of α -amylase inhibitor protein.

The α -amylase inhibitor from the seeds of transformed and untransformed chickpea seeds was quantified following the method of Piergiovanni (1992). In a blind experiment, the first generation (F₁) of transformed and untransformed chickpea seeds were placed in separate glass jars (8 x 6 x 4 cm) and tested for insect resistance. Ten pairs of newly emerged adults, obtained from the stock culture, were introduced into glass jars containing chickpea seeds. After one week dead insects were removed and the percentage of mortality was calculated.

The F₁ adult emergence was noted after 26 days and the weights of the newly emerged adults were recorded using a




Figure 1. T-DNA map of plasmid pAPSK α AI.




Figure 3. PCR amplification of *GUS* gene from transformed chickpea embryonic axis explants resulting from infection with *Agrobacterium* LBA 4404 strain. Lane 1, Marker lambda DNA/*Hind*III digest; lanes 2-3, negative control (undigested DNA from plants); lanes 4-5, positive control (plasmid DNA); lanes 6-7, negative control (digested DNA from plants); lanes 8-15, transformed chickpea plants.




Figure 4. Southern hybridization of *Hind*III digested DNA from transgenic chickpea plants using α -amylase gene probe. Lanes 1-6, Genomic DNA from transformed chickpea plants: CP-1-10, CP-1-12, CP-2-14, CP-2-16, CP-3-18, CP-3-20; lane 7, genomic DNA from untransformed control chickpea plant.

to be significantly reduced in comparison to insects reared on untransformed seeds.

4. Discussion

A major application of gene transfer technology is the introduction of agronomically useful traits into crop plants.

Even though there is one report available on the introduction of *CryIA(c)* gene (Kar *et al* 1997) into chickpea to protect crop from feeding larvae of *Helicoverpa armigera* in field conditions, reports on the resistance to stored product pests are scanty (Sarmah *et al* 2004).

In this study α -amylase inhibitor gene has been introduced into chickpea (K850) through *Agrobacterium*-mediated transfer to provide resistance to bruchid weevil.




Figure 5. Western blot analysis of α -amylase inhibitor protein from transgenic chickpea plants. Total protein extracts from the transgenic chickpea seeds were fractionated by SDS/PAGE and blotted onto a nitrocellulose membrane and probed with polyclonal antibody to α -amylase inhibitor protein. Lane 1, Marker; lane 2, untransformed chickpea plant; lanes 3-8, transformed chickpea plants: CP-1-10, CP-1-12, CP-2-14, CP-2-16, CP-3-18, CP-3-20.

Agrobacterium-mediated transformation of chickpea has been attempted earlier (Krishnamurthy *et al* 2000; Sanyal *et al* 2003; Tewari-Singh *et al* 2004) but only one transformation work has been done for stored product insect resistance (Sarmah *et al* 2004). In the present study, a good number of chickpea plants carrying α -amylase inhibitor gene has been produced in K850 cultivar by *Agrobacterium tumefaciens* mediated transformation. This study also proves the efficiency and effectiveness of the virulent *A. tumefaciens*

LBA 4404 (pAPSK α AI1) strain in transforming chickpea embryonic axis.

The *GUS* gene expression under the control of a strong seed-specific promoter (CaMV 35S) in transformed chickpea shoots was stable. Similar results have been reported earlier by other workers (Krishnamurthy *et al* 2000; Jaiwal *et al* 2001; Saini *et al* 2003).

Molecular analysis through PCR amplification confirmed the presence of *GUS* gene in transgenic shoots. Genomic analysis of six randomly selected transgenic plants confirmed the integration of α -amylase inhibitor gene. The number of hybridization signals indicated that single copy of the T-DNA had been integrated. Similar results have been obtained by other workers in various plant species using α -amylase inhibitor gene (Schroeder *et al* 1995; Ishimoto and Chrispeels 1996; Ishimoto *et al* 1996; Morton *et al* 2000). Western blot analysis confirmed the presence of α -amylase inhibitor protein. Transgenic plants produced fertile seeds. Segregation of α -amylase inhibitor gene to next generation demonstrated that inheritance occurred in Mendelian ratio of 3:1.

In this study, transformed chickpea plants contained good amount of α -amylase inhibitor protein which confirmed the presence of α -amylase gene. The results of bioassay study

Table 1. Quantitative estimation of α -amylase inhibitor protein content in control and transformed chickpea seeds.

Control and transformed chickpea plants	Percent α -amylase inhibitor protein in the seeds
Control	0.0 ^a
CP-1-10	0.65 \pm 0.12 ^b
CP-1-12	0.68 \pm 0.24 ^b
CP-2-14	0.71 \pm 0.19 ^b
CP-2-16	0.63 \pm 0.16 ^b
CP-3-18	0.72 \pm 0.31 ^{bc}
CP-3-20	0.65 \pm 0.8 ^b

Numbers with the same alphabet do not differ significantly.

Table 2. Segregation of kanamycin resistant gene and *GUS* gene to R₁ generation of chickpea transformed with *Agrobacterium tumefaciens* LBA 4404 (pAPSK α AI1).

Transformed chickpea plants	No. of seeds tested	Kanamycin resistant				GUS+			
		Resistant	Sensitive	Ratio	χ^2 Value	Positive	Negative	Ratio	χ^2 Value
CP-1-10	56	40	16	3:1	0.39	40	16	3:1	0.38
CP-1-12	42	32	10	3:1	0.10	32	10	3:1	0.10
CP-2-14	60	43	17	3:1	0.35	43	17	3:1	0.35
CP-2-16	37	26	11	3:1	0.44	26	11	3:1	0.44
CP-3-18	45	32	13	3:1	0.36	32	13	3:1	0.36
CP-3-20	64	49	15	3:1	0.08	49	15	3:1	0.08

Table 3. Biological performance of *C. maculatus* reared on control and transformed chickpea seeds.

Control and transformed chickpea plants	Mortality (%)	No. of adult emergence	Mean F ₁ adult insect weight (mg)	Adult longevity (days)
Control	7.0 ± 1.5 ^a	324 ± 53.0 ^c	3.4 ± 0.45 ^c	146 ± 23.5 ^a
CP-1-10	75.4 ± 5.4 ^b	25 ± 3.5 ^a	2.4 ± 0.1 ^a	30 ± 5.1 ^{ab}
CP-1-12	80.7 ± 9.1 ^{bc}	19 ± 4.0 ^a	2.6 ± 0.21 ^{ab}	27 ± 4.6 ^a
CP-2-14	78.2 ± 8.6 ^b	28 ± 3.0 ^{ab}	1.8 ± 0.14 ^a	34 ± 5.2 ^b
CP-2-16	74.6 ± 6.7 ^b	21 ± 5.5 ^a	2.5 ± 0.20 ^{ab}	19 ± 3.2 ^a
CP-3-18	69.3 ± 7.2 ^b	34 ± 6.0 ^{ab}	2.4 ± 0.16 ^a	28 ± 4.4 ^{ab}
CP-3-20	66.5 ± 8.3	26 ± 4.5 ^a	2.0 ± 0.22 ^a	25 ± 4.0 ^a

Numbers with the same alphabet do not differ significantly.

revealed a significant reduction in the survival of bruchid weevil *C. maculatus* reared on transgenic chickpea seeds. Bioassay study proved that the transgenic seeds showed increased resistance to the bruchid weevil than the control seeds. Similar results have been reported earlier (Sarmah *et al* 2004).

Acknowledgements

We are greatly indebted to Prof. T J V Higgins, CSIRO, Australia, for providing the plasmid harbouring alpha amylase inhibitor gene and antibody and Prof. R Terada, National Institute for Basic Biology, Okazaki, Japan, for providing the *Agrobacterium* strains.

References

- Boulter D 1993 Insect pest control by copying nature using genetically engineered crops; *Phytochemistry* **34** 1453–1466
- Dayal S, Lavanya M, Devi P and Sharma K K 2003 An efficient protocol for shoot regeneration and genetic transformation of pigeon pea [*Cajanus cajan* (L.) Millsp.] using leaf explants; *Plant Cell Rep.* **21** 1072–1079
- Ditta G, Stanfield S, Corbin D and Helinski D R 1980 Bruchid host range cloning system for gram-negative bacterium. Construction of gene bank of *Rhizobium meliloti*; *Proc. Natl. Acad. Sci. USA* **77** 7347–7351
- Geervani P and Umadevi T 1989 Effect of maturation of nutrient composition of selected vegetable legumes; *J. Sci. Food Agric.* **46** 243–248
- Grant J E, Thomsons L M J, Joyce M D P, Dale T M and Cooper P A 2003 Influence of *Agrobacterium tumefaciens* strain on the production of transgenic peas (*Pisum sativum* L.); *Plant Cell Rep.* **12** 1207–1210
- Ishimoto M and Chrispeels M J 1996 Protective mechanism of the Mexican bean weevil against high levels of α -amylase inhibitor in the common bean; *Physiol. Plant.* **111** 393–401
- Ishimoto M, Sato T, Chrispeels M J and Kitamura K 1996 Bruchid resistance of transgenic azuki bean expressing seeds α -amylase inhibitor of the common bean; *Entomol. Exp Appl.* **79** 309–315
- Jaiwal P K, Kumari R, Ignacimuthu S, Potrykus I and Sautter C 2001 *Agrobacterium tumefaciens*-mediated genetic transformation of mungbean (*Vigna radiata* L. Wilczek) – a recalcitrant grain legume; *Plant Sci.* **161** 239–247
- Jefferson R A 1987 Assaying Chimeric genes in plants: the *Gus* gene fusion system; *Plant Mol. Biol. Rep.* **5** 387–405
- Kar S, Basu D, Das S, Ramakrishnan N A, Mukherjee P, Nayak P and Sen S K 1997 Expression of cry1A(c) gene of *Bacillus thuringiensis* in transgenic chickpea plants inhibits development of podborer (*Heliothis armigera*) larvae; *Trans. Res.* **6** 177–185
- Krishnamurthy K V, Suhasini K, Sagare A P, Meixner M, de Kathen A, Pickardt T, Schieder O and Kathen de A 2000 *Agrobacterium* mediated transformation of chickpea (*Cicer arietinum* L.) embryo axes; *Plant Cell Rep.* **19** 235–240
- Morton R L, Schroeder H E, Bate K S, Chrispeels M J, Armstrong G E and Higgins T J V 2000 Bean α -amylase inhibitor 1 in transgenic peas (*Pisum sativum*) provides complete protection from pea weevil (*Bruchus pisorum*) under field conditions; *Proc. Natl. Acad. Sci. USA* **97** 3820–3825
- Piergiorgio A R 1992 Effects of some experimental parameters on the activity of cowpea alpha amylase inhibitors; *Lebens. Wiss. Technol.* **25** 321–324
- Rogers S O and Bendich A J 1994 Extraction of total cellular DNA from plants, algae and fungi; in *Plant molecular biology* (eds) S B Gelvin and R A Schilperoort (The Netherlands: Kluwer) 2nd edition, D1, pp 1–8.
- Ryan C A 1990 Protease inhibitors in plants. Genes for improving defense against insects and pathogens; *Annu. Rev. Phytopathol.* **28** 25–45
- Saini R, Jaiwal S and Jaiwal P K 2003 Stable genetic transformation of *Vigna mungo* L. Hepper via *Agrobacterium tumefaciens*; *Plant Cell Rep.* **21** 701–705
- Sanyal I, Singh A K and Amala D V 2003 *Agrobacterium tumefaciens*-mediated transformation of chickpea (*Cicer arietinum* L.) using mature embryonic axes and cotyledonary nodes; *Indian J. Biotechnol.* **2** 524–532
- Sarmah B K, Moore A, Tate W, Molvig L, Morton R L, Rees D P, Chiaiese P, Chrispeels M J, Tabe L M and Higgins T J V 2004

- Transgenic chickpea seeds expressing high levels of a bean α -amylase inhibitor; *Mol. Breed.* **14** 73–82.
- Schroeder H E, Gollash S and Moore A 1995 Bean α -amylase inhibitor confers resistance to the pea weevil (*Bruchus pisorum*) in transgenic peas (*Pisum sativum* L.); *Plant Physiol.* **107** 1233–1239
- Shade R E, Schroeder R E, Poueyo J J, Tabe L M, Murdock L I, Higgins T J V and Chrispeels M J 1994 Transgenic pea seeds expressing the α -amylase inhibitor of the common bean are resistant to bruchid beetles; *Bio/Technology* **12** 793–796
- Singh K B, Malhotra R S, Halila H M, Knights E J and Verma M M 1994 Current status and future strategy in breeding chickpea for resistance to biotic and abiotic stresses; *Euphytica* **73** 137–149
- Soni G L, George M and Singh R 1982 Role of common Indian pulses as hypocholesterolemic agents; *Indian J. Nutr. Diebet.* **19** 184–189
- Tewari-Singh N, Sen J, Kiesecker H, Reddy V S, Jacobsen H J and Guha-Mukherjee S 2004 Use of herbicide or lysine plus threonine for non-antibiotic selection of transgenic chickpea; *Plant Cell Rep.* **22** 576–583
- Ussuf K K, Laxmi N H and Mita R 2001 Protease inhibitors: plant derived genes of insecticidal protein for developing insect resistant transgenic plants; *Curr. Sci.* **80** 847–853

MS received 7 December 2005; accepted 27 April 2006

ePublication: 4 July 2006

Corresponding editor: RENU KHANNA CHOPRA