

## Supplementary data:

### Determination of gene action for some biometrical traits in *Lens culinaris* Medik. under mid-hill conditions of northwestern Himalayas

Naresh Kumar, B. C. Sood, T. R. Sharma, R. K. Chahota and Salej Sood

*J. Genet.* **90**, 493–497

**Table 1.** Pedigree/source of genotypes used.

Genotype	Pedigree/source	Yield (q/ha)	Days to maturity
L-652	L-4145 × PL-639	8–10	160–165
L-651	L-4145 × PL-639	8–10	160–165
L-658	L-4145 × PL-639	8–10	160–165
L-635	Vipasa × PL-639	8–10	160–165
L-407	PL-406 × L-259	8–10	160–165
L-630	Vipasa × PL-639	8–10	160–165
L-617	Vipasa × PL-639	8–10	160–165
L-666	L-4145 × PL-639	8–10	160–165
L-649	L-4145 × PL-639	8–10	160–165
L-620	Vipasa × PL-639	8–10	160–165
L-354	L-4145 × L-259	8–10	160–165
L-403	PL-406 × L-259	8–10	160–165
L-412	PL-639 × L-259	8–10	160–165
L-642	L-4145 × PL-639	8–10	160–165
L-737	L-2991 × PL-406	8–10	160–165
Vipasa (T <sub>1</sub> )	Pureline selection from exotic germplasm	10–12	170–175
PL-406 (T <sub>2</sub> )	Selection of P-495	15–20	130–135