

Quantum information paradox: Real or fictitious?

ABHAS MITRA

Theoretical Astrophysics Section, Bhabha Atomic Research Centre, Mumbai 400 085,
India

E-mail: amitra@barc.gov.in

Abstract. One of the outstanding puzzles of theoretical physics is whether quantum information indeed gets lost in the case of black hole (BH) evaporation or accretion. Let us recall that quantum mechanics (QM) demands an upper limit on the acceleration of a test particle. On the other hand, it is pointed out here that, if a Schwarzschild BH exists, the acceleration of the test particle would blow up at the event horizon in violation of QM. Thus the concept of an exact BH is in contradiction with QM and quantum gravity (QG). It is also reminded that the mass of a BH actually appears as an integration constant of Einstein equations. And it has been shown that the value of this integration constant is actually zero! Thus even classically, there cannot be finite mass BHs though zero mass BH is allowed. It has been further shown that during continued gravitational collapse, radiation emanating from the contracting object gets trapped within it by the runaway gravitational field. As a consequence, the contracting body attains a quasi-static state where outward trapped radiation pressure gets balanced by inward gravitational pull and the ideal classical BH state is never formed in a finite proper time. In other words, continued gravitational collapse results in an ‘eternally collapsing object’ which is a ball of hot plasma and which is asymptotically approaching the true BH state with $M = 0$ after radiating away its entire mass energy. And if we include QM, this contraction must halt at a radius suggested by the highest QM acceleration. In any case no event horizon (EH) is ever formed and in reality, there is no quantum information paradox.

Keywords. Quantum information paradox; black hole; eternally collapsing object.

PACS Nos 03.67.-a; 04.20.Dw; 04.40.-b

1. Introduction: Upper limit on acceleration in quantum gravity

Quantum gravity imposes a fundamental unit of proper length called Planck length:

$$l_p = \sqrt{\frac{\hbar G}{c^3}} \quad (1)$$

and a fundamental unit of time called Planck time $\tau_p = l_p/c$. Accordingly, any QG should impose a maximal acceleration:

$$a_p = \frac{\text{Maximum speed}}{\text{Minimum proper time}} \sim \frac{c^2}{l_p} = \frac{c^{7/2}}{\sqrt{\hbar G}}. \quad (2)$$

Detailed considerations yield [1,2]

$$a_p = \frac{c^2}{2l_p} = \frac{c^{7/2}}{2\sqrt{\hbar G}}. \quad (3)$$

Thus QM does not allow the occurrence of infinite proper acceleration. Some preliminary formulation of QG indeed mentioned maximal acceleration [3].

1.1 Proper acceleration due to gravity

The invariant proper acceleration on a test particle lying on the surface of a spherical body of mass M and radius R_0 [4]

$$a = \sqrt{-a^i a_i} = \frac{GM}{R_0^2 \sqrt{1 - 2GM/R_0 c^2}}. \quad (4)$$

And if the body is instead a Schwarzschild black hole with $R_0 = 2GM/c^2$, one would have $a = \infty$ [4]!! Note that since a is an invariant/scalar, its value is independent of coordinates and even a free falling test particle would experience infinite acceleration. Clearly, such a situation would be in violation of the inherent concept of a natural QG upper limit of acceleration a_p . Thus no properly formulated QG should admit the existence of BHs. Hence in properly formulated QG theories, there should not be any horizon, BH, or Hawking radiation. Thus, as per QG, either there must be some physical process to stop continued gravitational collapse at a certain $R_0 = R_p$ where $a \leq a_p$ or else, in the case of Schwarzschild BHs, the mass of the BH should be zero ($M = 0$). This is due to two reasons:

- The proper time of formation of a zero mass BH is infinite: $\tau_{\text{formation}} \propto M^{-1/2} = \infty$.
- Even if mathematically, one would assume the existence of such a zero mass BH, the proper time of infall of a test particle for reaching the event horizon (EH) $\tau_{\text{infall}} \propto M^{-1/2} = \infty$.

In this way, the conflict of avoiding an infinite acceleration would be avoided.

1.2 Hint from string theories

There are some solutions in string theories which do not admit any horizon or singularity. For instance

- ‘If one applies the transformation (2.13) to time transformation in Schwarzschild, one obtains a solution in which the horizon becomes a singularity’ [5].

Note, horizon itself becomes the singularity only when mass of the BH: $M = 0$ because radius of the BH, $R_g = 2M$ ($G = c = 1$).

- ‘D1-D5 solutions have neither any horizon nor any singularity’ [5].
- ‘Straight strings have no event horizon’ [5].

Some solutions of the string theories specifically suggest that point particle are massless:

- ‘The consistency of our picture requires that there is one and only one supermultiplet which becomes massless at the singularity’ [6].

“At first sight it may seem surprising that classical black holes can be massless. However, this phenomenon has an appealing explanation from 10D perspective. The IIA(IIB) theory has extremal black two-brane (three-brane) solutions whose mass is proportional to their area. After Calabi–Yau compactification these may wrap around minimal two-(three-) surface and appear as a 4D black hole. As the area of the surfaces around which they wrap goes to zero, the corresponding BH becomes massless...” [6].

When BH is massless, its horizon area $A = 0$, and this is a unique state with entropy

$$S = \ln 1 = 0. \tag{5}$$

Physically a massless BH with zero horizon area confines no information. There is no question of evaporation or information loss in such a case. Nor does a test particle ever reach the horizon of a massless BH, $\tau_{\text{infall}} \propto M^{-1/2} = \infty$.

In fact, there are specific indications that QG does not actually allow any Hawking radiation (even if the existence of a BH is assumed).

‘A new argument is presented confirming the point of view that a Schwarzschild BH formed during a gravitational collapse process does NOT radiate’ [7].

- ‘Hawking effect may be only a mathematical artifact because it demands singularity of the wave function at event horizon in violation of QM’ [8].

2. Does general relativity allow finite mass BHs?

The existence of the vacuum Schwarzschild solution apparently suggests the existence of BHs of arbitrary mass. But what is forgotten here is that

(i) The ‘mass’ of the BH appears through an integration constant: $\alpha = 2GM/c^2$ or $M = c^2\alpha/2G$.

(ii) Although symbolically ‘integration constants’ may look to assume an arbitrary value, in reality, they have a definite or precise value.

2.1 Fixing the value of this integration constant

One may consider the exterior Schwarzschild BH solution in two different coordinates:

- The original Schwarzschild solution:

$$ds^2 = (1 - \alpha/R)dT^2 - (1 - \alpha/R)^{-1}dR^2 - R^2(d\theta^2 + \sin^2\theta d\phi^2). \tag{6}$$

- The Eddington–Finkelstein solution [9–12]:

$$ds^2 = (1 - \alpha/R)dT_*^2 \pm (2\alpha/R)dT_* dR - (1 + \alpha/R)dR^2 - R^2(d\theta^2 + \sin^2\theta d\phi^2) \tag{7}$$

where

$$dT_* = dT + \frac{\alpha}{R - \alpha} dR \quad (8)$$

and apply the condition that proper 4-volume remains invariant for any coordinate transformation, i.e.,

$$\sqrt{-g_*} dR dT_* d\phi d\theta = \sqrt{-g} dR dT d\theta d\phi, \quad (9)$$

where $g = \det |g_{ik}|$ and happens to be same in both the cases: $g = g_* = -R^4 \sin^2 \theta$. Then the foregoing equation yields $dT_* = dT$ which, by virtue of eq. (8) demands

$$\alpha = \frac{2GM}{c^2} = 0 \quad (10)$$

or $M = 0$. This means that though the mass of a Schwarzschild BH appears to be arbitrary, actually, it is unique: $M \equiv 0!$ It is important to note that if instead of a ‘point particle’ we would be considering a finite static spherical body of gravitational mass M , the above derivation would be invalid even though by Birchoff’s theorem, the exterior space-time would still be described by metric (2). This happens because the coordinate transformation (8) is obtained by integrating the vacuum null geodesic all the way from $R = 0$, and this is allowed only when the space-time is indeed due to a point mass and not due to a finite body. For a finite body, the interior metric would be different from what is indicated by eqs (6) and (7). As explained above, $M = 0$ BHs are never formed, and even if they would be assumed to be there, there is no Hawking radiation!

More importantly, this means that the observed BH candidates having finite masses cannot be exact BHs though they may resemble dark/black-like theoretical BHs. This would be so if the gravitational redshift of the real BH candidates, $z \gg 1$.

Why? This is so because the observed luminosity of an object falls off as

$$L_{\text{observed}} \sim (1 + z)^{-2} \rightarrow 0 \quad \text{as } z \gg 1, \quad (11)$$

where one defines

$$z = (1 - 2GM/Rc^2)^{-1/2} - 1. \quad (12)$$

Recall here that for ideal BHs, $z = \infty$ and consequently $L_{\text{observed}} = 0$.

3. Consistency with gravitational collapse

The metric of a spherically evolving fluid is given by

$$ds^2 = g_{00}dt^2 + g_{rr}dr^2 - R^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (13)$$

where $R = R(r, t)$ is the invariant circumference coordinate (also called as area coordinate) which happens to be a scalar. Further, for radial motion with $d\theta = d\phi = 0$, the metric becomes

$$ds^2 = g_{00} dt^2(1 - x^2), \quad (14)$$

where

$$x = \frac{\sqrt{-g_{rr}} dr}{\sqrt{g_{00}} dt}. \quad (15)$$

We may recast eq. (14) as

$$(1 - x^2) = \frac{1}{g_{00}} \frac{ds^2}{dt^2}. \quad (16)$$

The co-moving observer at $r = r$ is free to do measurements of not only the fluid element at $r = r$ but also of other objects: If the co-moving observer is compared with a static floating boat in a flowing river, the boat can monitor the motion of other boats or the pebbles fixed on the river bed. Here the fixed markers on the river bed are like the background $R = \text{fixed}$ markers against which the river flows. Alternatively, the co-moving observer may be seen as the driver of a car (local fluid) while the fixed milestones on the roadside constitute the $R = \text{fixed} = \text{scalar grid}$ naturally definable for any spherically symmetric problem. If we intend to find the parameter x for such a $R = \text{constant}$ milestone, we will have, $dR = 0$, or

$$dR(r, t) = 0 = \dot{R} dt + R' dr, \quad (17)$$

where an overdot denotes a partial derivative with respect to t and a prime denotes a partial derivative with respect to r . Therefore, for the $R = \text{constant}$ marker, we find that

$$\frac{dr}{dt} = -\frac{\dot{R}}{R'} \quad (18)$$

and the corresponding $x = x_c$ is

$$x = x_c = \frac{\sqrt{-g_{rr}} dr}{\sqrt{g_{00}} dt} = -\frac{\sqrt{-g_{rr}} \dot{R}}{\sqrt{g_{00}} R'}. \quad (19)$$

Using eq. (16), we also have

$$(1 - x_c^2) = \frac{1}{g_{00}} \frac{ds^2}{dt^2}. \quad (20)$$

Now let us define [13–15] two auxiliary parameters

$$\Gamma = \frac{R'}{\sqrt{-g_{rr}}}, \quad U = \frac{\dot{R}}{\sqrt{g_{00}}} \quad (21)$$

so that the combined eqs (18), (19) and (21) yield

$$x_c = -\frac{U}{\Gamma}, \quad U = -x_c \Gamma \quad (22)$$

As is well known, the gravitational mass of the collapsing (or expanding) fluid is defined by the equation [4,13–15]

$$\Gamma^2 = 1 + U^2 - \frac{2M(r, t)}{R}. \quad (23)$$

Then the two foregoing equations can be combined and transposed to obtain

$$\Gamma^2(1 - x_c^2) = 1 - \frac{2M(r, t)}{R}. \quad (24)$$

By using eqs (20) and (22) in the foregoing equation, we obtain

$$\frac{R'^2}{-g_{rr}g_{00}} \frac{ds^2}{dt^2} = 1 - \frac{2M(r, t)}{R}. \quad (25)$$

Recall that the determinant of the metric tensor is always negative [4,13–15]:

$$g = R^4 \sin^2 \theta g_{00} g_{rr} \leq 0 \quad (26)$$

so that we must always have

$$-g_{rr}g_{00} \geq 0. \quad (27)$$

As mentioned before, all worldlines of the material particles or observers must be time-like because the co-moving metric (or for that matter even the appropriate Schwarzschild metric) has no supposed ‘coordinate singularity’ unlike the vacuum Schwarzschild case. For the signature chosen here, this means, $ds^2 > 0$ for all observers. Then by noting eq. (27), it follows that the LHS of eq. (25) is always positive. So must be the RHS of the same equation which implies that

$$\frac{2M(r, t)}{R} < 1. \quad (28)$$

This shows in the utmost general fashion that trapped surfaces are not formed in spherical collapse or expansion [4,13–16]. Thus the crucial assumption behind singularity theorems of Hawking, Penrose, Geroch, i.e., ‘formation of a trapped surface’ is actually not realized in General Relativity. Therefore, if a fluid would indeed collapse to a singularity at $R \rightarrow 0$, one must have $M \rightarrow 0$ so that the inequality is honoured, and this is consistent with the result that for BHs, $\alpha = 2GM/Rc^2 = 0$ and BHs have unique mass $M \equiv 0$!

4. Eternally collapsing objects

Two natural questions here would be:

1. What about the upper limit of white dwarfs and neutron star masses?

Answer: These limits are intended for objects supported by cold quantum degeneracy pressure. On the other hand, there is no upper limit for hot objects supported by radiation pressure [17–20].

2. If there cannot be any finite mass BH, what are the true nature of the observed BH candidates?

Answer: It has been showed that during continued gravitational collapse, the radiation (photons/neutrinos) emanating from the collapsing object would get almost trapped by the runaway gravitational field. Finally a state would be reached when [17–21]:

$$\text{Inward pull of gravitation} = \text{Outward trapped radiation pressure} \quad (29)$$

and consequently, the collapsing object becomes a hot ball of quasi-static plasma where extreme radiation pressure balances the gravity. However, it is still a quasi-static stage and is contracting at infinitesimally slow rate to attain the exact BH state with $M = 0$. It has also been shown that this contraction process continues indefinitely and hence the BH candidates are eternally collapsing objects (ECOs) rather than true BHs!

5. Observational prediction and verification

As an astrophysical plasma contracts, the frozen in magnetic field increases as $B \propto R^{-2}$ and this is the basic reason that young neutron stars have $B \sim 10^{12-13}$ G. Further, in the extreme relativistic stages with $z \gg 1$, local value of B may additionally increase as $B \propto (1+z)$. On this basis, the present author had predicted that the observed BH candidates, i.e., ECOs, will have strong intrinsic magnetic field [4,13,20]. In contrast, true chargeless BHs have no intrinsic magnetic field. Consequently, some of the properties as well as accretion geometries around ECOs could resemble, in some way, the magnetized neutron stars rather than (true) BHs.

This prediction has been verified for both stellar mass (galactic) and extragalactic BH candidates (i.e., ECOs) [22–26]. In particular, for the extragalactic cases, microlensing observations spanning 20 years have given almost direct proof that, some of the quasars, in particular Q0957 + 561, harbour magnetized ECOS (MECOS) rather than true BHs [25,26].

6. Conclusion

Though BH solution is a correct exact solution, the integration constant defining the mass of a Schwarzschild BH has the unique value $M = 0$. This means that the exact BH solution represents only an ideal state which can be attained asymptotically during continued gravitational collapse. At any finite proper time, the collapsing object is never, an ideal, exact BH. On the other hand, in the context of classical general relativity, the actual BH candidates are likely to be (magnetized) ECOs, a ball of relativistic plasma where inward pull of gravity is balanced by outward radiation pressure. Since all meaningful quantum gravity theories must reduce to GR in the low energy regime, the apparent BH-like solutions too must be pointing only ideal limiting/asymptotic states which can never exactly occur in Nature. Since there is no exact BH, no event horizon, there is no Hawking radiation and quantum information paradox. As mentioned in the beginning, the existence of

an upper limit of a finite proper acceleration a_p directly contradicts the notion of exact BHs.

In an important paper [27] it was shown that if BHs would be assumed to be present, quantum information paradox would be even more severe than what is generally considered. However, rather than noting that, the most physical resolution to this severe problem would lie in the realization that there cannot be finite mass BHs or Hawking radiation, the authors unfortunately suggested the most unlikely and farfetched possibilities. But it is hoped that authors working in QM/QG would be adequately aware of the developments in astrophysics and general relativity and would accordingly realize that quantum information paradox is actually non-existent.

References

- [1] Arun Pati, *Euro. Phys. Lett.* **18**, 285 (1992)
- [2] Arun Pati, *Il Nuovo Cimento* **B107**, 895 (1992)
- [3] E R Caianiello *et al*, *Int. J. Mod. Phys.* **D3(2)**, 485 (1994)
- [4] A Mitra, *Found. Phys. Lett.* **15(5)**, 439 (2002)
- [5] G Horowitz, hep-th/9210119 (1992)
- [6] A Strominger, *Nucl. Phys.* **B451**, 109 (1995), hep-th/954090
- [7] V A Belinski, *Phys. Lett.* **A354**, 249 (2006), gr-qc/060713
- [8] M Rabinowitz, astro-ph/0412101 (2004)
- [9] A Mitra, *Adv. Sp. Res.* **38(12)**, 2917 (2006), astro-ph/0510162
- [10] A Mitra, *J. Math. Phys.* **50(4)**, 0242502 (2009), arXiv:0904.4754
- [11] A Mitra, physics/0504076 (2005)
- [12] A Mitra, *Proc. 11th Marcel Gross Conf. on General Relativity* (World Scientific, Singapore, 2008) p. 2276
- [13] A Mitra, *Found. Phys. Lett.* **13(6)**, 543 (2000)
- [14] A Mitra, astro-ph/0408323 (2004)
- [15] A Mitra, gr-qc/0512006 (2005)
- [16] D Leiter and S R Robertson, *Found. Phys. Lett.* **16**, 143 (2003)
- [17] A Mitra, *Mon. Not. R. Astron. Soc. Lett.* **367**, L66 (2006), gr-qc/061025
- [18] A Mitra, *Mon. Not. R. Astron. Soc.* **369**, 492 (2006), gr-qc/0603055
- [19] A Mitra, *Phys. Rev.* **D74(2)**, 02010 (2006), gr-qc/0606066
- [20] A Mitra, *New Astronomy* **12(2)**, 146 (2006), astro-ph/0608178
- [21] A Mitra and N K Glendenning, eScholarship Repository, Lawrence Berkeley National Laboratory, University of California (2006)
see, <http://repositories.cdlib.org/lbnl/LBNL-59320>.
- [22] S Robertson and D Leiter, *Astrophys. J.* **565**, 447 (2002)
- [23] S Robertson and D Leiter, *Astrophys. J.* **569**, L203 (2003)
- [24] S Robertson and D Leiter, *Mon. Not. R. Astron. Soc.* **350**, 1391 (2004)
- [25] R Schild, D Leiter and S Robertson, *Astronom. J.* **132(1)**, 420 (2006)
- [26] R Schild, D Leiter and S Robertson, *Astronom. J.* **135(3)**, 947 (2008)
- [27] Arun Pati and S L Baurstein, *Phys. Rev. Lett.* **98(8)**, 080502 (2007)