

Mango ginger (*Curcuma amada* Roxb.) – A promising spice for phytochemicals and biological activities

RS POLICEGOUDRA^{1,*}, SM ARADHYA² and L SINGH¹

¹Department of Biotechnology, Defence Research Laboratory, Tezpur 784 001, India

²Fruit and Vegetable technology Department, Central Food Technological Research Institute, Mysore 570 020, India

*Corresponding author (Fax, +91-3712-258534; Email, poligene@rediffmail.com)

Mango ginger (*Curcuma amada* Roxb.) is a unique spice having morphological resemblance with ginger but imparts a raw mango flavour. The main use of mango ginger rhizome is in the manufacture of pickles and culinary preparations. Ayurveda and Unani medicinal systems have given much importance to mango ginger as an appetizer, alexteric, antipyretic, aphrodisiac, diuretic, emollient, expectorant and laxative and to cure biliousness, itching, skin diseases, bronchitis, asthma, hiccough and inflammation due to injuries. The biological activities of mango ginger include antioxidant activity, antibacterial activity, antifungal activity, anti-inflammatory activity, platelet aggregation inhibitory activity, cytotoxicity, antiallergic activity, hypotriglyceridemic activity, brine-shrimp lethal activity, enterokinase inhibitory activity, CNS depressant and analgesic activity. The major chemical components include starch, phenolic acids, volatile oils, curcuminoids and terpenoids like difurocumenonol, amadannulen and amadaldehyde. This article brings to light the major active components present in *C. amada* along with their biological activities that may be important from the pharmacological point of view.

[Policegoudra RS, Aradhya SM and Singh L 2011 Mango ginger (*Curcuma amada* Roxb.) – A promising spice for phytochemicals and biological activities. *J. Biosci.* 36 739–748] DOI 10.1007/s12038-011-9106-1

1. Introduction

Several epidemiological studies established a link between phytochemicals and the range of biological activities that impart health benefits in human beings. Scientific research supports the biological activity of many of the phytochemicals more in their native forms. They were copiously used in Ayurveda and other traditional medicine (Moon *et al.* 2010), which dates back to Charaka Samhita (Gupta *et al.* 2010). Amongst the phytochemicals, several groups of polyphenols (anthocyanins, proanthocyanidins, flavanones, isoflavones, resveratrol and ellagic acid), non-nutrient chemical and dietary constituents are currently used in the pharmaceutical industry. The spices are considered to be the storehouse of active phytochemicals. The various spices belonging to the genus *Curcuma* are well known for their multiple uses as medicines, cosmetics, dyes, flavourings and nutraceuticals. Extensive work has been

carried out on *Curcuma longa* (turmeric) and *Zingiber officinale* (ginger), but *Curcuma amada* (mango ginger) is an untapped medicinal plant of the ginger family. This review explores the phytochemicals and their biological activities of *C. amada*.

2. Botany and taxonomy of *Curcuma amada*

Curcuma amada is a unique spice having morphological resemblance with ginger (*Zingiber officinale*) but imparts a raw mango (*Mangifera indica*) flavour. The genus name *Curcuma* was coined by Linnaeus in 1753 in his *Species Plantarum*. The word probably derives from the Arabic word 'kurkum', which means yellow colour (Salvi *et al.* 2000; Shirgurkar *et al.* 2001). *Curcuma amada* Roxb. is commonly known as mango ginger. It is a perennial, rhizomatous, aromatic herb belonging to the family Zingiberaceae. This family is composed of 70–80 species of rhizomatous annual

Keywords. Antimicrobials; antioxidants; biological activities; biochemical constituents; *Curcuma amada*; Mango ginger

Supplementary materials pertaining to this article are available on the *Journal of Biosciences* Website at <http://www.ias.ac.in/jbiosci/Sep2011/pp739-748/suppl.pdf>

or perennial herbs (Purseglove 1974; Aminul 2004). The genus originated in the Indo-Malayan region, and is widely distributed in the tropics of Asia to Africa and Australia (Sasikumar 2005). The plant grows to a height of 1 m (figure 1A). The leaves are long, oblong, lanceolate, radical, sheathed, petiolate and in tufts. Each plant bears 5 to 6 pairs of leaves. Mango ginger rhizomes (figure 1B and C) are fleshy, buff coloured, 5–10 cm long, 2–5 cm in diameter and demarcated into nodes and internodes. At the rhizome nodes scaly leaves are arranged circularly giving the appearance of growth rings with scars on the surface. The rhizomes are branched, and the branching is sympodial. The rhizomes emulate a raw mango flavour and taste pungent.

The following Sankrit shloka explains the synonyms of mango ginger:

*Darvibheda, Amragandha, Surabhidaru,
Karpura padmapatra, Surimat and Surataraka.*

“दर्वीभेदाग्रगन्धा च सुरभीदारु दारु च ।
कर्पूर पद्मपत्रा स्यात् सुरीमत् सुरतारका ॥” (भा.प्र.)
[“Dārvībhēdāmrāgandhā ca surabhīdāru dāru ca
Karpūrā padmapatrā syāt surīmat suratārakā” (Bhā.pra.)]

The taxonomical hierarchy of mango ginger is as follows:

Kingdom: Plantae

Super division: Spermatophyta

Division: Magnoliophyta

Class: Monocotyledonae

Order: Zingiberales

Family: Zingiberaceae

Genus: *Curcuma*

Species: *C. amada* Roxb.

3. Occurrence and distribution

The geographical distribution of this genus ranges from India to Thailand, Indo-China, Malaysia, Indonesia and northern Australia. *C. amada* is found the wild in parts of West Bengal, and is cultivated in Gujarat, Uttar Pradesh, Kerala, Karnataka, Tamil Nadu and the north-eastern states. They originated in the Indo-Malayan region and distributed widely in the tropics from Asia to Africa and Australia (Sasikumar 2005). Out of 10 *Curcuma* species, 2 species, *C. amada* and *C. zedoaria*, are distributed throughout India in the wild and cultivated forms; 4 species, *C. aeruginosa*, *C. brog*, *C. caesia* and *C. sylvatica*, occur in wild conditions and distributed throughout north-eastern part of India. *C. malabarica* and *C. aromatica* occur in south India, while *C. raktakanta* and *C. harita* are distributed throughout Kerala (Velayudhan *et al.* 1999).

4. Chemical constituents

Proximate and nutrient analysis of edible rhizome plays a crucial role in assessing their nutritional significance and nutraceutical quality. The mango ginger rhizome was found to be a rich source of fibres and starch (Lakshminarayana *et al.* 1963).

4.1 Volatile constituents

There are many reports on the composition of mango ginger volatile oil. The mango flavour is mainly attributed to presence of car-3-ene and *cis*-ocimene (figure 2) among the 68 volatile aroma components present in the essential oil of mango ginger rhizome (Dutt and Tayal 1941; Golap and Bandyopadhyaya 1984; Rao *et al.*



Figure 1. Mango ginger plant (A), rhizomes (B) and TS of rhizome (C).




Figure 2. Volatile constituents of mango ginger.

1989; Choudhury *et al.* 1996; Srivastava *et al.* 2001; Singh *et al.* 2003a,b; Mustafa *et al.* 2005). The *cis*- and *trans*-hydroocimene, ocimene and myrcene were found to be the major compounds present in the volatile oils of *C. amada*, which indicates that the aroma of mango ginger is a mixture of characteristic compounds found in both raw mango and turmeric (Rao *et al.* 1989). The acetone extract of mango ginger is composed of colourless oil, curcumin, phytosterol and azulenogenic oil containing pinene, camphor, curcumene and ar-turmerone (Jain and Mishra 1964). There are more than 100 phytochemicals reported from fresh and dried extracts of *C. amada* as compiled by Jatoi *et al.* (2007).

4.2 Curcuminoids in mango ginger

The well-known curcumin, demethoxy curcumin and bis-demethoxy curcumin (figure 3) are the major constituents from acetone extract of *C. amada* (Gupta *et al.* 1999).

4.3 Phenolic content in mango ginger extracts

The free phenolic acids (figure 4) present in mango ginger are caffeic (26%, 195 mg/g), gentisic (24%, 180 mg/g) and ferulic (20%, 150 mg/g) followed by gallic (10%, 75 mg/g), cinnamic (7%, 52.5 mg/g), protocatechuic (7%, 52.5 mg/g) and small amounts of syringic (4%, 30 mg/g) and p-coumaric acids (2%, 15 mg/g) (Siddaraju and Dharmesh 2007). It also contains bound phenolic compounds like ferulic acid (47%, 391.5 mg/g) and cinnamic acid (29%, 237 mg/g), p-coumaric acid (11%, 95 mg/g), syringic acid (5%, 38.8 mg/g), caffeic acid (4%, 30.7 mg/g), gallic acid (1%, 11.5 mg/g) and gentisic acid (1%, 4.9 mg/g).

4.4 Terpenoid bioactive molecules

Three terpenoid bioactive compounds, viz. difurocumenol, amadannulen and amadaldehyde (figure 5), were successfully isolated and characterized from chloroform extract of *C. amada* rhizome (Policegoudra *et al.* 2007a,b, 2010). The bioactive compounds are antibacterial as well as antioxidant in nature with DPPH radical scavenging activity, superoxide radical scavenging activity, lipid peroxidation inhibitory activity, metal chelating activity and total reducing power. The above bioactive compounds were also potential platelet aggregation inhibitors and toxic for cancer cell lines (Policegoudra 2008; Policegoudra *et al.* 2010).




Figure 3. Curcuminoids from mango ginger.




Figure 4. Phenolic acids from mango ginger.

4.5 Functional attributes of *Curcuma amada* starch

It appears that starch from mango ginger has potential functional properties. The starch from an unconventional source like mango ginger was isolated and characterized by Policegoudra and Aradhya (2008). It has a distinct structural and biochemical features of its own. Mango ginger contains 1.3% ash, 9.8% moisture and 45% starch with 43% amylose. Morphologically, mango ginger starch resembles ginger starch granules, but it differs by the absence of fissures on the surface and its X-ray diffractogram pattern. Scanning electron micrograph (SEM) revealed the variations in shape of granules that appeared as round, elliptic, irregular and polygonal (figure 6). The granule size was between 3 and 20 μm for small granules and 20 and 48 μm for large

granules and belongs to 'B' type of starch, a characteristic feature of *Curcuma* species. Thus, it occupies a unique position between turmeric and ginger starch. The solubility and water holding capacity of mango ginger starch was linear with increase in temperature. High amylose content and low solubility are interesting attributes of mango ginger starch that need to be explored for preparation of nutraceutical products of metabolic advantages.

5. Accumulation pattern of bioactive compounds during development

It may have been a biological inevitability for mango ginger rhizome to develop compounds of multifunctional activity to counteract the diversified underground abiotic and biotic




Figure 5. Terpenoids from mango ginger.

challenges. Difurocumenonol proved to be one such compound with multifunctional properties found in mango

ginger. A distinct pattern of accumulation of phytochemicals and difurocumenonol was demonstrated along with




Figure 6. Starch granules of mango ginger rhizomes.

other visual parameters in mango ginger, which provided optimum period for harvesting of mango ginger rhizomes, endowed with phytochemicals for various medicinal properties mentioned earlier (Policegoudra *et al.* 2007c). Biosynthesis and accumulation pattern of difurocumenol, phenolics and protein concentrations are maximum in 150- to 180-day-old rhizomes. High concentration of bioactive compounds and other storage components are critical and play important role as a biomarkers to determine the quality index of *C. amada* rhizomes for its utility in food and pharmaceutical industry (Policegoudra *et al.* 2007c).

6. Fate of bioactive compounds at different temperatures

Stability of the phytochemical compounds and their unaltered biological activities in prolonged storage condition are the preferred qualities for production of nutraceutical and pharmaceutical products. Selection of plant material plays a crucial role in deciding the product quality. Less information is available in this regard especially with regard to bioactive phytochemicals. For the first time, difurocumenol,

a multi-functional bioactive terpenoid compound, has been successfully shown as biomarker to assess the bioactive quality changes in mango ginger rhizome during storage at different temperatures (Policegoudra 2008) (figure 7). The highest concentration of difurocumenol in 70 days stored rhizomes at ambient conditions becomes useless for product preparation due to decrease in its concentration of major phytochemicals. Retention of major phytochemicals and difurocumenol for longer period (120 days) can be achieved by storing mango ginger in low temperature ($14\pm 1^\circ\text{C}$). It also retained the mango flavour and minimized the changes in phytochemical composition and antioxidant properties. Thus, phytochemical and biological activity mango ginger was observed to be directly governed by temperature and time of storage (Policegoudra *et al.* 2007c).

7. Mango ginger and its use in traditional medicines (Ayurveda and Unani)

The ancient testimony for the use of plants as medicine was well documented in the treatise of *Ayurveda* and dates back to the pre-historic Vedic era. Accordingly, the medicinal




Figure 7. Changes in bioactive principles of mango ginger at different temperature.

properties of mango ginger are depicted in the following Sankrit shloka:

Raw mango flavoured ginger has a cooling effect on the body. It aggravates Vata.¹

It also pacifies deranged Pitta,² cures all types of itching and skin diseases.

आम्रगन्धिहरिद्रा या सा शीता वातला मता ।
पित्तहृत् मधुरा तिक्ता सर्वकण्डूविनाशिनी । ।
Āmragandhiharidrā yā sā śītā vātālā matā;
pittahṛt madhurā tiktā sarvakaṇḍūvināśinī.

Many species belonging to this genus have significant value as medicines, dyes and spices. Ayurveda, the oldest system of medicine in India, attributed multiple uses of rhizome as an appetizer, alexteric, antipyretic, aphrodisiac and laxative. Additional health benefits of *C. amada* rhizome reported were biliousness, itching, skin diseases, asthma and inflammation due to injuries (CSIR 1950). According to the Unani systems of medicine, it is a diuretic, maturant, emollient, expectorant, antipyretic and appetizer. Moreover, several reports have demonstrated the *C. amada* rhizome's ability against inflammation in the mouth and ear, gleet, ulcers on the male sex organs, scabies, lumbago and stomatitis (Kirtikar and Basu 1984; Warriar *et al.* 1994; Hussain *et al.* 1992).

8. Use of mango ginger in food industry

Mango ginger has a typical exotic flavour of raw unripe mango. Therefore, it is used as a basic ingredient in pickles, preserves, candies, sauces, curries, salads and so on (Verghese 1990; Shankaracharya 1982).

9. Biological activities of *Curcuma amada*

Rhizome, being a storehouse of bioactive compounds, has extensive use. Mango ginger is used medicinally as a coolant, aromatic and astringent and to promote digestion. A rhizome paste has traditionally been used for healing of wounds, cuts and itching (Srivastava *et al.* 2006). The external use of the rhizome paste for sprains and skin diseases is also an old practice (Gupta *et al.* 1999a,b). The rhizome has carminative properties, as well as being useful as a stomachic (Hussain *et al.* 1992).

¹ *Vata* – an Ayurvedic principle necessary to co-ordinate the function of the nervous system.

² *Pitta* – an Ayurvedic principle that uses bile to direct digestion and hence metabolism into the venous system.

Very few reports are available on the aerial parts of the plant. However, a whole-plant paste with crushed long peppers (*Piper longum*) is reported to be effective for the treatment of piles, and a decoction of the rhizome with common salt is an effective treatment for colds and coughs and is used to improve blood quality (Kapoor 1990). Topical use of native extract of leaves for contusions and sprains are also reported by several authors (Kirtikar and Basu 1984; Nadkarni 1954; Rao *et al.* 1989).

9.1 Antioxidant activity

The antioxidant activity of aqueous methanol extract of mango ginger leaves and rhizomes by β -carotene bleaching method are reported (Prakash *et al.* 2007). Leaf extract was more active than rhizomes extract. The antioxidant activity of sequential extracts of mango ginger with increasing polarity of solvents was reported by Policegoudra *et al.* (2007a). They reported different antioxidant activity assays like DPPH radical scavenging activity, superoxide radical scavenging activity, metal chelating activity and lipid peroxidation activity. Among hexane, chloroform, ethyl acetate, acetone and methanol extracts, ethyl acetate and acetone extracts showed good DPPH radical scavenging activity. The non-polar extracts showed good lipid peroxidation inhibitory activity. The antioxidant activity of mango ginger rhizomes in fresh and cooked form has demonstrated that there are no significant differences in the utilization of mango ginger for preparation of culinary preparations (Tarwadi and Agte 2005).

9.2 Antibacterial activity

The aqueous and organic solvent extracts of mango ginger are antibacterial against *Escherichia coli*, *Bacillus subtilis* and *Staphylococcus aureus* (Chandarana *et al.* 2005). The heated aqueous extract exhibited high antibacterial activity compared to the unheated aqueous extract. Chandarana *et al.* also reported that the solvent extracts (1,4-dioxan and DMF) were less effective against bacteria.

Antibacterial activity of free and bound phenolics from mango ginger rhizomes has been reported by Siddaraju and Dharmesh (2007). Both free and bound phenolic fractions of mango ginger are effective in inhibiting H⁺, K⁺-ATPase activity and *Helicobacter pylori* growth. Cinnamic and ferulic acids present in phenolic fractions of mango ginger contribute significantly to H⁺, K⁺-ATPase as well as *H. pylori* growth inhibition. The different extracts like hexane, chloroform, ethylacetate, acetone and methanol extracts are highly antibacterial against *Bacillus cereus*, *B. subtilis*, *Micrococcus luteus*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Enterococcus fecalis* and *Salmonella typhi* (Policegoudra *et al.* 2007a, b). These extracts showed antibacterial activity against wide

range of bacteria. The chloroform extract was more effective when compared to other extracts. However, *E. coli*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Proteus mirabilis* and *Yersinia enterocolitica* were not inhibited by any of the extracts of mango ginger.

9.3 Antifungal activity

The volatile oil from mango ginger rhizomes has antifungal in nature (Singh *et al.* 2002). Myrcene (4.6%) and pinene (80.5%) are the major components of volatile oils responsible for antifungal activity against the wide range of fungi, viz. *Curvularia palliscens*, *Aspergillus niger*, *A. terreus*, *Fusarium moniliforme* and *F. falcatum*.

9.4 Anti-inflammatory activity

The ethyl alcohol extract of mango ginger rhizome has anti-inflammatory activity in acute and chronic administration in albino rats (Mujumdar *et al.* 2000). Mujumdar *et al.* also reported the presence of chemical compounds with hydroxyl, ester, carbonyl and olefin functional groups in ethyl alcohol extract. It was found to be significant at higher concentrations in acute carrageenan-induced rat paw oedema model.

9.5 Platelet aggregation inhibitory activity

Platelet aggregation inhibitory activity of ethyl acetate extract and acetone extract is reported to be very high compared to methanol extract (Policegoudra 2008). The high platelet aggregation inhibitory activity of ethyl acetate, acetone and methanol extracts appears to be correlated with high phenolic content and to be concentration dependent.

9.6 Cytotoxicity

Cytotoxicity of the hexane, chloroform, ethyl acetate, acetone and methanol extracts of mango ginger towards both normal and cancer cell cultures were reported (Policegoudra 2008). All the extracts showed comparatively higher toxicity towards cancer cells when compared with normal cells, which is a good indicator of the anticancer property of extracts. Among the five extracts tested, the ethyl acetate extract showed greater toxicity, followed by chloroform, hexane, and acetone and methanol extract. The cytotoxicity results of different extracts of mango ginger indicate that the extracts are less toxic towards the normal cell lines.

9.7 Antiallergic activity

C. amada is reportedly used in various herbal preparations, including antiallergy formulations (Pushpangadan *et al.* 2006).

9.8 Biopesticide activity

C. amada is highly effective insecticide or pesticide as reported by several studies. *C. amada* was exhibited 100% adult mortality and a reduction in oviposition, even at 0.5% concentration. Essential oils from *C. amada* showed 100% repellent activity (Singh and Singh 1991). Biochemical studies of these promising oils may lead to the development of future pesticides. Insect mortality was 100% at 1% levels after 45 days. In addition, *C. amada* was highly effective in inhibiting the emergence of the F1 generation of weevils (Ahmad and Ahmad 1991).

9.9 Hypotriglyceridemic activity

Mango ginger extract showed hypotriglyceridemic activity and influences on both liver synthesis and blood clearance (Srinivasan and Chandrashekharan 1992, 1993). They reported hypotriglyceridemic activity on Triton-induced hyperlipidemic rats.

9.10 Brine-shrimp lethal activity

The water extract of mango ginger rhizomes showed brine-shrimp (*Artemia salina*) lethal activity (Krishnaraju *et al.* 2006). The lethality value (LC₅₀=6,600 µg, 24 h) was determined by a plot of the percentage of the shrimps killed against the concentrations of the extracts. The degree of lethality was found to be directly proportional to the concentration of the extract.

9.11 CNS depressant and analgesic activity

A fraction obtained from ethanol extract of mango ginger rhizome exhibited CNS depressant and analgesic activity (Mujumdar *et al.* 2004). The active fraction showed reduction in exploratory activity of barbiturate sleeping time, indicating CNS depressant activity. Further, it also showed reduction in acetic acid-induced writhings, tail-flick response, and carrageenan-induced inflammation, indicating potential antinociceptive and antiphlogistic activity.

9.12 Enterokinase inhibitory activity and antitubercular activity

The mango ginger also has enterokinase inhibitory activity (Bhat *et al.* 1981) and antitubercular activity. The labdane-type diterpenoid, labda-8 (17),12-diene-15,16-dial and its modified analogues have antitubercular properties (Singh *et al.* 2010).

10. Conclusions

The currently available drugs in the allopathic system of medicine are not so effective in combating a wide variety of

complications. The remedial measure may lie in the phytochemicals. Mango ginger appears to be highly potential and had remained unexplored for their bioactive phytochemicals. The available literature on phytochemicals and biological and pharmaceutical properties activities are very impressive. Very less information is available on the aerial parts of the plant. An array of phytochemicals like phenolics, terpenoids and other bioactive constituents were reported in mango ginger rhizome. They were recognized for various biological activities such as antimicrobial, antioxidant, anticancer, anti-inflammatory, antidepressant, antitubercular and platelet aggregation inhibitory activities. It may be concluded that *C. amada* contains a number of bioactive divergent ingredients in which one ingredient may act to enhance the action of other ingredient. The varied phytochemical and biological activities of *C. amada* reported in the present review may confirm the therapeutic value, for its combating abilities and for its use against multiple diseases. Further, it may be scientific validation for bioactive properties of mango ginger rhizome and its usage in Ayurveda and other traditional medicines. However, the structure–activity relationships and pharmacological activity of these constituents is the need of the hour. Further synthesis of active principles can lead to development of pharmacological products for health benefits.

References

- Ahmad PYA and Ahmad SM 1991 Potential of some rhizomes of Zingiberaceae family as grain protectants against storage insect pests. *J. Food Sci. Technol.* **28** 375–377
- Aminul I 2004 Genetic diversity of the genus *Curcuma* in Bangladesh and further biotechnological approaches for *in vitro* regeneration and long-term conservation of *C. longa* germplasm. PhD Thesis, University of Hannover, Germany
- Bhat PG, Jacob RT and Pattabiraman TN 1981 Enzyme inhibitors from plants: Enterokinase inhibitors in tubers and seeds. *J. Biosci.* **3** 371–378
- Chandarana H, Baluja S and Chanda SV 2005 Comparison of antibacterial activities of selected species of Zingiberaceae family and some synthetic compounds. *Tur. J. Biol.* **29** 83–97
- Choudhury SN, Rabha LC, Kanjilal PB, Ghosh AC and Leclercq PA 1996 Essential oil of *Curcuma amada* Roxb. from northeastern India. *J. Ess. Oil Res.* **8** 79–80
- Council of Scientific and Industrial Research (CSIR) 1950 Raw materials; In *Wealth of India* (New Delhi: CSIR) pp 401
- Dutt S and Tayal JN 1941 Chemical examination of the essential oil derived from the rhizomes of *Curcuma amada* Roxb. *Ind. Soap J.* **7** 200–205
- Golap SG and Bandyopadhyaya C 1984 Characterization of mango-like aroma in *Curcuma amada* Roxb. *J. Agric. Food Chem.* **32** 57–59
- Gupta AP, Gupta AM and Sushil Kumar K 1999 Simultaneous determination of curcuminoids in curcuma samples using high performance thin layer chromatography. *J. Liq. Chromatog. Technol.* **22** 1561–1569
- Gupta AP, Gupta MM and Kumar SJ 1999 *Liq. Chromatogr. Rel. Technol.* **22** 1561–1569
- Gupta M, Shaw BO and Mukherjee A 2010 A new glycosidic flavonoid from *J warhar* mahakashay (antipyretic) Ayurvedic preparation. *Int. J. Ayur. Res.* **1** 106–111
- Hussain A, Virmani OP, Popli SP, Misra LN and Gupta MM 1992; in *Dictionary of Indian medicinal plants* (Lucknow: CIMAP) pp 161–162
- Jain MK and Mishra RK 1964 Chemical examination of *C. amada* Roxb. *Indian J. Chem.* **2** 39
- Jatoi SA, Kikuchi A, Gilani SA, Watanabe KN 2007 Phytochemical, pharmacological and ethnobotanical studies in mango ginger (*Curcuma amada* Roxb., Zingiberaceae). *Phytother. Res.* **21** 507–516
- Kapoor LD 1990 *CRC Handbook of Ayurvedic medicinal plants* (Boca Raton, FL: CRC Press)
- Kirtikar KR and Basu BD 1984; in *Indian medicinal plants* 2nd edition (Dehra Dun: Bishen Singh Mahendra Pal Singh) pp 2422–2423
- Krishnaraju AV, Rao TVN, Sundararaju D, Vanisree M, Tsay HS and Subbaraju GV 2006 Biological Screening of Medicinal Plants Collected from Eastern Ghats of India Using *Artemia salina* (Brine Shrimp Test). *Int. J. Appl. Sci. Eng.* **4** 115–125
- Lakshminarayana S, Siddappa GS, SubbaRao MS and Johar DS 1963 Preservation of mango ginger. *Food Sci.* **12** 11–14
- Moon K, Khadabadi SS, Deokate UA and Deore SL 2010 *Caesalpinia bonducella* F – An overview. *Report Opin.* **2** 83–90
- Mujumdar AM, Naik DG, Dandge CN and Puntambekar HM 2000 Anti-inflammatory activity of *Curcuma amada* Roxb. in albino rats. *Indian J. Pharmacol.* **32** 375–377
- Mujumdar AM, Naik DG, Misar AV, Puntambekar HM and Dandge CN 2004 CNS depressant and analgesic activity of a fraction isolated from ethanol extract of *Curcuma amada* rhizomes. *Pharma. Biol.* **42** 542–546
- Mustafa A, Ali M and Khan ZN 2005 Volatile oil constituents of the fresh rhizomes of *Curcuma amada* Roxb. *J. Ess. Oil Res.* **17** 490–491
- Nadkarni KM 1954 *The Indian materia medica* (Bombay: Popular Book Depot) p 412
- Policegoudra RS 2008 Functional properties of bioactive molecules from mango ginger (*Curcuma amada* Roxb.) and its applications in food. PhD Thesis, Mysore University, Mysore, India
- Policegoudra RS and Aradhya SM 2007c Biochemical changes and antioxidant activity of mango ginger (*Curcuma amada* Roxb.) rhizomes during postharvest storage at different temperatures. *Postharvest Biol. Technol.* **46** 189–194
- Policegoudra RS and Aradhya SM 2008 Structure and biochemical properties of starch from an unconventional source - a mango ginger (*Curcuma amada* Roxb.) rhizome. *Food Hydrocoll.* **22** 513–519
- Policegoudra RS, Divakar S and Aradhya SM 2007a Identification of Difurocumenonol, a new antimicrobial compound isolated from mango ginger (*Curcuma amada* Roxb.) rhizome. *J. Appl. Microbiol.* **102** 1594–1602

- Policegoudra RS, Abiraj K, Channe Gowda D and Aradhya SM 2007b Isolation and characterization of antioxidant and antibacterial compound from mango ginger (*Curcuma amada* Roxb.) rhizome. *J. Chromatogr. B* **852** 40–48
- Policegoudra RS, Swaroop Kumar HM and Aradhya SM 2007d Accumulation of bioactive compounds and their bioactivity during developmental stages of mango ginger (*Curcuma amada* Roxb.) rhizome. *J. Agric. Food Chem.* **55** 8105–8111
- Policegoudra RS, Rehna K, Rao LJ and Aradhya SM 2010 Antimicrobial, antioxidant, cytotoxicity and platelet aggregation inhibitory activity of a novel molecule isolated and characterized from mango ginger (*Curcuma amada* Roxb.) rhizome. *J. Biosci.* **35** 231–240
- Prakash D, Suri S, Upadhyay G and Singh B 2007 Total phenol, antioxidant and free radical scavenging activities of some medicinal plants. *Int. J. Food Sci. Nutr.* **58** 18–28
- Purseglove JW 1974 Tropical crops: monocotyledons. Longman Group Ltd, London
- Pushpangadan PR, Chandana VR, Ajay KS, Ojha SK and Reddy GD 2006 Anti-allergic herbal formulation(s). Patent No. WO 2006067802
- Rao AS, Rajanikanth B and Seshadri R 1989 Volatile aroma components of *Curcuma amada* Roxb. *J. Agric. Food Chem.* **37** 740–743
- Salvi ND, George L and Eapen S 2000 Direct regeneration of shoots from immature inflorescence cultures of turmeric. *Plant Cell Tiss. Org.* **62** 235–238
- Sasikumar B 2005 Genetic resources of *Curcuma*: diversity, characterization and utilization. *Plant Gen. Res.* **3** 230–251
- Shankaracharya NB 1982 Mango ginger. *Indian Cocoa, Arecanut Spices J.* **5** 78–80
- Shirgurkar MV, John CK and Nadgouda RS 2001 Factors affecting microrhizome production in turmeric. *Plant Cell Tiss. Org.* **64** 5–11
- Siddaraju MN and Dharmesh SM 2007 Inhibition of Gastric H⁺,K⁺-ATPase and *Helicobacter pylori* Growth by Phenolic Antioxidants of *Curcuma amada*. *J. Agric. Food Chem.* **55** 7377–7386
- Singh D and Singh AK 1991. Repellent and insecticidal properties of essential oils against housefly (*Musca domestica* L). *Insect Sci. Applic.* **12** 487–492
- Singh G, Singh OP and Maurya S 2002 Chemical and biocidal investigations on essential oils of some Indian *Curcuma* species. *Prog. Crystal Growth Charact.* **45** 75–81
- Singh G, Singh OP, Lampasona M P and Catalan C 2003a *Curcuma amada* Roxb – Chemical composition of rhizome oil. *Ind. Perfum.* **47** 143–146
- Singh N, Singh J, Kaur L, Sodhi NS and Gill BS 2003b Morphological, thermal and rheological properties of starches from different botanical sources - a review. *Food Chem.* **81** 219–231
- Singh S, Kotesch Kumar J, Saikia D, Shanker K, Thakur J, Negi A and Banerjee S 2010 A bioactive labdane diterpenoid from *Curcuma amada* and its semisynthetic analogues as antitubercular agents. *Eur. J. Med. Chem.* **45** 4379–4382
- Srinivasan MR and Chandrashekhara N 1992 Effect of Mango ginger on lipid status in normal and hypertriglyceridemic rats. *J. Food Sci. Technol.* **29** 130–132
- Srinivasan MR and Chandrashekhara N 1993 Effect of Mango ginger on Triton WR-1338 Induced hyperlipidemia and plasma lipases activity in the rats. *Nutr. Res.* **13** 1183–1190.
- Srivastava AK, Srivastava SK and Shah NC 2001 Constituents of the rhizome essential oil of *Curcuma amada* Roxb. from India. *J. Ess. Oil Res.* **13** 63–64
- Srivastava S, Srivastava M, Rawat AKS and Mehrotra S 2006 Pharmacognostic evaluation of *Curcuma amada* Roxb. *Proc. Nat. Acad. Sci. Ind.* **76** 153–160
- Tarwadi K and Agte V 2005 Antioxidant and micronutrient quality of fruit and root vegetables from the Indian subcontinent and their comparative performance with green leafy vegetables and fruits. *J. Sci. Food Agric.* **85** 1469–1476
- Velayudhan KC, Muralidharan VK, Amalraj VA, Gautam PL, Mandal S and Kumar D 1999 *Curcuma* genetic resources. Scientific monograph No. 4. National Bureau of Plant Genetic Resources Regional Station, Thrissur, Kerala, India, pp 1–27
- Vergheze J 1990 Mango ginger - an exotic flavourant. *Indian Spices* **27** 15–16
- Warrier PK, Nambiar VPK and Ramankutty C 1994; in *Indian medicinal plants – a compendium of 500 species* (Chennai: Orient Longman Pvt Ltd) p 106

MS received 15 December 2010; accepted 13 June 2011

ePublication: 16 August 2011

Corresponding editor: SUDHA BHATTACHARYA