



Review

An update and perspectives on the use of promoters in plant genetic engineering

DIVYA KUMMARI^{1,2}, SUDHAKAR REDDY PALAKOLANU^{1*} ,
P B KAVI KISHOR³, POOJA BHATNAGAR-MATHUR¹, PRASANTH SINGAM²,
VINCENT VADEZ¹ and KIRAN K SHARMA¹

¹International Crops Research Institute for the Semi-Arid Tropics (ICRISAT),
Patancheru, Hyderabad 502 324, India

²Department of Genetics, Osmania University, Hyderabad 500 007, India

³Department of Biotechnology, Vignan's Foundation for Science, Technology and Research Deemed
to be University, Vadlamudi, Guntur 522 213, India

*Corresponding author (Email, p.sudhakarreddy@cgiar.org, palakolanusreddy@gmail.com)

Genetically engineered plants have varied applications in agriculture for enhancing the values of food and feed. Genetic engineering aims to introduce selected genetic regions with desirable traits into target plants for both spatial and temporal expressions. Promoters are the key elements responsible for regulating gene expressions by modulating the transcription factors (TFs) through recognition of RNA polymerases. Based on their recognition and expression, RNA polymerases were categorized into *RNA pol II* and *pol III* promoters. Promoter activity and specificity are the two prime parameters in regulating the transgene expression. Since the use of constitutive promoters like *Cauliflower mosaic virus (CaMV) 35S* may lead to adverse effects on non-target organisms or ecosystem, inducible/tissue specific promoters and/or the *RNA pol III* promoters provide myriad opportunities for gene expressions with controlled regulation and with minimum adverse effects. Besides their role in transgene expression, their influence in synthetic biology and genome editing are also discussed. This review provides an update on the importance, current prospects, and insight into the advantages and disadvantages of promoters reported thus far would help to utilize them in the endeavour to develop nutritionally and agronomically improved transgenic crops for commercialization.

Keywords. *CaMV35S* promoter; constitutive promoters; synthetic promoters; genetic engineering; *RNA pol II* promoters; *RNA pol III* promoters; *U3* promoter; *U6* promoter

1. Introduction

Promoters are gene switches located upstream of gene coding regions, which turn on and off the functional activity of genes and contain specific *cis*-acting elements which are binding targets for proteins involved in the

initiation and regulation of transcription. Promoters are molecular biological clocks crucial for choice of the targeted gene expression (Potenza *et al.* 2004), that act as key regulatory check points for transcription of genes that are recognized by transcription factors (TFs) (Smale and Kadonaga 2003). TFs bind to specific *cis*-acting elements present on the respective promoter sequences through RNA polymerase and regulate expression of the downstream genes (Hernandez-Garcia and Finer 2014). Promoters of coding genes often contain core, proximal

This article is part of the Topical Collection: Genetic Intervention in Plants: Mechanisms and Benefits.

and distal regions. Further, proximal and distal regions of the promoter contain different regulatory sequences such as enhancers, silencers, insulators, and *cis*-elements that contribute to fine regulation of gene expression at the transcriptional level (Hernandez-Garcia and Finer 2014). Detailed structural properties of promoters and gene regulatory elements were discussed earlier by Hernandez-Garcia and Finer (2014), Porto *et al.* (2014) and Shah *et al.* (2015). Promoter sequences that regulate the controlled transgene expression in plants are useful for developing the genetically modified (GM) crops with improved agronomical and nutritional traits (Mittler and Blumwald 2010). There are few promoter prediction bioinformatics tools; one among them is TSSPlant, a novel tool that predicts both TATA and TATA-less promoters in sequences of a wide spectrum of plant genomes (Shahmuradov *et al.* 2017). As complete genome sequences of most of the plant species are becoming available, several promoters are being identified, isolated and evaluated, while many more are likely to be elucidated in the near future. Predicted regulatory sequences may or may not be functionally active or necessitating the confirmation of the role of specific elements for promoter activity. Promoters can be isolated through different methods some of which being: genome walking, inverse PCR (IPCR) plasmid rescue, screening of genomic DNA library constructed from mutant plant, the thermal asymmetric interlaced PCR TAIL-PCR (Reddy *et al.* 2008; Liu and Whittier 1995). Quick, efficient, predictable, and high-throughput analysis of gene expressions and their promoters will be crucial for validating the functional regulatory element sequences.

Promoters can be selected to develop transgenic plants based on the type of trait and target tissue to be regulated (Bilas *et al.* 2016a, b). A variety of plant promoters that regulate the degree of expression of a transgene can be obtained from various sources. These can be categorized into *pol II* and *pol III* that are activated upon recognition by the RNA polymerases II and III. *Pol II* promoters can in turn be classified into constitutive, tissue-specific, stress-inducible and synthetic (Bilas *et al.* 2016a, b), whereas *pol III* promoters are U3 and U6 (Marshall *et al.* 1992). *CaMV35S* (commonly referred to as 35S) constitutive promoter is the most widely used for gene expressions in transgenic plants and in basic functional genomic studies (Porto *et al.* 2014). Hence, specific or inducible promoters not only provide increased gene expression in a tissue of interest or at specific developmental stage(s), but also provide more predictable gene expression, with minimal or no penalties on final yield. The discovery of proteins with programmable DNA-binding specificities has triggered an array of

applications in synthetic biology, regulation of transcription, including genome editing, and epigenetic modifications. Among those, transcription activator-like effectors (TALENs) and the clustered regularly interspaced short palindromic repeats (CRISPR/Cas9) system played high attention due to their in-built function as transcription regulators for the control of gene expression. Recent studies have shown that the expression of Cas9/gRNA with the help of specific promoters is important for highly efficient genome editing in plants (Hyun *et al.* 2015; Wang *et al.* 2015; Yan *et al.* 2015; Mao *et al.* 2016). Here, we attempt to provide a comprehensive overview of promoters in plants, their selection, specificity, cross activity, and how they can be useful in maximizing the transgene expression for potential applications in crop improvement programs.

2. Types of promoters

Based on the type of recognition of RNA polymerases, they are mainly classified into RNA polymerase (*pol II* and *pol III*) promoters. Detailed information and study of these promoters are provided in the subsequent sections of this review.

2.1 RNA *pol II* promoters

RNA *pol II* promoters are the shortest sequences where RNA *pol II* polymerase binds to the DNA to initiate transcription with help of the TATA box, a common component located at -35bp upstream of the transcription start codon. Based on the type of expression, *pol II* promoters are further categorized into constitutive, organ- or tissue-specific, stress-inducible and synthetic (figure 1).

2.1.1 Constitutive promoters: Constitutive promoters express constitutively throughout the plant life cycle, irrespective of the external and developmental factors which is beneficial for expression of the industrial enzymes, insect resistance and selectable marker genes (Jiang *et al.* 2018). Based on the source of origin, they are further classified into viral, bacterial and plant constitutive promoters.

2.1.1.1 Viral: In 1980s, Chua and collaborators isolated a promoter, responsible for the transcription of the whole genome of a Cauliflower mosaic virus (CaMV) infecting turnips. Later, this was named as *CaMV35S*




Figure 1. A flow chart showing different types of RNA Pol II promoters categorized based on their mode of expression into four major classes: constitutive, tissue-specific, stress-inducible and synthetic promoters. Each of these promoters are further categorized into different types; constitutive promoters (viral, bacterial, plant), tissue-specific promoters (root, fruit, seed), stress-inducible promoters (abiotic {light, wound, heat, drought, salt, cold, hormone and chemical} and biotic {pathogen, pest, weed}).

promoter due to the sedimentation coefficient of the viral transcript whose expression is driven by *CaMV35S* promoter and used for constitutive overexpression of genes (Odell *et al.*, 1985). There is not much variation in the activities between the full-length *CaMV35S* promoter (−941 to +9 bp) (Odell *et al.* 1985) and −343bp deletion fragment (Fang *et al.* 1989). Although the *CaMV35S* promoter and its derivatives can drive high levels of transgene expression in dicotyledonous plants (Benfey *et al.* 1990a, 1990b), their activities are substantially lower in monocotyledonous plants (Gupta *et al.* 2001). The expression of *uidA* gene under *CaMV35S* promoter was observed in different parts of transgenic tobacco with maximum levels in leaf and root tissues (Malik *et al.* 2002). In contrast, *uidA* gene expression under the *CaMV35S* promoter in transgenic canola and *Arabidopsis* showed expression in all plant tissues (Malik *et al.* 2002). Sharma *et al.* (2006) demonstrated a higher activity of the *cryIAb* gene under the constitutive *CaMV35S* promoter in flowers of pigeonpea, although other parts like leaves, seeds and pod walls also showed expression of the transgenes to varying degrees.

Apart from *CaMV35S*, *peanut chlorotic streak caulimovirus (PCISV)* and *figwort mosaic virus (FMV)* have also been shown to be very useful for generating GM plants. *PCISV* is a strong constitutive promoter

and comparative analysis with *FMV* promoter in GM plants showed similar expression pattern. Although functionally analogous, the nucleotide sequence of *PCISV* promoter has limited homology with other caulimovirus promoters (Maiti and Shepherd 1998). The *FMV Sgt* (Figwort mosaic virus sub-genomic transcript) promoter is another constitutive promoter whose expression was observed in all tissues (Bhattacharyya *et al.* 2002). A comparative analysis of the *FMV Sgt* promoter with that of *CaMV35S* promoter showed that it was 2-folds stronger than the latter, and was also less active in monocots like maize where its expression was about 27.5-folds lower than in tobacco (Bhattacharyya *et al.* 2002). Gemini viral promoters are in general located within the intergenic region, though promoters have also been identified within the genes. In the same way, Gemini virus-associated beta satellite harbors a promoter element for driving the expression of its only ORF, which can be useful in the regulation of plant genes (Borah *et al.* 2016). TaB virus-derived promoters may be useful for the high level constitutive expression of transgenes either in monocotyledonous or dicotyledonous species (Yang *et al.* 2003), thus proving its versatility. *Rep* promoter from *cotton leaf curl Burewala virus (CLCuBuV)* showed consistent strong transient expression in tobacco and cotton leaves

as compared to the *CaMV35S* promoter, whereas other *CP* promoter showed lower expression (Khan *et al.* 2015). Promoters originating from the *tomato leaf curl virus* (*TLCV*) drive both constitutive as well as tissue-specific expression in transgenic tobacco (Seemanpillai *et al.*, 2003). But it is not widely used and accepted from the research community. Comparative analysis of the *CaMV35S* and the enhanced *CaMV35S* (*E35S*), *CsVMV*, *FMV*, and the *Strawberry vein banding virus* (*SVBV2*) promoters indicated that the *FMV* promoter facilitated a strong expression of target genes in soybean hairy roots and root nodules (Govindarajulu *et al.* 2008). Transgenic maize with *aryloxyalkanoate dioxygenase* (*aad-1*) expression under the influence of different promoters demonstrated that viral promoters such as *CaMV35S* and *SCBV* produce lower transformation efficiencies, but higher percentages of low copy number events contrary to plant constitutive promoters like *OsAct1* and *ZmUbi1* (Beringer *et al.* 2017). These promoters are recommended for wide use since they promote low copy number transgenic events. Various constitutively expressed viral promoters used in the development of transgenic plants are listed in table 1.

2.1.1.2 Bacterial: Bacterial-origin promoters such as *nopaline synthase* (*nos*), *octopine synthase* (*ocs*) and *mannopine synthase* (*mas*) have been isolated from *Agrobacterium tumefaciens* and used for the control the transgene expression. Although their level of expression and activity can be affected by hormones and wounding, they are repeatedly being used for the transformation of plants. *mas* promoters can be used as enhancers or silencers, since they have the ability to bind with the factors of nuclear protein from different plants (Shah *et al.* 2015). Thus, it appears that both *mas* and *CaMV35S* promoters can be used for the construction of improved plant transformation vectors. *Tn5 neomycin phosphotransferase* (*nptH*) gene was expressed with either of these promoters (Kevin *et al.* 1989). Sequence analysis of the *AV3* promoter from *Ageratum yellow vein virus* (*AYVV*) showed that it might be a remnant of prokaryotic ancestors that could be related to certain promoters of bacteria from marine or freshwater environments (Wang *et al.* 2013). Bacterial promoters show differential expression in transgenic plants, and regulation of *rol* gene expression plays a role in the biological effects that are caused by the *rol A*, *B*, and *C* genes containing the *uidA* reporter gene under the control of *rol A*, *B*, and *C* promoters of *Agrobacterium rhizogenes* (Schmullig *et al.* 1989). However, their use for generating transgenic crop plants is limited.

2.1.1.3 Plant: Promoters derived from plant origin are used for the constitutive transgene expressions in plants (Dhankher *et al.* 2002). Plant constitutive promoters such as rice *actin* (*OsAct1*) (McElroy *et al.* 1991), maize *alcohol dehydrogenase1* (*ZmAdh1*) (Fromm *et al.* 1990), and maize *ubiquitin* (*ZmUbi1* and *ZmUbi2*) (Christensen *et al.* 1992) are the most commonly used in crop plants. *Act2* promoter was obtained from the *actin* gene family, which is the cytoskeletal component and expressed in every plant cell (An *et al.* 1996). Similarly, the *OsActin1* promoter also displayed expression in almost all tissues when transformed back into rice (McElroy *et al.* 1991). The *OsAct* promoter was shown to drive high levels of expression of *HVA1* gene in the leaf and root tissues of rice, thereby conferring salt and water stress tolerance (Xu *et al.* 1996). *ZmAdh* is another constitutive promoter expressed in specific tissues like root, shoot meristems, endosperms and pollen (Kyoizuka *et al.* 1991). When *MtHP* promoter from *Medicago* was fused to a *uidA* gene, it showed expression in various plant parts in *Medicago* and *Arabidopsis* with an expression pattern similar to that of the *CaMV35S* promoter (Xiao *et al.* 2005). Expression of *uidA* gene driven by the *CaMV35S* and *MpEF1 α* promoters were compared throughout plant development. While *CaMV35S* promoter resulted in inadequate expression in the meristematic tissues and a strong expression in the callus, the *MpEF1 α* -promoter caused a strong meristematic *uidA* gene expression and was more active in female sexual tissues. It appears therefore, *MpEF1 α* - promoter is a better option for obtaining strong and ubiquitous transgene expression compared to the *CaMV35S* promoter as has also been pointed out by Althoff *et al.* 2014.

Expression of *uidA* under the influence of *UBQ1* or *UBQ2* rice ubiquitin promoters were 8 to 35-folds higher in transgenic rice plants, respectively, when compared to *CaMV35S* (Wang and Oard 2003). This indicates that *ubiquitin* promoters are superior for activating the transgenes. Expression of the *gfp* gene mediated by a *GmUbi* showed high levels of constitutive expression in soybean tissues, thereby providing an alternative to viral promoters for driving gene expression in soybean (Hernandez-Garcia *et al.* 2009). The native *ubil* promoter is a promising genetic element for constitutive expression of any gene in rice tissues (Bhattacharyya *et al.* 2011). *OsCon1* promoter exhibited comparable activity with *OsCcl1*, *OsAct1* or *ZmUbi* promoters in most tissues, and even more active than the *CaMV35S* promoter in roots, seeds and calluses indicating that it is a novel constitutive promoter which could potentially be used for developing

Table 1. Constitutively expressed viral promoters used in development of transgenic plants

Promoter	Source	Host	References
<i>SVFLt</i>	<i>Peanut chlorotic streak caulimo virus (PCISV)</i>	Tobacco	Maiti and Shepherd (1998)
<i>Enhanced 35S (E35S)</i>	<i>Cauliflower mosaic virus 35S</i>	Soybean	Li <i>et al.</i> (2001) and Bhattacharyya <i>et al.</i> (2002)
<i>Sgt</i>	<i>Figwort mosaic virus (FMV)</i>	Tobacco/maize	Bhattacharyya <i>et al.</i> (2002)
<i>CsVMV</i>	<i>Cassava vein mosaic virus (CsVMV)</i>	Soybean/grape	Seemanpillai <i>et al.</i> (2003)
<i>sgRNA β 1, β 2, and γ</i>	<i>Barley stripe mosaic virus (BSMV)</i>	Tobacco	Johnson <i>et al.</i> (2003)
<i>SV (antisense)</i>	<i>Peanut chlorotic streak caulimovirus (PCISV)</i>	Tobacco	Bhattacharyya <i>et al.</i> (2003)
<i>35S</i>	<i>Cauliflower mosaic virus 35S</i>	<i>Arabidopsis</i> /Tobacco/Soybean/grape/ <i>Phaeodactylum</i> / <i>Jatropha</i>	Yang <i>et al.</i> (2003)
<i>CLCuBuV Rep/CLCuBuV CP</i>	<i>Cotton leaf curl Burewala virus (CLCuBuV)</i>	Tobacco/cotton	Obertello <i>et al.</i> (2005)
<i>FMV</i>	<i>Figwort mosaic virus (FMV)</i>	Soybean	Govindarajulu <i>et al.</i> (2008)
<i>Pptca1</i>	<i>Cytomegalo virus (CMV)</i>	<i>Phaeodactylum tricornutum</i>	Sakaue <i>et al.</i> (2008)
<i>e35S-4ocs</i>	<i>Cauliflower mosaic virus 35S</i>	<i>Allocasuarina verticillata</i>	Govindarajulu <i>et al.</i> (2008)
<i>AV3</i>	<i>Ageratum yellow vein virus (AYVV)</i>	<i>E. coli</i>	Wang <i>et al.</i> (2013)
<i>TLCV</i>	<i>Tomato leaf curl virus</i>	Tobacco	Khan <i>et al.</i> (2015)
<i>SVBV2</i>	<i>Strawberry vein banding virus</i>	Soybean	–
<i>T500/T600/ T1200</i>	<i>Taro bacilliform virus (TaBV)</i>	Banana, tobacco	–

transgenics (Li *et al.* 2014). Incorporation of *AtTCTP* promoter into creeping bent grass showed that it can be used as a plant-derived constitutive promoter for the expression of selectable marker genes, and as an alternative to the *CaMV35S* promoter for developing GM crops (Han *et al.* 2015). The plant-derived *JcUEP* promoter could be another alternative to the *CaMV35S* promoter for directing constitutive transgene expression in *Jatropha* and other plants (Tao *et al.* 2015). Pineapple *SUI1* and *L36* promoters seem to drive *uidA* expression in all tissues of *Arabidopsis* at levels comparable to that of the *CaMV35S* promoter (Koia *et al.* 2013). Studies on rice *APX* promoter show edits constitutive expression in the seed, root, blade, flower and leaf of transgenic rice (Park *et al.* 2010). The inducible activities of *PvUbi1* and *PvUbi2* promoters in switchgrass, rice and tobacco are strong constitutive promoter candidates that can be used in genetic transformation of both monocots and dicots (Mann *et al.* 2011). Promoters such as *pBdEF1 α* and *pBdUBI10* are constitutively expressed and highly active in maize, whereas *pBdGLUI* isolated from *Brachypodium* was clearly endosperm-specific, thereby indicating that it is an excellent resource for promoters for transgenic research in heterologous cereal species (Coussens *et al.* 2012).

The *KST1* partial promoter was shown to drive constitutive expression in the guard cells of monocots and dicots, as well as in both annual and perennial plants (Kelly *et al.* 2017). Three *Citrus sinensis* constitutive gene promoters confirmed their role in vegetative tissues (Erpen *et al.* 2018), thus enabling the researchers to use it specifically in vegetative tissues. On the other hand, Jiang *et al.* 2018 reported that *AtSCPL30* promoter could be an alternative for the *CaMV35S* in terms of reducing transgene silencing and also a good source for the multigene transformation. *GAPC2* and *EF1* promoter fragments from *Citrus sinensis* were expressed constitutively in the transgenic tobacco plants (Erpen-Dalla Corte *et al.* 2020). Description of constitutively expressed plant promoters used for the development of transgenic plants is listed in the table 2.

2.1.2 Tissue-specific promoters: Tissue-specific promoters drive the expression of a targeted gene in a specific tissue(s) or organ at a specific stage(s) of plant growth and development (figure 2b). Since the constitutive promoters cause potential negative effects such as metabolic burden on the transgenics, specific promoters are used to regulate and target gene expressions in an effective manner for improving traits like grain

Table 2. Description of constitutively expressed plant promoters used for the development of transgenic plants

Promoter	Source	Host	References
<i>UBQ1/UBQ2</i>	Rice	Rice	Wang and Oard (2003)
<i>UBQ1</i>	<i>Arabidopsis</i>	<i>Allocauarina verticillata</i>	Obertello et al. (2005)
<i>VR-ACSI</i>	Mung bean	Tobacco/ <i>Arabidopsis</i>	Cazzonelli et al. (2005)
<i>Ubi</i>	Soybean	Soybean	Hernandez-Garcia et al. (2009)
<i>Ubi1</i>	Rice	Rice	Bhattacharyya et al. (2011)
<i>Ubi1</i>	Corn	Rice	Bhattacharyya et al. (2011)
<i>GAI</i>	Rice	Rice	Bhattacharyya et al. (2011)
<i>Ubi1/Ubi2</i>	Switchgrass	Switchgrass/ Rice/ Tobacco	Mann et al. (2011)
<i>uceApro2</i>	Cotton	Cotton/ <i>Arabidopsis</i>	Viana et al. (2011)
<i>SUI1</i>	Pineapple	<i>Arabidopsis</i>	Koia et al. (2013)
<i>L36</i>	Pineapple	<i>Arabidopsis</i>	Koia et al. (2013)
<i>EF1α</i>	<i>Marchantia polymorpha</i>	<i>Marchantia polymorpha</i>	Althoff et al. (2014)
<i>EF1α/UBI10/GLU1</i>	<i>Brachypodium distachyon</i>	Maize	Coussens et al. (2012)
<i>Con1</i>	Rice	Rice	Li et al. (2014)
<i>TCTP</i>	<i>Arabidopsis</i>	<i>Agrostis stolonifera</i>	Han et al. (2015)
<i>RD29A/RD29B</i>	<i>Arabidopsis</i>	Soybean	Bihmidine et al. (2013)
<i>UEP</i>	<i>Jatropha curcas</i>	<i>Jatropha curcas</i> , <i>Arabidopsis</i>	Tao et al. (2015)
<i>KST1</i>	Potato	Potato, Tobacco, Cucumber, Grape, Barley	Kelly et al. (2017)
<i>CsCYP,CsGAPC2, CsEF1</i>	<i>C. sinensis</i>	<i>C. sinensi</i>	Erpen et al. (2018)
<i>AtSCPL30</i>	<i>Arabidopsis</i>	Tobacco	Jiang et al. (2018)
<i>CsGAPC2, CsEF1</i>	<i>Citrus sinensis</i>	Tobacco	Erpen-Dalla Corte et al. (2020)

nutritional quality. Due to the demand from biosafety regulators for less intrusive transgene expression, targeted expression of transgenes has become more important for the future development of value-added crops (Bucchini and Goldman 2002).

2.1.2.1 Promoters that help in biomass production: *Eucalyptus* transgenics with AtEXP4-pro:CBM2a showing increased plant height, enlargement of xylem, xylem fiber and vessel cells can be an attractive choice for plant biomass improvement (Keadtidumrongkul et al. 2017).

2.1.2.2 Promoters specific to guard cells and vascular tissues: A partial promoter of *KST1* from potato is active in a monocot and the first promoter reported to confer guard cell expression in barley and cucumber (Kelly et al. 2017). Thus, *KST1* appears to be a cell specific promoter and highly useful for manipulating proteins in the guard cells. The use of phloem specific promoters (*Citrus phloem protein 2-CsPP2*, *Arabidopsis thalianaphloem protein 2-AtPP2*, *A. thaliana sucrose transporter 2-AtSUC2*, and *sucrose synthase*

1-SUS1) might increase the chances of producing more bacterial disease resistant transgenic cultivars in many crop plants (Singer et al. 2011; Miyata et al. 2012).

2.1.2.3 Root-specific promoters: Root-specific promoters are of special interest since they help in understanding the root architecture and in alleviating drought and salt stress conditions. The *pNtRELI* promoter can be used to direct root-specific expression of target genes to protect the root from different abiotic stresses (Zhang et al. 2016a, b). Alternatively, Tau Class *Glutathione-S-Transferase (SbGSTU)* gene promoter of *Salicornia brachiata* can be used for both constitutive as well as stress-inducible expression of transgenes (Tiwari et al. 2016). More data are needed if such promoters help to mitigate the stress better in comparison with the constitutive promoters with minimum or no yield penalty. A root-preferential promoter *GmPRP2* isolated from soybean has been found useful in developing transgenics of novel soybean cultivars (Chen et al. 2014). Rice promoters such as *Os03g01700* and *Os02g37190* are highly active in the root tissues of rice and can be helpful for the root-




Figure 2. Schematic depiction of different types of promoters. (a) Stress-inducible promoters can be activated under any biotic or abiotic stress by binding the corresponding transcription factors and *cis*-acting elements. (b) Tissue-specific promoters are expressed in a particular tissue by binding the corresponding transcription factors and *cis*-acting elements. (c) Bidirectional promoters drive the transcription of two genes simultaneously that flank the promoter.

specific expression of target gene (Li *et al.* 2013a, b, c, d, e). Novel nematode-responsive root-specific promoter (NRRS) from *Arabidopsis* is seen to be expressed high in gall and root of transgenic *Arabidopsis* plants in defence to the infection of *Meloidogyne incognita* nematode (Kakrana *et al.* 2017).

2.1.2.4 Flower-specific promoters: High-level expression was recorded in a floral-enhanced manner for 1.45 kb promoter of endogenous ubiquitin extension protein (UEP1), although 9-fold lower *uidA* expression was noticed in the leaves (Annadana *et al.* 2002). Comparison between *CaMV35S* and *UEP1* promoters revealed that the *UEP1* has 40- to 85-fold higher expression in the flower, indicating its high tissue-specificity. The *Arabidopsis CER6* gene when transformed into *Chrysanthemum* resulted in high, specific expression in flower petals due to the *CER6* promoter, but had high variability as compared to the *UEP1* promoter (Hannoufa *et al.* 1996).

2.1.2.5 Fruit-specific promoters: For the fruit-specific expression, promoters related to fruit ripening such as *1-aminocyclopropane-1-carboxylate (ACCoxidase)*, *E8* and *polygalacturonase (PG)* have been characterized from apple (Atkinson *et al.* 1998) and tomato (Nicholass *et al.* 1995). A fruit-ripening-induced promoter 1(*RIP1*) of tomato is perhaps more crucial for controlled manipulation of ripening-related agronomic traits not only in tomato but also in many other fruit crops (Agarwal *et al.* 2017). In transgenic tomato, the promoter *ACC oxidase* was able to drive ripening-specific expression of a reporter gene in the fruit only with no detectable activities in other tissues and mature green fruits (Atkinson *et al.* 1998). Transgenic tomato plants with enhanced sweetness, flavor and nutritive values were developed by expressing the *monellin* gene under the control of *E8* promoter (Reddy *et al.* 2015). For the expression of target genes and directing expression of the virus-F protein as an oral vaccine to induce systemic immunity in mice, the promoter of the tomato *E8* gene was used successfully in a number of instances (Lewinsohn *et al.* 2001). Although this promoter drives the expression of transgenes in flowers (predominantly anthers), its invariable expression throughout the ripening fruit makes it very valuable. The improved quality of ripened tomatoes in terms of vitamins and micronutrients content makes it valuable for use in transgenic research (Hadley *et al.* 2002). For the production of biopharmaceuticals, the use of organ-specific promoters are important so as to express the

gene of interest in those organs that are able to produce protein(s) in an appropriate manner (Twyman 2003). Deletion analysis showed improvement in the quality and nutritional value of the fruits when the stearyl-acyl-carrier-protein desaturase (Des) promoter region located between -590 and +10 was used for transforming oil palm and also dicots (Saed *et al.* 2012). However, comparative studies between constitutive and inducible promoters elaborating the specific merits of inducible promoters especially in the fruits are very meager in the literature. Therefore, such a study is vital to pinpoint the advantages of inducible promoters over that of constitutive (table 3).

2.1.2.6 Seed-specific promoters: Seed-specific transgene expression has been used in many applications including nutritive value improvement, milled grain quality and production of industrial or pharmaceutical compounds (Ye *et al.* 2000; Cahoon and Shanklin 2000). Promoter's specific to seed storage genes is an attractive target for such uses in GM crops. Soybean β -conglycinin gene promoter was used for seed-specific expression, where gene expression was restricted to the embryo during the mid-to-late maturation phase (Lesard *et al.* 2002). Likewise, groundnut seed promoter (GSP) can potentially be used for modification of seed phenotypes in agronomically important crops (Sunkara *et al.* 2014). While the sunflower *helianthinin* full-length promoter was used to drive seed-specific expression in tobacco (Nunberg *et al.* 1994), the expression of the β -phaseolin promoter in transgenic tobacco plants was restricted to the developing seed only (van der Geest and Hall 1997). Several endosperm-specific promoters like those from *hordein* in barley (Forde *et al.* 1985), *glutenin* in wheat (Lamacchia *et al.* 2001) and *zein* in maize (Marzabal *et al.* 1998) have been studied extensively. The guar *MS* promoter could manifest its usage for directing

Table 3. Bacterial constitutively expressed promoters expressed in different plants

Promoter	Source	Host	References
<i>Rol A/B/C</i>	<i>Agrobacterium rhizogens</i>	-	Schmulling <i>et al.</i> (1989)
<i>AV3</i>	<i>Ageratum yellow vein virus (AYVV)</i>	<i>E.coli</i>	Wang <i>et al.</i> (2013)
<i>Ocs</i>	<i>Agrobacterium</i>	-	Zhou <i>et al.</i> (2013)
<i>Mas</i>	<i>Agrobacterium</i>	-	Zhou <i>et al.</i> (2013)
<i>Nos</i>	<i>Agrobacterium</i>	Rice	Zhou <i>et al.</i> (2013)

endosperm-specific expression of transgenes in legume species (Naoumkina and Dixon 2011). The expression of a 4kb promoter fragment from *granule-bound starch synthase1* (*gbss1*) gene in wheat is restricted to endosperm and developing kernel pericarp tissues only (Kluth *et al.* 2002).

The *WM403* promoter from watermelon may be useful in driving nucellus-specific gene expression in plants inclusive of candidate genes for important nucellus-specific attributes such as apospory or adventitious embryony (Dwivedi *et al.* 2010). *HaFAD2-1* promoter is as strong as the *CaMV35S* promoter even though it is a tissue-specific promoter and its activity is derived from the embryo, thereby confirming that it can be considered as a strong, highly specific seed promoter useful for genetic manipulation applications in the seed (Zavallo *et al.* 2010). The *Arabidopsis At4g12960* (*AtGILT*) promoter was employed as a canola seed coat outer integument-specific promoter after the production and selection of desired transformants from many transgenic lines (Wu *et al.* 2011). Seed coat-specific promoters can also be used to assess the effects of many pathway enzymes/proteins and cell-wall modifying proteins on mucilage structure as has been pointed out by Dean *et al.* (2017). Maize *BD1*, *Def1* and *Def2* promoters were active and reproduced the expression patterns of both *Def1* and *Def2* genes in transformed immature maize embryos, as well as in developing seeds of transgenic maize (Liu *et al.* 2016). Thus, an array of tissue-specific promoters has been used in diverse taxa with many beneficial effects. Table 4 highlights a list of tissue-specific promoters that have been used so far in plant genetic engineering studies.

2.1.2.7 Underground storage-tissue-specific promoters: Promoters like *pDJ3S* and to a lesser extent *pMe1* from yam and casava respectively drive the high and preferential expression of genes in carrot storage roots (Arango *et al.* 2010). Such promoters help in improving the biomass and nutritional quality of the underground storage tissues.

2.1.3 Abiotic stress inducible: Plant performance is not conditioned to endogenous factors alone, but also to environmental factors and external stimuli. Several promoters are induced by hormones, chemicals, and environmental stresses (figure 2a) (Chakravarthi *et al.* 2016). Based on the source and type of cells in which they regulate gene expression, hundreds of inducible promoters have been identified (Singh *et al.* 2002). For induction of promoters under different abiotic stresses,

Table 4. List of tissue-specific promoters expressed in different plant species

Promoter	Source	Host	References
<i>aPAL</i>	<i>Loblolly pine</i>	Tobacco	Osakabe <i>et al.</i> (2009)
<i>DJ3S</i>	Yam	Carrot	Arango <i>et al.</i> (2010)
<i>Me1</i>	Casava	Carrot	Arango <i>et al.</i> (2010)
<i>WM403</i>	Watermelon	<i>Arabidopsis</i>	Dwivedi <i>et al.</i> (2010)
<i>GILT</i>	<i>Arabidopsis</i>	Canola	Wu <i>et al.</i> (2011)
<i>HaFAD2-1/HaAP10</i>	Sunflower	<i>Arabidopsis</i>	Zavallo <i>et al.</i> (2010)
<i>CCR</i>	<i>Eucalyptus gunnii</i>	Grapevine	Gago <i>et al.</i> (2011)
<i>MS</i>	Guar	<i>Medicago sativa</i>	Naoumkina and Dixon (2011)
<i>CsSUS1p-1/CsSUS1p-2</i>	<i>Citrus sinensis</i>	<i>Arabidopsis/tobacco</i>	Singer <i>et al.</i> (2011)
<i>Des</i>	Oil palm	Tomato	Saed <i>et al.</i> (2012)
<i>DX1</i>	Rice	Rice	Ye <i>et al.</i> (2012)
<i>Os03g01700/Os02g37190</i>	Rice	Rice	Li <i>et al.</i> (2013b)
<i>tCUP1</i>	Tobacco	Rice	Zhou <i>et al.</i> (2013)
<i>RIP1</i>	Tomato	Tomato/ <i>Arabidopsis</i>	Agarwal <i>et al.</i> (2017)
<i>PZmBD1/PZmDef1/PZmDef2</i>	Maize	Maize	Liu <i>et al.</i> (2016)
<i>AtMYB60</i>	<i>Arabidopsis</i>	<i>Arabidopsis/tobacco/tomato/rice</i>	Cominelli <i>et al.</i> (2011), Meyer <i>et al.</i> (2010) Oh <i>et al.</i> (2005) and Rusconi <i>et al.</i> (2013)
<i>CYP86A2</i>	<i>Arabidopsis</i>	<i>Arabidopsis</i>	Francia <i>et al.</i> (2008)
<i>GCI</i>	<i>Arabidopsis</i>	Tobacco	Wang <i>et al.</i> (2014)
<i>NRRS</i>	<i>Arabidopsis</i>	<i>Arabidopsis</i>	Kakrana <i>et al.</i> (2017)

cis-acting elements like dehydration responsive element (DRE) (Yamaguchi-Shinozaki and Shinozaki 2006), an abscisic acid-responsive element (ABRE) (Bonetta and Mccourt 1998), and heat shock element (HSE) have been identified. The necessity of such a system for transgene expression may be vital to

develop promoter set-ups where precise temporal regulation of transgene expression is a requisite. Examples include situations where unwanted gene expression is harmful or lethal, especially during the development of the plant (Guo *et al.* 2003). For example, peanut transgenics overexpressing *AtDREB1A* transcription factor under the influence of the *CaMV35S* promoter showed severe stunting that could be overcome by using the *rd29A* stress-inducible promoter to drive the expression of *DREB1A* (Bhatnagar-Mathur *et al.* 2007). Overexpression of a stress-related gene under an inducible promoter may confer better tolerance to stress than when under a constitutive one, while causing minimal or no growth retardation (Nakashima *et al.* 2007; Li *et al.* 2013a, b, c, d, e). The growth and development of *CaMV35S*-TaEXPB23 transgenic tobacco plants were altered under normal conditions, with a faster growth rate at the seedling stage, early flowering and maturation, and a shorter plant height compared to wild-type plants. On the other hand, RD29A-TaEXPB23 transgenic tobacco plants exhibited greater tolerance to water stress than the wild-type. Therefore, the use of stress-inducible promoters, such as *asrd29A* may minimize the negative effects of constitutive transgene expression and improve the water-stress tolerance of plants (Shen *et al.* 2003; Bhatnagar-Mathur *et al.* 2007; Li *et al.* 2013a, b, c, d, e). These studies infer that stress-inducible promoters are superior for enhancing plant productivity under stress conditions in comparison with that of the constitutive promoters. However, it must be ensured that the stress-inducible promoter expression is very tightly controlled in response to the desired stress to avoid any unintended effects on plant growth and development.

Rd29X promoters have been demonstrated to be useful in controlling targeted transgenes to mitigate abiotic stress in soybean (Bihmidine *et al.* 2013) and peanut (Bhatnagar-Mathur *et al.* 2007; Rao *et al.* 2017). However, the usefulness of the endogenous inducible promoters is often limited since they are leaky. There may be a possibility for the activation of other endogenous genes by the same inducer. Srivastava *et al.* 2014 reported that *PsSEOF1* promoter from pea could serve as an important candidate for tissue-specific promoter for engineering plants for both biotic and abiotic stress conditions. *HsfB2c* and *PM19* promoters from rice are highly heat-inducible and further characterization and reorganization of *cis*-acting elements in their promoters could lead to the development of highly effective, heat-inducible promoters (Reksiri *et al.* 2013). It was noticed that *PR10* promoter from *Erianthus* though highly constitutive, was quickly

induced upon wounding as well as on treatment with ABA and methyl jasmonate (Chakravarthi *et al.* 2016). The *OsABA2* promoter was shown to drive a low constitutive transgene expression under normal conditions but high induction in response to ABA, salt and drought stresses (Rai *et al.* 2009). The *ALSAP* promoter from the halophyte grass *Aeluropus litoralis* directs a stress-inducible expression pattern in transgenic rice plants making it an interesting candidate for engineering abiotic stress tolerance in cereals (Ben-Saad *et al.* 2015). Bang *et al.* 2012 similarly demonstrated that *OsNCED3* promoter was stress-inducible in whole rice plant except in the aleurones and endosperm and stably active over three generations. Stress-inducible expression of *Hsc70*, *Lea*, *Hsp10*, *Dhn* and *Apx* promoters from *Pennisetum glaucum* have been shown to confer abiotic stress tolerance in different tissues (leaf, stem and root) of the transgenic tobacco plants when exposed to different abiotic stresses (Divya *et al.* 2019; 2020, Divya *et al.* unpublished data). Hou *et al.* 2016 demonstrated that maize *Type-II H⁺-pyrophosphatase* promoter has higher expression under drought and salinity conditions compared to the *CaMV35S* promoter. The *AtUSP* promoter is highly inducible by phytohormones as well as multiple abiotic stresses that can be exploited as a stress-inducible promoter to develop multi-stress tolerant crops with least adverse effects on other important traits (Bhuria *et al.* 2016).

Promoter like the *AhMTP1* having a variety of *cis*-acting elements, particularly the MYB-binding sites are involved in the evolution of zinc tolerance (Fasani *et al.* 2017). The three cadmium-inducible gene promoters such as *OsGSTU5*, *OsGSTU37* and an *OsHsp20* from rice could be potentially used for bio-environmental contamination and improving heavy metal tolerance in crops (Qiu *et al.* 2015). Recent study indicated that *pGAL-2kb* could be a useful in developing drought-tolerant cultivars by driving transgene expression (Conforte *et al.* 2017). *CcHyPRP* promoter in *Arabidopsis* has been shown to be regulated by different stress factors which can be deployed for enhancing abiotic stress tolerance in transgenics (Srinath *et al.* 2017). *ZmPIS* and *PZ7* promoter fragments in tobacco would be ideal candidates for overexpression of drought- and salinity-responsive genes to improve crop tolerance (Zhang *et al.* 2016a, b).

Regulated gene expression systems would also be valuable in GE applications such as conditional expression of herbicide, flowering and fruit ripening genes that emphasize the need for developing transgenics by deploying inducible promoter systems (Liu *et al.* 2013). Myb-related protein-like promoter from

rice showed ethanol-inducible promoter activity and could be used to generate transgenic crops with desirable traits as demonstrated by Khanthapok *et al.* (2018). Afforementioned studies indicate that a wide spectrum of stress-inducible promoters exist, however their effectiveness and relative performance under multiple stress conditions in different taxa needs to be investigated. Recent study by Divya *et al.* (2019) it's observed that *PgApx* and *PgDhn* promoters are upregulated in drought, heat, cold and salt stresses and *PgHsc70* promoter is active in heat and drought stress. In another study, *PgHsp10* promoter from same plant i, e., *Pennisetum glaucum* seen to be expressed in heat and drought stress in transgenic tobacco plants (Divya *et al.* 2020). Different stress-inducible promoters used in varied studies so far are listed in table 5.

2.1.4 Synthetic: Recent advances in plant promoter engineering are making strides to generate more constitutive, bidirectional or inducible synthetic promoters for a proper transcriptional modulation of transgene expression in plants. Synthetic promoters provide enormous advantages over their counterparts with respect to transgene expression at a specific developmental stage, strength and tissue specificity. Synthetic promoters can be rationally sketched and constructed using specific type, copy number and positioning of motifs upstream of synthetic or native core promoters. To date, most synthetic promoters tested were either hybrids of multiple promoter parts or fusions of specific *cis*-regulatory elements with a core promoter. Typically, synthetic motif sequence is derived from extant sequences that are multiplied or recombined. The selection, copy number and spacing of *cis*-elements ascertain the strength, temporal and spatial expression patterns of synthetic promoters. Selection of motifs with known functions can be conducted with the help of previous studies (Banerjee *et al.* 2015) or from the databases, synthetic motif library screening (Roccaro *et al.* 2013) and bioinformatics based *de novo* motif discovery (Tompa *et al.* 2005). As the deconstructive analysis of plant natural promoters for functional motif discovery has slowly increased and the effectiveness of using systems biology tools for *denovo* motif discovery has been experimentally demonstrated in *Arabidopsis* (Koschmann *et al.* 2012) and soybean (Liu *et al.* 2014). Once motifs of interest have been selected for synthetic promoter construction, copy number and spacing of motif need to be optimized. Usually, motif copy number often correlates with synthetic promoter strength which has been demonstrated in various plant species like rice (Wu *et al.*

Table 5. List of stress-inducible promoters studied in transgenic plants

Promoter	Source	Host	References
<i>ABA2/ rab16A, HP1</i>	Rice	Rice	Rai <i>et al.</i> (2009)
<i>DREBa</i>	<i>Chrysanthemum dichrum</i>	<i>Arabidopsis</i>	Chen <i>et al.</i> (2012)
<i>Rab16A</i>	Rice	Rice	Ganguly <i>et al.</i> (2011)
<i>NCED3</i>	Rice	Rice	Bang <i>et al.</i> (2012)
<i>HsfB2cp, PM19p, Hsp90p</i>	Rice	Rice	Rerksiri <i>et al.</i> (2013)
<i>DXS, GGPPS</i>	<i>Ginkgo biloba</i>	-	Xu <i>et al.</i> (2013)
<i>hsp82</i>	Rice	Rice	Company <i>et al.</i> (2014)
<i>SOS1-AB</i>	<i>Triticum aestivum</i>	<i>Arabidopsis</i>	Feki <i>et al.</i> (2015)
<i>SEOF1</i>	Pea	Tobacco	Srivastava <i>et al.</i> (2014)
<i>SEOF1</i>	<i>Pisum sativum</i>	Tobacco	Srivastava <i>et al.</i> (2014)
<i>PR10</i>	<i>Erianthus arundinaceus</i>	Tobacco/ rice/ sugarcane	Chakravarthi <i>et al.</i> (2016)
<i>DREB1</i>	Buckwheat	Tobacco	Fang <i>et al.</i> (2016)
<i>ERF3</i>	Soybean	Soybean/ tobacco	Hernandez-Garcia and Finer (2016)
<i>UGT71C5</i>	<i>Arabidopsis</i>	<i>Arabidopsis</i>	Liu <i>et al.</i> (2015)
<i>GAPP</i>	Maize	Tobacco	Hou <i>et al.</i> (2016)
<i>pGAL</i>	Soybean	<i>Arabidopsis</i>	Conforte <i>et al.</i> (2017)
<i>HyPRP</i>	Cajanus cajan	<i>Arabidopsis</i>	Srinath <i>et al.</i> (2017)
<i>SnRK2.7</i>	<i>Triticum aestivum</i>	<i>Arabidopsis</i>	Wang <i>et al.</i> (2018)
<i>Myb-related Protein</i>	<i>Oryza sativa</i>	<i>Oryza sativa</i>	Khanthapok <i>et al.</i> (2018)
<i>PgAPX</i>	<i>Pennisetum glaucum</i>	Tobacco	Divya <i>et al.</i> (2019)
<i>PgDhn</i>	<i>Pennisetum glaucum</i>	Tobacco	Divya <i>et al.</i> (2019)
<i>PgHsc70</i>	<i>Pennisetum glaucum</i>	Tobacco	Divya <i>et al.</i> (2019)
<i>PgHsp10</i>	<i>Pennisetum glaucum</i>	Tobacco	Divya <i>et al.</i> (2020)

1998), tobacco (Sawant *et al.* 2005) and *Arabidopsis* (Sahoo *et al.* 2014). The motif dosage effect in synthetic promoters is not surprising, since congruent

findings have been observed in native promoters like in *Malus* (Espley et al. 2009) and *Arabidopsis* (Cao et al. 2014). When many motifs are accommodated in to a single synthetic promoter, spacing among motifs should be proper that is required for the hierarchical arrangement of their corresponding TFs in order to obtain full synergistic interactions with the RNA polymerase II complex (Sawant et al. 2005). Interestingly, two copies of the ACGT motif in synthetic promoters was shown to result in salicylic acid inducibility when separated by five nucleotides, but were ABA-inducible when separated by 25 nucleotides (Mehrotra and Mehrotra 2010).

The parsley (*Petroselinum crispum*) protoplast system was used for analyzing MAMP-responsive synthetic promoters which can be used by other plant systems to respond in cases of microbial pathogen attack (Kanofsky et al. 2016). Li et al. (2013a) reported that the synthetic promoter pCL made from *Arabidopsis* and potato was found optimal use for gene function research in potato tubers in response to low temperature. Altering the structure of CRT/DRE enhanced the CBF-associated transcription complex formation and thus improved the activity of this pCL, synthetic tuber-specific and cold-inducible promoter (Li et al. 2015). Comparative studies amongst the three natural promoters from rice (*Rab16A*) and two synthetically designed promoters, viz., 4X ABRE (abscisic acid-responsive element) having four tandem repeats of ABRE, and 2X ABRC (abscisic acid-responsive complex) having two tandem repeats of ABRE and two copies of coupling elements showed that 2XABRC make a better salinity/ABA-inducible promoter. The studies of Ganguly et al. (2011) indicated strong *GUS* expression in the whole seed (both embryo and aleurone layer of endosperm) only by 2X ABRC, while it was localized in the embryo for the other two promoters. Wang et al. (2015) demonstrated the synthesis of rice tissue-specific promoters and also developed a novel, feasible method for screening as well as for functional characterization of tissue-specific *cis*-acting elements with their flanking sequences at the genome-wide level in rice. This synthetic promoter including the 35S core sequence and two binding sites for cold-inducible CBF transcription factors (PDRE::35S) exhibited transient expression under chilling conditions using synthetic cold-inducible promoter which enhanced the target protein accumulation, and may decrease greenhouse heating expenses (Gerasymenko and Sheludko 2017). Synthetic *SynP16* promoter designed from *cis*-motifs of soybean viz. *ABF*, *ABRE*, *ABRE-Like*, *CBF*, *E2F-VARIANT*, *G-box*, *GCC-Box*,

MYB1, *MYB4*, *RAV1-A*, and *RAV1-B* (in multiple copies and various combination) with a minimal 35S core promoter and a 222 bp synthetic intron sequence. This *synp16* promoter induced *GUS* expression in stress inducible manner and tissues specifically in transgenic soybean and *Arabidopsis* (Jameel et al. 2020). Thus, synthetic promoters that are rationally designed have become highly efficient components for precise regulation of target gene expression. Different synthetic promoters used thus far are shown in table 6.

2.2 RNA pol III promoters

RNA polymerase III (pol III) has been used for transcription of structural and catalytic RNAs including 5S rRNA, tRNA and most small nuclear RNAs. Promoters for pol III have diverse structural features including reinitiation property and transcribing of at least four different types of genes, such as tRNA, 5S rRNA, U6 snRNA, 7SL-Sc-RNA, 7SK RNA. Thus *pol III* transcribes housekeeping genes required at all times. Different types of *pol III* promoters such as type 1 (5S RNA), type 2 (tRNA *Leu*) promoters of the *Xenopus laevis*, type 3 promoter of the *Homo sapiens* U6 snRNA gene and the type 4 Sc type of the *Saccharomyces cerevisiae* promoters were noticed. Other *pol III* promoters are *U3* and *U6* which are used in monocots and dicots, respectively (Belhaj et al. 2013). *Pol III* promoters are generally used to express small RNAs, short hairpin RNA, and guide RNA in the CRISPR/Cas9 system adapted for genome editing. *Pol III* promoters have not been much characterized excepting in model plants. Also, their relative advantage over that of other promoters need to be explored further.

3. Promoters in genome editing (CRISPR/Cas9 system)

Genome editing is a type of GE where the specific DNA sequence can be inserted, deleted or replaced. This can be accomplished by introducing site-specific double-strand breaks (DSBs) at specific sites in the targeted genome. The induced DSBs are repaired either through non-homologous end joining (NHEJ) or homologous recombination (HR), resulting in specific and targeted mutations. Currently, four families of engineered nucleases are being used: meganucleases (MGNs), zinc finger nucleases (ZFNs), transcription activator-like effector-based nucleases (TALEN), and the clustered regularly interspaced short palindromic

Table 6. Synthetic promoters that have been designed for use in plant systems

Promoter	Type	Source	Host	References
<i>pCL</i>	Tuber-specific and cold-inducible	At cor15a promoter region and potato patatin promoter region	Potato	Li <i>et al.</i> (2013a)
<i>Sab/sba</i>	Cold-inducible	At cor15a promoter regions and cor15b promoter regions	Tobacco	Li <i>et al.</i> (2013a)
<i>4 x CCTC</i>	Fungal colonization under low-Pi condition-inducible	Potato Piransporter 3 (StPT3) promoter regions; 35S core promoter	Potato/Lotus	Lota <i>et al.</i> (2013)
<i>4xRE/B4REA</i>	Hormonal and bacterial pathogen inducible	Hormone-response elements; 35S core promoter	Tobacco, <i>Arabidopsis</i>	Liu <i>et al.</i> (2014)
<i>FsFfCBD</i>	Bidirectional	FsCP; 35S core promoter; a tri-hybrid enhancer FsEFfECE	Tobacco	Patro <i>et al.</i> (2013)
<i>4 x GCC</i>	Jasmonic acid inducible	AtPDF1.2 promoter; 35S core promoter	<i>Arabidopsis</i>	Van der Does <i>et al.</i> (2013)
<i>4 x ROSE1 ~ 7</i>	ROS-inducible	AtROS-responsive elements; 35S core promoter	<i>Arabidopsis</i>	Wang <i>et al.</i> (2013)
<i>FSgt-PFIt;MSgt-PFIt;PFIt-UAS-2X</i>	Constitutive	FSgt, Msgt, PFIt-UAS; PFIt core promoter (PFIt)	Tobacco, <i>Petunia</i> , <i>Arabidopsis</i> , Tomato, Spinach	Acharya <i>et al.</i> (2014)
<i>4 x RSRE</i>	Stress-inducible	<i>Arabidopsis</i> rapid stress response elements; NOS core promoter	<i>Arabidopsis</i>	Benn <i>et al.</i> (2014)
<i>p35S-PCHS-Ω;p35S-Ω;LCHSΩ;pOCS-PCHS-Ω;pOCS-LCHS-Ω</i>	Flower-specific	35S or OCS enhancer; petunia CHSA core promoter; lily CHS core promoter, an omega element	<i>Toreniafournier</i>	Du <i>et al.</i> (2014)
<i>CL</i>	Tuber-specific and cold-inducible	<i>Arabidopsis</i> cor15a promoter region and potato patatin promoter region; 7 nucleotides mutation in the 5' and 3' flanking sequences of CRT/DRE.	Potato	Li <i>et al.</i> (2015)
<i>MAMP-responsive synthetic promoter P_DRE::35S</i>	Microbial pathogen attack chilling temperatures	Four copies of a potential MAMP-responsive cis-sequence two binding sites for CBF transcription factors (CRT/DRE), CaMV35S promoter core sequence and 5-leader sequence of TMV omega	<i>Petroselinum crispum</i> <i>N. excelsior</i>	Kanofsky <i>et al.</i> (2016) Gerasymenko and Sheludko (2017)
<i>SynP16</i>	Soybean	Soybean <i>ABF</i> , <i>ABRE</i> , <i>ABRE-Like</i> , <i>CBF</i> , <i>E2F-VARIANT</i> , <i>G-box</i> , <i>GCC-Box</i> , <i>MYB1</i> , <i>MYB4</i> , <i>RAV1-A</i> , and <i>RAV1-B</i> (in multiple copies and various combination) with a minimal 35s core promoter and a 222 bp synthetic intron sequence	Soybean, <i>Arabidopsis</i>	Aysha <i>et al.</i> (2020)

repeats (CRISPR). While the MGNs and ZFNs were costly and hard to engineer, TALE nucleases are flexible, specific and relatively complex and CRISPR/Cas9 nucleases are user-friendly and cost-effective (Belhaj *et al.* 2013). The CRISPR/Cas9 system derived from bacterial immune system and has been widely used for targeted genome editing in diverse organisms including mammalian and most of the plant species (Hsu *et al.*

2014; Ma *et al.* 2015; Guo *et al.* 2018). The CRISPR/Cas9 system uses either single or multi-guide RNA system to induce the specific edits and this efficiency depends on the endonuclease activity of the sgRNA/Cas9 complex and selection of the promoters (Guo *et al.* 2018). It has been observed that targeted efficiency of the CRISPR/Cas9 depends on the codon optimization of the Cas9 and the promoters. Targeted

efficiency increases when expressing *Cas9* gene other than *CaMV35S* promoter such as the dividing cell specific *INCURVATA2* (Hyun et al. 2015), egg cell-specific promoters (Wang et al. 2015), the cell division-specific *YAO* (Yan et al. 2015), and the germ-line-specific *SPOROCTELESS* (Mao et al. 2016) and embryo-specific promoter DD45 (Miki et al., 2018). Targeted efficiency increased significantly when *ZmUbi1* promoter was used instead of *CaMV35S* (Feng et al. 2018). Feng et al. (2018) reported that *Zmdmcl* promoter could be an alternative to the *CaMV35S* for expressing the CRISPR/Cas9 system in maize for generating highly efficient targeted genome editing.

The *pol III* promoters such as *U3* and *U6* are commonly used to express the small RNAs and guide RNAs of the CRISPR/Cas9 system. Due to their long-term regulation of target genes and their defined sites for transcription initiation and termination, *U3* and *U6* promoters gained special attention over the others. Characterization of *pol III* promoters in many organisms was not properly explored and this becomes difficult in choosing the right promoters for CRISPR/Cas9 gene edited targeted mutagenesis. Hence, in most of the studies, *pol III* promoters such as *U3* or *U6* from *Arabidopsis* or rice or maize or wheat are often chosen to drive gRNAs (Nekrasov et al. 2013; Bortesi and Fischer 2015; Feng et al. 2018). Maize *ubi1* gene promoter in combination with two rice *U6* promoters performed well in maize with mutation efficiency up to 70% (Char et al. 2016). Processing of a *pol II* transcript into functional gRNA has been found successful using ribozyme or Csy4RNA cleavage systems. Mikami et al. (2017) have shown that functional gRNAs can be efficiently processed using SpCas9 protein and plant RNA cleavage systems without any need for a specific RNA processing. Recent studies have shown that *AtU6* promoter from *Arabidopsis* and *Fve U6* promoter of wild strawberry are equally good for the high-efficiency genome editing (Zhou et al. 2013). Hashimoto et al. (2018) demonstrated increased efficiency and also improved multiplex genome editing using *Cas9* gene expression with *SIEF1 α* promoter. Ordon et al. (2020) demonstrated that *DD45* and *RPS5a* promoter-driven Cas9 showed higher mutational frequency. Promoters used for genome editing are shown in the table 5.

4. Bidirectional promoters (BDPs)

A synthetic bidirectional expression module was prepared by placing a computationally designed minimal promoter sequence at the 5' and 3' sides of a

transcription activation module. BDPs were identified in diverse organisms including plants. Structural features and functional consequences of BDPs received much attention in plant biotechnology due to their effective usage in gene-stacking/pyramiding and targeting the complex traits (Que et al. 2010). However, the underlying mechanisms responsible for the bidirectional transcription and co-expression of BDPs has remained poorly understood in plants where they drive the transcription of the two target genes (Patro et al. 2013). Such promoters play a pivotal role in the transcription of bidirectional gene pairs where the two genes are positioned head-to-head on opposite strands of DNA (figure 2c). The BDPs efficiently regulate two transgenes simultaneously which can be evaluated in multiple model-plant systems. The BDPs representing *Rep* and *coat protein (CP)* genes of CLCuBuV were characterized and their efficacy assayed (Khan et al. 2015). They also showed that the strong constitutive CLCuBuV *Rep* promoter could be of use for higher expression of transgenes in a variety of plant cells. Wang et al. (2016a, b) combined RNA-seq data and cDNA microarray data to discover the potential BDPs in rice genome. Comparative analysis between *ZmBD1*, *ZmDef1* and *ZmDef2* promoters revealed that *PZmBD1* shared most of the expression characteristics of the two polar promoters, but displayed more stringent embryo specificity, delayed expression initiation, and asymmetric promoter activity (Liu et al. 2016).

The initiation of transcription from both the unidirectional and bidirectional promoters made from the same sequence elements were evaluated by using the *uidA* and *gfp* reporter genes. The investigation based on transient and stable transformation of tobacco exhibited that the artificially designed multifactorial activation module activated the transcription simultaneously to proportionate levels in both the directions (Chaturvedi et al. 2006). The transcription regulatory module responded to elicitors like salicylic acid (SA), NaCl and IAA in the forward as well as reverse directions. It implied that constitutive and chemically-inducible bidirectional promoters can be deployed for predictable simultaneous regulation of two genes for genetic engineering in plants (Chaturvedi et al. 2006). Fang et al. 2016 elucidated the unique epigenetic mechanism of BDPs and regulation of the bidirectional gene pairs and eventually used for GE. The bidirectional promoters *At4g35985* (P85) and *At4g35987* (P87) in both orientations display up-regulation under salt stress. Such regulatory elements of BDPs showing spatial and stress-inducible promoter and functioning in heterologous systems might be an important tool for

Table 7. List of the promoters that have been designed for use in CRISPR/Cas9 genome editing

Species	Cas9 codon optimization	Promoters for Cas9	Promoter for gRNA	Target gene	Reference
<i>A. thaliana</i>	Human	<i>AtUBQ1</i>	<i>AtU6</i>	<i>GUUS</i>	Mao <i>et al.</i> (2016)
<i>A. thaliana</i>	<i>Arabidopsis</i>	<i>PcUBI4-2</i>	<i>AtU6</i>	<i>GUUS, UGUS</i>	Fausser <i>et al.</i> (2014)
<i>A. thaliana</i>	<i>Arabidopsis</i>	<i>PcUBI4-2</i>	<i>AtU6</i>	<i>GUUS, UGUS</i>	Fausser <i>et al.</i> (2014)
<i>A. thaliana</i>	Human	<i>2xCaMV35S</i>	<i>AtU6</i>	<i>YFFP</i>	Feng <i>et al.</i> (2013)
<i>A. thaliana</i>	<i>Arabidopsis</i>	<i>PcUBI4-2</i>	<i>AtU6</i>	<i>ADH1</i>	Schimpl <i>et al.</i> (2014)
<i>N. benthamiana</i>	<i>Arabidopsis</i>	<i>CaMV35SPDK</i>	<i>AtU6</i>	<i>NbPDS3</i>	Li <i>et al.</i> (2013a, b, c, d, e)
<i>O. sativa</i>	Rice	<i>2xCaMV35S</i>	<i>OsU3</i>	<i>OsPDS, OsBADH2</i>	Shan <i>et al.</i> (2013)
<i>O. sativa</i>	Rice	<i>ZmUbi</i>	<i>OsU3</i>	<i>GUUS</i>	Mao <i>et al.</i> (2016)
<i>A. thaliana</i>	Plant	<i>35S PPKK</i>	<i>AtU6</i>	<i>AtPDS3, AtFL2, AtRACK1b, AtRACK1c</i>	Li <i>et al.</i> (2013a, b, c, d, e)
<i>A. thaliana</i>	<i>Arabidopsis</i>	<i>ICU2</i>	<i>AtU6</i>	<i>AtFT, AtSPL4</i>	Hyun <i>et al.</i> (2015)
<i>N. benthamiana</i>	Human	<i>CaMV35</i>	<i>PEBV</i>	<i>NbPDS, NbPCNA</i>	Ali <i>et al.</i> (2015)
<i>G. max</i>	Soybean	<i>EF1A2</i>	<i>GmU6</i>	<i>GmDD20, GmDD43</i>	Li <i>et al.</i> (2015)
<i>Z. mays</i>	Human	<i>ZmDMC1</i>	<i>ZmU3</i>	<i>Zmzb7</i>	Feng <i>et al.</i> (2018)
<i>Gossypium hirsutum</i>		<i>2xP35s</i>	<i>GhU6.3, AtU6-29</i>	<i>GUS</i>	Long <i>et al.</i> (2018)
<i>A. thaliana</i>		<i>CaMV35s</i>	<i>AtU6-26</i>	<i>AtM20</i>	Ma <i>et al.</i> (2018)
Tomato, <i>N. benthamiana</i>	Human	<i>CaMV35s</i>	<i>U6-26s</i>	<i>CP, Rep</i>	Tashkandi <i>et al.</i> (2018)
<i>Fragaria vesca</i>	<i>Arabidopsis</i> and maize	<i>AtUBQ10, CaMV35s</i>	<i>FveU6-2, AtU6-26</i>	<i>TAA1, ARF8</i>	Zhou <i>et al.</i> (2013)
<i>Solanum lycopersicum</i>	<i>Arabidopsis</i>	<i>SIEF1α, Slp16, Pcubi4, 2xCaMV35S</i>	<i>AtU6-26</i>	<i>SINADK2A</i>	Hashimoto <i>et al.</i> (2018)
<i>A. thaliana</i>	<i>Arabidopsis</i>	<i>DD45, RPS5a</i>		<i>Lhcb1</i>	Ordon <i>et al.</i> (2020)

plant biotechnology and gene stacking applications (Banerjee *et al.* 2013). Recently, Araceli *et al.* (2017) have shown that BDPs function can be regulated by degree/intensity of the abiotic stresses like Pi availability. *GhZU* promoter from *Gossypium hirsutum* is seen to regulate the expression of *gus* and *gfp* genes in both the directions in transgenic *Arabiposis* plants (Yang *et al.* 2018). From these studies, it can be stated that BDPs have a potential role to play in plant genetic engineering (Table 7).

5. Conclusions and future perspectives

The selection of a promoter suitable for the targeted transgene expression is one of the most important criteria for developing the GM crops. There are some important implications for producing transgenic plants with higher yield without any compromise on environmental and biosafety concerns. Improved expression of transgenes has been carried by hybrids or combined promoters that are from the constitutive promoters. Constitutive promoters are often seen to be beneficial for a high-level expression of

selectable marker genes, necessary for efficient selection and generation of transgenics. Constitutively active promoters are not necessarily always desirable for GE plants as the constitutive overexpression of a transgene may compete for energy and building blocks for synthesis of proteins, RNA and others. Transgenic plants constitutively expressing TFs have been shown to exhibit mild/severe growth retardation in the aerial parts, shorter petioles, rounder leaves, delayed flowering and a dwarf phenotype. In these situations, use of tissue-specific or stress-inducible promoters of moderate strength may be more desirable.

Tissue-specific and stress-inducible promoters can be exploited to minimize the unintended effects of adverse environmental conditions such as heat, cold, drought, and salinity. Inducible promoters are very powerful resources for GE plants, since the expression of genes operably linked to them can be regulated to function at certain stages of growth and development of an organism or a particular tissue which help in reduction of energy expenditure of the plant. Further, inducible promoters will be activated only when required, thereby maintaining yield or productivity even under adverse conditions. On the other hand, synthetic

promoters paired with synthetic TFs can be used to provide a coordinated transcriptional control of multiple genes, which would be needed for successful metabolic engineering.

Genome-wide computational prediction and analysis of core promoter elements across plant monocots and dicots will be useful for future work related to genome annotation projects and can inspire research efforts aimed to better understand regulatory mechanisms of transcription. Genome-wide transcriptomic analysis such as RNA-seq will also help in gene regulation as well as in the identification of promoters and *cis*-acting elements. Advancement in the area of systems biology and the development of new tools would be beneficial in the discovery of novel promoters and their *cis*-acting elements. Identification and deployment of promoters that can help under diverse stress conditions are desirable, ultimately for the generation of transgenics with better survival and productivity. CRISPR technology has become popular and an alternative to the plant breeding techniques. Genome editing efficiency can be improved by optimizing the diverse range of the promoters. Hence, crop varieties developed by above mentioned technologies are considered as non-GM in USA and other regions. Relaxing in usage of such crop varieties from the scope of the GMO legislation may have a positive impact on the development of the plant biotechnology and breeding sector for the betterment of humans.

Acknowledgements

PSR acknowledges financial support from the Department of Science and Technology, Govt. of India, New Delhi, for a fellowship and research grant through the INSPIRE Faculty Award No. IFALSPA-06 and Young Scientist Award SB/YS/LS-12/2013. This work was undertaken as part of the CGIAR Research Program on Grain Legumes & Dryland Cereals (CRP-GLDC).

References

Acharya S, Ranjan R, Pattanaik S, Maiti IB and Dey N 2014 Efficient chimeric plant promoters derived from plant infecting viral promoter sequences. *Planta* **239** 381–396

Agarwal P, Kumar R, Pareek A and Sharma AK 2017 Fruit preferential activity of the tomato RIP1 gene promoter in transgenic tomato and Arabidopsis. *Mol. Genet. Genom.* **292** 145–156

Ali Z, Abulfaraj A, Idris A, Ali S, Tashkandi M and Mahfouz MM 2015 CRISPR/Cas9-mediated viral interference in plants. *Genome Biol.* **16** 238

Althoff F, Kopischke S, Zobell O, Ide K, Ishizaki K, Kohchi T and Zachgo S 2014 Comparison of the MpEF1alpha and CaMV35 promoters for application in *Marchantia polymorpha* overexpression studies. *Transgenic Res.* **23** 235–244

An YQ, McDowell JM, Huang S, McKinney EC, Chambliss S and Meagher RB 1996 Strong, constitutive expression of the *Arabidopsis* ACT2/ACT8 actin subclass in vegetative tissues. *Plant J.* **10** 107–121

Annadana S, Beekwilder MJ, Kuipers G, Visser PB, Outchkourov N, Pereira A, Udayakumar M, De Jong J and Jongsma MA 2002 Cloning of the chrysanthemum UEP1 promoter and comparative expression in florets and leaves of *Dendranthema grandiflora*. *Transgenic Res.* **11** 437–445

Araceli OA, Alfredo CR, Javier MM and Luis HE 2017 A phosphate starvation-driven bidirectional promoter as a potential tool for crop improvement and in vitro plant biotechnology. *Plant Biotechnol. J.* **15** 558–567

Arango J, Salazar B, Welsch R, Sarmiento F, Beyer P and Al-Babili S 2010 Putative storage root specific promoters from cassava and yam: cloning and evaluation in transgenic carrots as a model system. *Plant Cell Rep.* **29** 651–659

Atkinson RG, Bolitho KM, Wright MA, Iturriagoitia-Bueno T, Reid SJ and Ross GS 1998 Apple ACC-oxidase and polygalacturonase:ripening-specific gene expression and promoter analysis in transgenic tomato. *Plant Mol. Biol.* **38** 449–460

Bang SW, Park SH, Jeong JS, Kim YS, Jung H, Ha SH and Kim JK 2012 Characterization of the stress-inducible OsNCED3 promoter in different transgenic rice organs and over three homozygous generations. *Planta* **237** 211–224.

Banerjee J, Sahoo DK, Dey N, Houtz RL and Maiti IB 2013 An intergenic region shared by At4g35985 and At4g35987 in *Arabidopsis thaliana* is a tissue specific and stress inducible bidirectional promoter analyzed in transgenic arabidopsis and tobacco plants. *PLoS One.* **8** e79622

Banerjee J, Sahoo DK, Raha S, Sarkar S, Dey N and Maiti IB 2015 A region containing an as-1 element of Dahlia Mosaic Virus (DaMV) subgenomic transcript promoter plays a key role in green tissue and root-specific expression in plants *Plant Mol. Biol. Rep.* **33** 532–556

Belhaj K, Chaparral-Garcia A, Amount S and Nekrasov V 2013 Plant genome editing made easy: targeted mutagenesis in model and crop plants using the CRISPR/Cas system. *Plant Methods* **9** 1–10

Benn G, Wang CQ, Hicks DR, Stein J, Guthrie C and Dehesh K 2014 A key general stress response motif is regulated non-uniformly by CAMTA transcription factors. *Plant J.* **80** 82–92

Ben-Saad R, Meynard D, Ben-Romdhane W, Mieulet D, Verdeil JL, Al-Doss A, Guiderdoni E and Hassairi A 2015

- The promoter of the AISAP gene from the halophyte grass *Aeluropus litoralis* directs a stress-inducible expression pattern in transgenic rice plants. *Plant Cell Rep.* **34** 1791–1806
- Benfey PN, Ren L and Chua NH 1990 Tissue-specific expression from CaMV35S enhancer subdomains in early stages of plant development. *EMBO J.* **9** 1677–1684
- Benfey PN, Takatsuji H, Ren L, Shah DM and Chua NH 1990 Sequence requirements of the 5-enolpyruvylshikimate-3-phosphate synthase 5[prime]-upstream region for tissue-specific expression in flowers and seedlings. *Plant Cell* **2** 849–856
- Beringer J, Chen W, Garton R, Sardesai N, Wang PH, Zhou N, Gupta M and Wu H 2017 Comparison of the impact of viral and plant-derived promoters regulating selectable marker gene on maize transformation and transgene expression. *Plant Cell Rep.* **36** 519–528
- Bhattacharyya J, Chowdhury AH, Ray S, Jha JK, Das S, Gayen S, Chakraborty A, Mitra J, Maiti MK, Basu A and Sen SK 2011 Native polyubiquitin promoter of rice provides increased constitutive expression in stable transgenic rice plants *Plant Cell Rep.* **31** 271–279
- Bhattacharyya S, Dey N and Maiti IB 2002 Analysis of cis-sequence of subgenomic transcript promoter from the Figwort mosaic virus and comparison of promoter activity with the cauliflower mosaic virus promoters in monocot and dicot cells. *Virus Res.* **90** 47–62
- Bhattacharyya S, Pattanaik S and Maiti IB 2003 Intron-mediated enhancement of gene expression in transgenic plants using chimeric constructs composed of the Peanut chlorotic streak virus (PCISV) promoter-leader and the antisense orientation of PCISV ORF VII (p7R). *Planta* **218** 115–24
- Bhatnagar-Mathur P, Devi MJ, Reddy DS, Lavanya M, Vadez V, Serraj R, Yamaguchi-Shinozaki K and Sharma KK 2007 Stress-inducible expression of At DREB1A in transgenic peanut (*Arachis hypogaea* L) increases transpiration efficiency under water-limiting conditions. *Plant Cell Rep.* **26** 2071–2082
- Bhuria M, Goel P, Kumar S and Singh AK 2016 The Promoter of AtUSP is co-regulated by phytohormones and abiotic stresses in *Arabidopsis thaliana*. *Front Plant Sci.* **7** 1957
- Bihmidine S, Lin J, Stone JM, Awada T, Specht JE and Clemente TE 2013 Activity of the *Arabidopsis* RD29A and RD29B promoter elements in soybean under water stress. *Planta* **23** 755–64
- Biłas R, Szafran K, Hnatuszko-Konka K and Kononowicz AK 2016 Cis-regulatory elements used to control gene expression in plants *Plant Cell Tiss. Org. Cul.* **127** 269–287
- Bonetta D and McCourt P 1998 Genetic analysis of ABA signal transduction pathways. *Trends Plant Sci.* **3** 231–235
- Borah BK, Zarreen F, Baruah G and Dasgupta I 2016 Insights into the control of geminiviral promoters. *Virology* **495** 101–111
- Bortesi L and Fischer R 2015 The CRISPR/Cas9 system for plant genome editing and beyond. *Biotechnol. Adv.* **33** 41–52
- Bucchini L and Goldman LR 2002 Starlink corn: a risk analysis *Environ Health Perspect* **110** 5
- Biłas R, Szafran K, Hnatuszko-Konka K and Kononowicz AK 2016 Cis-regulatory elements used to control gene expression in plants *Plant Cell Tiss. Org. Cul.* **127** 269–287
- Cahoon EB and Shanklin J 2000 Substrate-dependent mutant complementation to select fatty acid desaturase variants for metabolic engineering of plant seed oils. *Proc Nat Acad Sci.* **97** 12350–12355
- Cao S, Kumimoto RW, Gnesutta N, Calogero AM, Mantovani R and Holt BF 2014 A distal CCAAT/NUCLEAR FACTOR Y complex promotes chromatin looping at the FLOWERING LOCUS T promoter and regulates the timing of flowering in *Arabidopsis*. *Plant Cell* **26** 1009–1017
- Cazzonelli CI, McCallum EJ, Lee R and Botella JR 2005 Characterization of a strong constitutive mung bean (*Vigna radiata* L) promoter with a complex mode of regulation in planta. *Transgenic Res.* **14** 941–967
- Chakravarthi M, Syamaladevi DP, Harunipriya P, Augustine SM and Subramonian N 2016 A novel PR10 promoter from *Erianthus arundinaceus* directs high constitutive transgene expression and is enhanced upon wounding in heterologous plant systems. *Mol. Biol. Rep.* **43** 17–30
- Chaturvedi CP, Sawant SV, Kiran K, Mehrotra R, Lodhi N, Ansari SA and Tuli R 2006 Analysis of polarity in the expression from a multifactorial bidirectional promoter designed for high-level expression of transgenes in plants. *J. Biotechnol.* **123** 1231–1312
- Char SN, Neelakandan AK, Nahampun H, Frame B, Main M, Spalding M H et al (2017) An Agrobacterium-delivered CRISPR/Cas9 system for high-frequency targeted mutagenesis in maize. *Plant Biotechnol. J.* **15** 257–268.
- Chen Y, Chen S, Chen F, Li P, Chen L, Guan Z and Chang Q 2012 Functional characterization of a *Chrysanthemum dichrum* stress-related promoter. *Mol. Biotechnol.* **52** 161–169
- Chen L, Jiang B, Wu C, Sun S, Hou W and Han T 2014 GmPRP2 promoter drives root-preferential expression in transgenic *Arabidopsis* and soybean hairy roots. *BMC Plant Biol.* **14** 245
- Christensen AH, Sharrock RA and Quail PH 1992 Maize polyubiquitin genes: structure hermal perturbation of expression and transcript splicing and promoter activity following transfer to protoplasts by electroporation. *Plant Mol. Biol.* **18** 675–689

- Cominelli E, Galbiati M, Albertini A, Fornara F, Conti L, Coupland G and Tonelli C 2011 DOF-binding sites additively contribute to guard cell-specificity of AtMYB60 promoter. *BMC Plant Biol.* **11** 162
- Company N, Nadal A, Ruiz C and Pla M 2014 Production of phytotoxic cationic alpha-helical antimicrobial peptides in plant cells using inducible promoters. *PLoS One* **9** e109990
- Conforte AJ, Guimaraes-Dias F, Neves-Borges AC, Bencke-Malato M, Felix-Whipps D and Alves-Ferreira M (2017) Isolation and characterization of a promoter responsive to salt osmotic and dehydration stresses in soybean. *Genet. Mol. Biol.* **40** 226–237
- Coussens G, Aesaert S, Verelst W, Demeulenaere M, De Buck S, Njuguna E, Inze D and Van Lijsebettens M 2012 *Brachypodium distachyon* promoters as efficient building blocks for transgenic research in maize. *J. Exp. Bot.* **63** 4263–4273
- Dean GH, Jin Z, Shi L, Esfandiari E, McGee R, Nabata K et al 2017 Identification of a seed coat-specific promoter fragment from the Arabidopsis MUCILAGE-MODIFIED4 gene. *Plant Mol. Bio.* **95** 33–50
- Dhankher OP, Li Y, Rosen BP, Shi J, Salt D, Senecoff JF, Sashti NA and Meagher RB 2002 Engineering tolerance and hyperaccumulation of arsenic in plants by combining arsenate reductase and gamma-glutamylcysteine synthetase expression. *Nat. Biotechnol.* **20** 1140–1145
- Divya K, Kishor PBK, Bhatnagar-Mathur P, Singam P, Sharma KK, Vadez V and Reddy PS 2019 Isolation and functional characterization of three abiotic stress-inducible (Apx, DhN and Hsc70) promoters from pearl millet (*Pennisetum glaucum* L.) *Mol. Biol. Rep.* **46** 6039–6052
- Divya K, Bhatnagar-Mathur P, Sharma KK, Vadez V and Reddy PS 2020 Functional characterization of the promoter of pearl millet heat shock protein 10 (PgHsp10) in response to abiotic stresses in transgenic tobacco plants. *Int. J. Biol. Macromol.* **12** 103–110.
- Du L, Lou Q, Zhang X, Jiao S, Liu Y and Wang Y 2014 Construction of flower-specific chimeric promoters and analysis of their activities in transgenic *Torenia*. *Plant Mol. Biol. Rep.* **32** 234–245
- Dwivedi KK, Roche D and Carman JG 2010 Expression in Arabidopsis of a nucellus-specific promoter from watermelon (*Citrullus lanatus*). *Plant Sci.* **179** 549–552
- Erpen L, Tavano ECR, Harakava R, Dutt M, Grosser JW, Piedade SMS and Mourão Filho FAA 2018 Isolation characterization and evaluation of three Citrus sinensis-derived constitutive gene promoters. *Plant Cell Rep.* **37** 1113–1125
- Erpen-Dalla Corte L, Mendes BMJ, Mourão Filho FAA, Grosser JW and Dutt M 2020 Functional characterization of full-length and 5' deletion fragments of Citrus sinensis-derived constitutive promoters in *Nicotiana benthamiana*. *In Vitro Cell. Dev. Biol. Plant* **56** 280–289
- Espley RV, Brendolise C, Chagne D, Kuty-Amma S, Green S, Volz R, Putterill J, Schouten HJ, Gardiner SE, Hellens RP and Allan AC 2009 Multiple repeats of a promoter segment causes transcription factor autoregulation in red apples. *Plant Cell* **21** 168–183
- Fang RX, Nagy F, Sivasubramaniam S and Chua NH 1989 Multiple cis regulatory elements for maximal expression of the cauliflower mosaic virus 35S promoter in transgenic plants. *Plant Cell*, **1** 141–150
- Fang Y, Wang L, Wang XY, Ou Q, Pan X, Xiao J and Zhang W 2016 Histone modifications facilitate the coexpression of bidirectional promoters in rice. *BMC Genomics* **17** 768
- Fasani E, DalCorso G, Varotto C, Li M, Visioli G, Mattarozzi M and Furini A 2017 The MTP1 promoters from *Arabidopsis halleri* reveal cis-regulating elements for the evolution of metal tolerance. *New Phytol.* **214** 1614–1630
- Fausser F, Schiml S and Puchta H 2014 Both CRISPR/Cas-based nucleases and nickases can be used efficiently for genome engineering in Arabidopsis thaliana. *Plant J.* **79** 348–359
- Feki K, Brini F, Ben Amar S, Saibi W and Masmoudi K 2015 Comparative functional analysis of two wheat Na(+)/H(+) antiporter SOS1 promoters in *Arabidopsis thaliana* under various stress conditions. *J. Appl. Genet.* **56** 15–26.
- Feng Z, Zhang B, Ding W, Liu X, Yang DL, Wei P and Zhu JK 2013 Efficient genome editing in plants using a CRISPR/Cas system. *Cell Res.* **23** 1229
- Feng C, Su H, Bai H, Wang R, Liu Y, Guo X and Han F 2018 High-efficiency genome editing using a dmcl promoter-controlled CRISPR/Cas9 system in maize. *Plant Biotechnol. J.* **16** 1848–1857
- Forde BG, Heyworth A, Pywell J and Kreis M 1985 Nucleotide sequence of a B1 hordein gene and the identification of possible upstream regulatory elements in endosperm storage protein genes from barley wheat and maize. *Nucleic Acids Res.* **13** 7327–7339
- Francia P, Simoni L, Cominelli E, Tonelli C and Galbiati M 2008 Gene trap-based identification of a guard cell promoter in *Arabidopsis*. *Plant Signal. Beh.* **3** 684–686
- Fromm ME, Morrish F, Armstrong C, Williams R, Thomas J and Klein TM 1990 Inheritance and expression of chimeric genes in the progeny of transgenic maize plants. *Biotechnology (NY)* **8** 833–839
- Gago J, Grima-Pettenati J and Gallego PP 2011 Vascular-specific expression of GUS and GFP reporter genes in transgenic grapevine (*Vitis vinifera* L cv Albarino) conferred by the EgCCR promoter of *Eucalyptus gunnii*. *Plant Physiol. Biochem.* **49** 413–419
- Ganguly M, Roychoudhury A, Sarkar SN, Sengupta DN, Datta SK and Datta K 2011 Inducibility of three salinity/abscisic acid-regulated promoters in transgenic rice with gusA reporter gene *Plant Cell Rep.* **30** 1617–1625

- Gerasyenko IM and Sheludko YV 2017 Synthetic cold-inducible promoter enhances recombinant protein accumulation during *Agrobacterium*-mediated transient expression in *Nicotiana excelsior* at chilling temperatures *Biotechnol. Lett.* **39** 1059–1067
- Govindarajulu M, Elmore JM, Fester T and Taylor CG 2008 Evaluation of constitutive viral promoters in transgenic soybean roots and nodules. *Mol. Plant Microbe Interact.* **21** 1027–1035
- Guo HS, Fei JF, Xie Q and Chua NH 2003 A chemical-regulated inducible RNAi system in plants. *Plant J.* **34** 383–392
- Guo J, Li K, Jin L, Xu R, Miao K, Yang F and Miao Y 2018 A simple and cost-effective method for screening of CRISPR/Cas9-induced homozygous/biallelic mutants. *Plant Methods* **14** 40
- Gupta P, Raghuvanshi S and Tyagi AK 2001 Assessment of the efficiency of various gene promoters via biolistics in leaf and regenerating seed callus of millets *Eleusine coracana* and *Echinochloa crusgalli*. *Plant Biotech.* **18** 275–282
- Hadley CW, Miller EC, Schwartz SJ and Clinton SK 2002 Tomatoes lycopene and prostate cancer: progress and promise. *Exp. Biol. Med.* **227** 869–880
- Han YJ, Kim YM, Hwang OJ and Kim JI 2015 Characterization of a small constitutive promoter from *Arabidopsis* translationally controlled tumor protein (AtTCTP) gene for plant transformation. *Plant Cell Rep.* **34** 265–275
- Hannoufa A, Negruk V, Eisner G and Lemieux B 1996 The CER3 gene of *Arabidopsis thaliana* is expressed in leaves stems roots flowers and apical meristems. *Plant J.* **10** 459–467
- Hashimoto R, Ueta R, Abe C, Osakabe Y and Osakabe K 2018 Efficient multiplex genome editing induces precise and self-ligated type mutations in tomato plants. *Front. Plant Sci.* **9** 916
- Hernandez-Garcia CM and Finer JJ 2014 Identification and validation of promoters and cis-acting regulatory elements. *Plant Sci.* **217–218** 109–119
- Hernandez-Garcia CM and Finer JJ 2016 A novel cis-acting element in the GmERF3 promoter contributes to inducible gene expression in soybean and tobacco after wounding. *Plant Cell Rep.* **35** 303–316
- Hernandez-Garcia CM, Martinelli AP, Bouchard RA and Finer JJ 2009 A soybean (*Glycine max*) polyubiquitin promoter gives strong constitutive expression in transgenic soybean. *Plant Cell Rep.* **28** 837–849
- Hou J, Jiang P, Qi S, Zhang K, He Q, Xu C, Ding Z, Zhang K and Li K 2016 Isolation and functional validation of salinity and osmotic stress inducible promoter from the maize Type-II H⁺-Pyrophosphatase gene by deletion analysis in transgenic tobacco Plants. *PLoS One* **11** e0154041
- Hsu PD, Lander ES and Zhang F 2014 Development and applications of CRISPR-Cas9 for genome engineering. *Cell* **157** 1262–1278
- Hyun Y, Kim J, Cho SW, Choi Y, Kim JS and Coupland G 2015 Site-directed mutagenesis in *Arabidopsis thaliana* using dividing tissue-targeted RGEN of the CRISPR/Cas system to generate heritable null alleles. *Planta* **241** 271–284
- Jameel A, Noman M, Liu W, Ahmad N, Wang F, Li X and Li H 2020 Tinkering cis motifs jigsaw puzzle led to root-specific drought-inducible novel synthetic promoters. *Int. J. Mol. Sci.* **21** 1357
- Jiang P, Zhang K, Ding Z, He Q, Li W, Zhu S and Li K 2018 Characterization of a strong and constitutive promoter from the *Arabidopsis* serine carboxypeptidase-like gene AtSCPL30 as a potential tool for crop transgenic breeding. *BMC Biotech.* **18** 59
- Johnson JA, Bragg JN, Lawrence DM and Jackson AO 2003 Sequence elements controlling expression of Barley stripe mosaic virus subgenomic RNAs in vivo. *Virology* **313** 66–80
- Kakrana A, Kumar A, Satheesh V, Abdin MZ, Subramaniam K, Bhattacharya RC and Jain PK 2017 Identification validation and utilization of novel nematode-responsive root-specific promoters in *Arabidopsis* for inducing host-delivered RNAi mediated root-knot nematode resistance. *Front Plant Sci.* **8** 2049
- Kanofsky K, Lehmeier M, Schulze J and Hehl R 2016 Analysis of microbe-associated molecular pattern-responsive synthetic promoters with the parsley protoplast system. *Methods Mol. Biol.* **1482** 163–174
- Khanthapok P, Sang-Awut N, Chakhonkaen S, Pitngam K, Osadencio A, Sukrong S and Muangprom A 2018 Identification of ethanol-inducible genes and isolation of the myb-related protein-like promoter in *Oryza sativa* L. *J. Plant Growth Regul.* **37** 452–470
- Keadtidumrongkul P, Suttangkakul A, Pinmanee P, Pattana K, Kittiwongwattana C, Apisitwanich S and Vuttipongchaikij S 2017 Growth modulation effects of CBM2a under the control of AtEXP4 and CaMV35S promoters in *Arabidopsis thaliana*, *Nicotiana tabacum* and *Eucalyptus camaldulensis*. *Transgenic Res.* **26** 447–463
- Kelly G, Lugassi N, Belausov E, Wolf D, Khamaisi B, Brandsma D, Kottapalli J, Fidel L, Ben-Zvi B, Egbaria A, Acheampong AK, Zheng C, Or E, Distelfeld A, David-Schwartz R, Carmi N and Granot D 2017 The *Solanum tuberosum* KST1 partial promoter as a tool for guard cell expression in multiple plant species. *J. Exp. Bot.* **68** 2885–2897
- Khan ZA, Abdin MZ and Khan JA 2015 Functional characterization of a strong bi-directional constitutive plant promoter isolated from cotton leaf curl Burewala virus. *PLoS One* **10** e0121656
- Kevin E, Mc Bride and Kristin R 1989 Improved binary vectors for *Agrobacterium*-mediated plant transformation. Summerfelt Calgene Inc. 1920 Fifth St., Davis, CA 95616, USA

- Kluth A, Sprunck S, Becker D, Lorz H and Lutticke S 2002 5' deletion of a *gbss1* promoter region from wheat leads to changes in tissue and developmental specificities. *Plant Mol. Biol.* **49** 669–682
- Koia J, Moyle R, Hendry C, Lim L and Botella JR 2013 Pineapple translation factor SUI1 and ribosomal protein L36 promoters drive constitutive transgene expression patterns in *Arabidopsis thaliana*. *Plant Mol. Biol.* **81** 327–336
- Koschmann J, Machens F, Becker M, Niemeyer J, Schulze J, Bulow L, Stahl DJ and Hehl R 2012 Integration of bioinformatics and synthetic promoters leads to the discovery of novel elicitor-responsive cis-regulatory sequences in *Arabidopsis*. *Plant Physiol.* **160** 178–191
- Kyozuka J, Fujimoto H, Izawa T and Shimamoto K 1991 Anaerobic induction and tissue-specific expression of maize *Adh1* promoter in transgenic rice plants and their progeny. *Mol. Gen. Genet.* **228** 40–48
- Lamacchia C, Shewry PR, Di Fonzo N, Forsyth JL, Harris N, Lazzeri PA, Napier JA, Halford NG and Barcelo P 2001 Endosperm-specific activity of a storage protein gene promoter in transgenic wheat seed. *J. Exp. Bot.* **52** 243–250
- Lessard PA, Kulaveerasingam H, York GM, Strong A and Sinskey AJ 2002 Manipulating gene expression for the metabolic engineering of plants. *Metab. Eng.* **4** 467–479
- Lewinsohn E, Schalechet F, Wilkinson J, Matsui K, Tadmor Y, Nam KH, Amar O, Lastochkin E, Larkov O, Ravid U, Hiatt W, Gepstein S and Pichersky E 2001 Enhanced levels of the aroma and flavor compound S-linalool by metabolic engineering of the terpenoid pathway in tomato fruits. *Plant Physiol.* **127** 1256–1265
- Li F, Han Y, Feng Y, Xing S, Zhao M, Chen Y and Wang W 2013 Expression of wheat expansin driven by the RD29 promoter in tobacco confers water-stress tolerance without impacting growth and development. *J. Biotechnol.* **163** 281–291
- Li JF, Norville JE, Aach J, McCormack M, Zhang D, Bush J, Church GM and Sheen J 2013 Multiplex and homologous recombination-mediated genome editing in *Arabidopsis* and *Nicotiana benthamiana* using guide RNA and Cas9. *Nat. Biotechnol.* **31** 688–691
- Li M, Wang X, Cao Y, Liu X, Lin Y, Ou Y, Zhang H and Liu J 2013a Strength comparison between cold-inducible promoters of *Arabidopsis cor15a* and *cor15b* genes in potato and tobacco. *Plant Physiol. Biochem.* **71** 77–86
- Li M, Xie C, Song B, Ou Y, Lin Y, Liu X, Zhang H and Liu J 2015 Construction of efficient tuber-specific and cold-inducible promoters in potato. *Plant Sci.* **23** 514–524
- Li T, Huang S, Zhou J and Yang B 2013 Designer TAL effectors induce disease susceptibility and resistance to *Xanthomonas oryzae* in rice. *Mol. Plant* **6** 781–789
- Li Y, Liu S, Yu Z, Liu Y and Wu P 2013b Isolation and characterization of two novel root-specific promoters in rice (*Oryza sativa* L.). *Plant Sci.* **207** 37–44
- Li J, Xu RF, Qin RY, Ma H, Li H, Zhang YP, Li L, Wei PC and Yang JB 2014 Isolation and functional characterization of a novel rice constitutive promoter. *Plant Cell Rep.* **33** 1651–60
- Li Z, Jayasankar S and Gray DJ 2001 Expression of a bifunctional green fluorescent protein (GFP) fusion marker under the control of three constitutive promoters and enhanced derivatives in transgenic grape (*Vitis vinifera*). *Plant Sci.* **160** 877–887
- Liu C, Fukumoto T, Matsumoto T, Gena P, Frascaria D, Kaneko T et al 2013 Aquaporin OsPIP1; 1 promotes rice salt resistance and seed germination. *Plant Physiol. Biochem.* **63** 151–158
- Liu W, Mazarei M, Peng Y, Fethe MH, Rudis MR, Lin J, Millwood RJ, Arelli PR and Stewart CN 2014 Computational discovery of soybean promoter cis-regulatory elements for the construction of soybean cyst nematode-inducible synthetic promoters. *Plant Biotechnol. J.* **12** 1015–1026
- Liu YG and Whittier RF 1995 Thermal asymmetric interlaced PCR: automatable amplification and sequencing of insert end fragments from P1 and YAC clones for chromosome walking. *Genomics* **25** 674–681
- Liu X, Yang W, Li Y, Li S, Zhou X, Zhao Q, Fan Y, Lin M and Chen R 2016 The intergenic region of the maize defensin-like protein genes *Def1* and *Def2* functions as an embryo-specific asymmetric bidirectional promoter. *J. Exp. Bot.* **67** 4403–4413
- Liu Z, Luo QH, Wang JM, Li XF and Yang Y 2015 Functional characterization and analysis of the *Arabidopsis* UGT71C5 promoter region. *Genet. Mol. Res.* **14** 191 73–83
- Long L, Guo DD, Gao W, Yang WW, Hou LP, Ma XN and Song CP 2018 Optimization of CRISPR/Cas9 genome editing in cotton by improved sgRNA expression. *Plant Methods* **14** 85
- Lota F, Wegmüller S, Buer B, Sato S, Bräutigam A, Hanf B and Bucher M 2013 The cis-acting CTTC-P1BS module is indicative for gene function of *LjVTI12* a Qb-SNARE protein gene that is required for arbuscule formation in *Lotus japonicas*. *Plant J.* **74** 280–293
- Ma X, Zhang Q, Zhu Q, Liu W, Chen Y, Qiu R et al. 2015 A robust CRISPR/Cas9 system for convenient high-efficiency multiplex genome editing in monocot and dicot plants. *Mol. Plant* **8** 1274–1284
- Ma F, Qi H, Hu Y, Jiang Q, Zhang LG, Xue P and Zhang Q 2018 The mitochondrial endonuclease M20 Participates in the down-regulation of mitochondrial DNA in pollen cells. *Plant Physiol.* **178** 1537–1550
- Maiti IB and Shepherd RJ 1998 Isolation and expression analysis of peanut chlorotic streak caulimovirus (PCISV) full-length transcript (FLt) promoter in transgenic plants. *Biochem. Biophys. Res. Commun.* **244** 440–444
- Malik K, Wu K, Li XQ, Martin-Heller T, Hu M, Foster E, Tian L, Wang C, Ward K, Jordan M, Brown D, Gleddie S, Simmonds D, Zheng S, Simmonds J and Miki B 2002 A

- constitutive gene expression system derived from the tCUP cryptic promoter elements. *Theor. Appl. Genet.* **105** 505–514
- Mann DG, King ZR, Liu W, Joyce BL, Percifield RJ, Hawkins JS, LaFayette PR, Artelt BJ, Burris JN, Mazarei M, Bennetzen JL, Parrott WA and Stewart CN 2011 Switchgrass (*Panicum virgatum* L) polyubiquitin gene (PvUbi1 and PvUbi2) promoters for use in plant transformation. *BMC Biotechnol.* **11** 74
- Mao Y, Zhang Z, Feng Z, Wei P, Zhang H, Botella JR and Zhu JK 2016 Development of germ-line-specific CRISPR-Cas9 systems to improve the production of heritable gene modifications in *Arabidopsis*. *Plant Biotechnol. J.* **14** 519–532
- Marshallsay C, Connelly S and Filipowicz W 1992 Characterization of the U3 and U6 snRNA genes from wheat: U3 snRNA genes in monocot plants are transcribed by RNA polymerase III. *Plant Mol. Biol.* **19** 973–983
- Marzabal P, Busk PK, Ludevid MD and Torrent M 1998 The bifactorial endosperm box of gamma-zein gene: characterisation and function of the Pb3 and GZM cis-acting elements. *Plant J.* **16** 41–52
- McElroy D, Blowers AD, Jenes B and Wu R 1991 Construction of expression vectors based on the rice actin 1 (*Act1*) 5' region for use in monocot transformation. *Mol. Gen. Genet.* **231** 150–160
- Mehrotra R and Mehrotra S 2010 Promoter activation by ACGT in response to salicylic and abscisic acids is differentially regulated by the spacing between two copies of the motif. *J. Plant Physiol.* **167** 1214–1218
- Meyer S, Mumm P, Imes D, Endler A, Weder B, Al-Rasheid KA and Hedrich R 2010 AtALMT12 represents an R-type anion channel required for stomatal movement in *Arabidopsis* guard cells. *Plant J.* **63** 1054–1062
- Miki D, Zhang W, Zeng W, Feng Z and Zhu JK 2018 CRISPR/Cas9-mediated gene targeting in *Arabidopsis* using sequential transformation. *Nat. Commun.* **9** 1967
- Mikami M, Toki S, and Endo M 2017 In Planta processing of the SpCas9-gRNA complex. *Plant Cell Physiol* **58** 1857–1867
- Miyata LY, Harakava R, Stipp LC, Mendes BM, Appezzato-da-Glória B, de Assis Alves Mourão and Filho F 2012 GUS expression in sweet oranges (*Citrus sinensis* L Osbeck) driven by three different phloem-specific promoters. *Plant Cell Rep.* **31** 2005–2013
- Mittler R and Blumwald E 2010 Genetic engineering for modern agriculture: challenges and perspectives. *Annu. Rev. Plant. Biol.* **61** 443–462
- Nakashima K, Tran LS, Van Nguyen D, Fujita M, Maruyama K, Todaka D, Ito Y, Hayashi N, Shinozaki K and Yamaguchi-Shinozaki K 2007 Functional analysis of a NAC-type transcription factor OsNAC6 involved in abiotic and biotic stress-responsive gene expression in rice. *Plant J.* **51** 617–630
- Naoumkina M and Dixon RA 2011 Characterization of the mannan synthase promoter from guar (*Cyamopsis tetragonoloba*). *Plant Cell Rep.* **30** 997–1006
- Nekrasov V, Staskawicz B, Weigel D, Jones JD and Kamoun S 2013 Targeted mutagenesis in the model plant *Nicotiana benthamiana* using Cas9 RNA-guided endonuclease. *Nat. Biotech.* **31** 691
- Nicholass FJ, Smith CJ, Schuch W, Bird CR and Grierson D 1995 High levels of ripening-specific reporter gene expression directed by tomato fruit polygalacturonase gene-flanking regions. *Plant Mol. Biol.* **28** 423–435
- Nunberg AN, Li Z, Bogue MA, Vivekananda J, Reddy AS and Thomas TL 1994 Developmental and hormonal regulation of sunflower helianthinin genes: proximal promoter sequences confer regionalized seed expression. *Plant Cell* **6** 473–486
- Obertello M, Santi C, Sy MO, Laplaze L, Auguy F, Bogusz D and Franche C 2005 Comparison of four constitutive promoters for the expression of transgenes in the tropical nitrogen-fixing tree *Allocasuarina verticillata*. *Plant Cell Rep.* **24** 540–548
- Odell JT, Nagy F and Chua NH 1985 Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter. *Nature* **313** 810–812
- Oh SJ, Song SI, Kim YS, Jang HJ, Kim SY, Kim M and Kim JK 2005 *Arabidopsis* CBF3/DREB1A and ABF3 in transgenic rice increased tolerance to abiotic stress without stunting growth. *Plant Physiol.* **138** 341–351
- Ordon J, Bressan M, Kretschmer C, Dall'Osto L, Marillonnet S, Bassi R and Stuttmann J 2020 Optimized Cas9 expression systems for highly efficient *Arabidopsis* genome editing facilitate isolation of complex alleles in a single generation. *Funct. Integr. Genomics* **20** 151–162
- Osakabe Y, Osakabe K and Chiang VL 2009 Characterization of the tissue-specific expression of phenylalanine ammonia-lyase gene promoter from loblolly pine (*Pinus taeda*) in *Nicotiana tabacum*. *Plant Cell Rep.* **28** 1309–1317
- Park SH, Yi N, Kim YS, Jeong MH, Bang SW, Choi YD and Kim JK 2010 Analysis of five novel putative constitutive gene promoters in transgenic rice plants. *J. Exp. Bot.* **61** 2459–2467
- Patro S, Maiti IB and Dey N 2013 Development of an efficient bi-directional promoter with tripartite enhancer employing three viral promoters. *J. Biotechnol.* **163** 311–317
- Porto MS, Pinheiro MP, Batista VG, dos Santos RC, Filho Pde A and de Lima LM 2014 Plant promoters: an approach of structure and function. *Mol. Biotechnol.* **56** 38–49
- Potenza C, Aleman L and Gopalan CS 2004 Targeting transgene expression in research agricultural and environmental applications: promoters used in plant transformation. *In Vitro Cell Dev. Biol. Plant* **40** 1–22
- Que Q, Chilton M-DM, de Fontes CM, He C, Nuccio M, Zhu T, Wu Y, Chen JS, Shi L 2010 Trait stacking in

- transgenic crops: challenges and opportunities. *GM Crops* **1** 220–229
- Qiu CH, Li H, Li J, Qin RY, Xu RF, Yang YC, Ma H, Song FS, Li L, Wei PC and Yang JB 2015 Isolation and characterization of three cadmium-inducible promoters from *Oryza sativa*. *J. Biotechnol.* **216** 1–19
- Rai M, He C and Wu R 2009 Comparative functional analysis of three abiotic stress-inducible promoters in transgenic rice. *Transgenic Res.* **187** 87–99
- Rao TSRB, Naresh JV, Reddy PS, Reddy MK and Mallikarjuna G 2017 Expression of *Pennisetum glaucum* eukaryotic translational initiation factor 4A (PgelF4A) confers improved drought salinity and oxidative stress tolerance in groundnut. *Front. Plant Sci.* **8** 453
- Reddy CS, Vijayalakshmi M, Kaul T, Islam T and Reddy MK 2015 Improving flavour and quality of tomatoes by expression of synthetic gene encoding sweet protein monellin. *Mol. Biotechnol.* **57** 448–53
- Reddy PS, Mahanty S, Kaul T, Nair S, Sopory SK and Reddy MK 2008 A high-throughput genome-walking method and its use for cloning unknown flanking sequences. *Anal. Biochem.* **381** 248–253
- Reksiri W, Zhang X, Xiong H and Chen X 2013 Expression and promoter analysis of six heat stress-inducible genes in rice. *Sci. World J.* 397401
- Roccaro M, Ahmadinejad N, Colby T and Somssich IE 2013 Identification of functional cis-regulatory elements by sequential enrichment from a randomized synthetic DNA library. *BMC Plant Biol.* **13** 164
- Rusconi F, Simeoni F, Francia P, Cominelli E, Conti L, Riboni M and Galbiati M 2013 The Arabidopsis thaliana MYB60 promoter provides a tool for the spatio-temporal control of gene expression in stomatal guard cells. *J. Exp. Bot.* **64** 3361–3371
- Saed TR, Ismail I, Zainal Z and Abdullah SN 2012 The stearyl-acyl-carrier-protein desaturase promoter (Des) from oil palm confers fruit-specific GUS expression in transgenic tomato. *J. Plant Physiol.* **1691** 290–300
- Sahoo DK, Sarkar S, Raha S, Maiti IB and Dey N 2014 Comparative analysis of synthetic DNA promoters for high-level gene expression in plants. *Planta* **240** 855–875
- Sakaue K, Harada H and Matsuda Y 2008 Development of gene expression system in a marine diatom using viral promoters of a wide variety of origin. *Physiol. Plant.* **133** 59–67
- Sawant SV, Kiran K, Mehrotra R, Chaturvedi CP, Ansari SA, Singh P, Lodhi N and Tuli R 2005 A variety of synergistic and antagonistic interactions mediated by cis-acting DNA motifs regulate gene expression in plant cells and modulate stability of the transcription complex formed on a basal promoter. *J. Exp. Bot.* **56** 2345–2353
- Schimpl S, Fauser F and Puchta H 2014 The CRISPR/C as system can be used as nuclease for in planta gene targeting and as paired nickases for directed mutagenesis in Arabidopsis resulting in heritable progeny. *Plant J.* **80** 1139–1150
- Schmulling T, Schell J and Spena A 1989 Promoters of the rolA B and C genes of *Agrobacterium rhizogenes* are differentially regulated in transgenic plants. *Plant Cell* **1** 665–670
- Seemanpillai M, Dry I, Randles J and Rezaian A 2003 Transcriptional silencing of geminiviral promoter-driven transgenes following homologous virus infection. *Mol. Plant Microbe Interact.* **16** 429–438
- Shah SH, Jan SA, Ahmad N, Khan SU, Kumar T, Iqbal A and Nasir F 2015 Use of different promoters in transgenic plant development: current challenges and future perspectives. *Am Eurasian J. Agric. Environ. Sci.* **15** 664–675
- Shahmuradov IA, Umarov RK and Solovyev VV 2017 TSSPlant: a new tool for prediction of plant Pol II promoters. *Nucleic Acids Res.* **45** e65
- Shan Q, Wang Y, Chen K, Liang Z, Li J, Zhang Y et al 2013 Rapid and efficient gene modification in rice and Brachypodium using TALENs. *Mol. Plant* **6** 1365–1368
- Sharma KK, Lavanya M and Anjaiah V 2006 Agrobacterium-mediated production of transgenic pigeonpea (*Cajanus cajan* L. Millsp.) expressing the synthetic Bt cry1Ab gene. *In Vitro Cell. Dev. Biol.-Plant* **42** 165–173
- Shen YG, Zhang WK, Yan DQ, Du BX, Zhang JS, Liu Q and Chen SY 2003 Characterization of a DRE-binding transcription factor from a halophyte *Atriplex hortensis*. *Theor. Appl. Genet.* **107** 155–161
- Singer SD, Hily JM and Cox KD 2011 The sucrose synthase-1 promoter from Citrus sinensis directs expression of the β -glucuronidase reporter gene in phloem tissue and in response to wounding in transgenic plants. *Planta* **234** 623–637
- Smale ST and Kadonaga JT 2003 The RNA polymerase II core promoter *Annu. Rev. Biochem.* **72** 449–479
- Singh KB, Foley RC and Onate-Sanchez L 2002 Transcription factors in plant defense and stress responses. *Curr. Opin. Plant Biol.* **5** 430–436
- Srinath T, Reddy VD and Rao KV 2017 Isolation and functional characterization of a novel stress inducible promoter from pigeonpea (*Cajanus cajan* L.). *Plant Cell Tiss. Org. Cult.* **128** 457–468
- Srivastava VK, Raikwar S and Tuteja N 2014 Cloning and functional characterization of the promoter of *PsSEOF1* gene from *Pisum sativum* under different stress conditions using *Agrobacterium*-mediated transient assay. *Plant Signal Behav.* **9** e29626
- Sunkara S, Bhatnagar-Mathur P and Sharma KK 2014 Isolation and functional characterization of a novel seed-specific promoter region from peanut. *Appl Biochem Biotechnol.* **172** 325–339
- Tao YB, He LL, Niu LJ and Xu ZF 2015 Isolation and characterization of an ubiquitin extension protein gene (JcUEP) promoter from *Jatropha curcas*. *Planta* **241** 823–836

- Tashkandi M, Ali Z, Aljedaani F, Shami A and Mahfouz MM 2018 Engineering resistance against Tomato yellow leaf curl virus via the CRISPR/Cas9 system in tomato. - *Plant Signal. Behav.* **13** e1525996
- Tiwari V, Patel MK, Chaturvedi AK, Mishra A and Jha B 2016 Functional characterization of the tau class glutathione-S-transferases gene (SbGSTU) promoter of *Salicornia brachiata* under salinity and osmotic stress. *PLoS One* **11** e0148494
- Tompa M, Li N, Bailey TL, Church GM, et al. 2005 Assessing computational tools for the discovery of transcription factor binding sites *Nat. Biotechnol.* **23** 137–144
- Twyman RM 2003 Growth and development: control of gene expression regulation of transcription; In *Encyclopedia of applied plant sciences* (Eds) Thomas B, Murphy DJ and Murray GB (Elsevier Science, London UK) pp 558–567
- van der Geest AH and Hall TC 1997 The β -phaseolin5 matrix attachment region acts as an enhancer facilitator. *Plant Mol. Bio.* **33** 553–557
- Van der Does D, Leon-Reyes A, Koornneef A, Van Verk MC, Rodenburg N, Pauwels L, Goossens A, Körbes AP, Memelink J, Ritsema T, Van Wees SC and Pieterse CM 2013 Salicylic acid suppresses jasmonic acid signaling downstream of SCFCO11-JAZ by targeting GCC promoter motifs via transcription factor ORA59. *Plant Cell* **25** 744–761
- Viana AA, Fragoso RR, Guimarães LM, Pontes N, Oliveira-Neto OB, Artico S et al 2011 Isolation and functional characterization of a cotton ubiquitination-related promoter and 5'UTR that drives high levels of expression in root and flower tissues. *BMC Biotechnol.* **11** 115
- Wang JY, Wang JP and Yang HF 2016a Identification and functional characterization of the NAC gene promoter from *Populus euphratica*. *Planta* **244** 417–427
- Wang P, Du Y, Zhao X, Miao Y and Song CP 2013 The MPK6-ERF6-ROS-responsive cis-acting Element7/GCC box complex modulates oxidative gene transcription and the oxidative response in *Arabidopsis*. *Plant Physiol.* **161** 1392–1408
- Wang R, Yan Y, Zhu M, Yang M, Zhou F, Chen H and Lin Y 2016b Isolation and Functional Characterization of bidirectional promoters in rice. *Front. Plant Sci.* **77** 66
- Wang R, Zhu M, Ye R, Liu Z, Zhou F, Chen H and Lin Y 2015 Novel green tissue-specific synthetic promoters and cis-regulatory elements in rice. *Sci. Rep.* **51** 825
- Wang J and Oard JH 2003 Rice ubiquitin promoters: deletion analysis and potential usefulness in plant transformation systems: Genetic transformation and hybridization. *Plant Cell Rep.* **221** 29–34
- Wang Y, Lin S, Song Q, Li K, Tao H, Huang J et al 2014 Genome-wide identification of heat shock proteins (Hsps) and Hsp interactors in rice: Hsp70s as a case study. *BMC Genom.* **15** 344
- Wang X, Wang L, Wang Y, Liu H, Hu D. Zhang N et al 2018 Arabidopsis PCaP2 plays an important role in chilling tolerance and ABA response by activating CBF- and SnRK2-mediated transcriptional regulatory network. *Frontiers in plant science* **9** 215
- Wu CY, Suzuki A, Washida H and Takaiwa F 1998 The GCN4 motif in a rice glutelin gene is essential for endosperm-specific gene expression and is activated by Opaque-2 in transgenic rice plants. *Plant J.* **14** 673–683
- Wu L, E-Mezawy A and Shah S 2011 A seed coat outer integument-specific promoter for *Brassica napus*. *Plant Cell Rep.* **30** 75–80
- Xiao K, Zhang C, Harrison M and Zeng YW 2005 Isolation and characterization of novel plant promoter that directs strong constitutive expression of transgene in plants. *Mol. Breed.* **15** 221–231
- Xu D, Duan X, Wang B, Hong B, Ho T and Wu R 1996 Expression of a late embryogenesis abundant protein gene HVA1 from barley confers tolerance to water deficit and salt stress in transgenic rice. *Plant Physiol.* **110** 249–257
- Xu F, Huang XH, Li LL, Deng G, Cheng H, Rong XF, Li JB and Cheng SY 2013 Molecular cloning and characterization of GbDXS and GbGGPPS gene promoters from *Ginkgo biloba*. *Genet. Mol. Res.* **4** 293–301
- Yamaguchi-Shinozaki K and Shinozaki K 2006 Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annu. Rev. Plant Biol.* **57** 781–803
- Yan L, Wei S, Wu Y, Hu R, Li H, Yang W and Xie Q 2015 High-efficiency genome editing in *Arabidopsis* using YAO promoter-driven CRISPR/Cas9 system. *Mol. Plant* **8** 1820–1823
- Yang IC, Iommarini JP, Becker DK, Hafner GJ, Dale JL and Harding RM 2003 A promoter derived from taro bacilliform badnavirus drives strong expression in transgenic banana and tobacco plants. *Plant Cell Rep.* **21** 1199–1206
- Yang J, Wang X, Hasi A and Wang Z 2018 Structural and functional analysis of a bidirectional promoter from *Gossypium hirsutum* in *Arabidopsis*. *Int. J. Mol. Sci.* **19** 3291
- Ye X, Al-Babili S, Klott A, Zhang J, Lucca P, Beyer P and Potrykus I 2000 Engineering the provitamin A (beta-carotene) biosynthetic pathway into (carotenoid-free) rice endosperm. *Science* **287** 303–305
- Ye R, Zhou F and Lin Y 2012 Two novel positive cis-regulatory elements involved in green tissue-specific promoter activity in rice (*Oryza sativa* L ssp). *Plant Cell Rep.* **31** 1159–72
- Zavallo D, Lopez Bilbao M, Hopp HE and Heinz R 2010 Isolation and functional characterization of two novel seed-specific promoters from sunflower (*Helianthus annuus* L). *Plant Cell Rep.* **29** 239–248
- Zhang C, Pan S, Chen H, Cai T, Zhuang C, Deng Y, Zhuang Y, Zeng Y, Chen S and Zhuang W 2016 Characterization of NtREL1 a novel root-specific gene from tobacco and

- upstream promoter activity analysis in homologous and heterologous hosts. *Plant Cell Rep.* **35** 757–769
- Zhang H, Hou J, Jiang P, Qi S, Xu C, He Q, Ding Z, Wang Z, Zhang K and Li K 2016 Identification of a 467 bp promoter of maize phosphatidylinositol synthase gene (ZmPIS) which confers high-level gene expression and salinity or osmotic stress inducibility in transgenic tobacco. *Front. Plant Sci.* **7** 42
- Zhou J, Yang Y, Wang X, Yu F, Yu C, Chen J, Cheng Y, Yan C and Chen J 2013 Enhanced transgene expression in rice following selection controlled by weak promoters. *BMC Biotechnol.* **27** 13–29