

Organic synthesis *via* photoinduced hydrogen and electron transfer reactions

S V KESSAR

Department of Chemistry, Panjab University, Chandigarh 160014, India

Abstract. Synthetic utility of photoinduced intramolecular hydrogen transfer reactions of ketones, amides, imines, nitrites and nitro compounds is highlighted by illustrative examples. Role of light-initiated electron transfer from amines, mercaptans, aryl groups and from lithiated ketones, amides and imines in reactions of synthetic interest is discussed.

Keywords. Ketones; amides; imines; nitrites; nitro compounds; photocyclisation; benzophenanthridines; berberines; aporphines.

1. Introduction


Adsorption of light by an organic molecule promotes it to a relatively high energy state and therein lies the strength as well as the weakness of photochemistry as a synthetic tool. Whereas a desired reaction pathway which is thermally inaccessible can become available in the excited state many side reactions may also occur from it. Nevertheless, numerous photoreactions showing good stereo-, regio, and chemoselectivity have been discovered. The present article is intended to highlight the utility of such reactions for synthesis of complex organic molecules. It is limited only to transformations which involve intramolecular hydrogen or electron transfer as the primary photoprocess. Thus many photoreactions of proven synthetic utility like cycloadditions, electrocyclic ring closures and rearrangements (Wender and Howbert 1982; Oppolzer and Godel 1978; Schulty *et al* 1977; Ninomiya and Naito 1981) are not covered. Further, the examples are chosen primarily from author's own area of interest *viz* the photochemistry of nitrogen containing functional groups.

2. Hydrogen transfer reactions

On irradiation, functional groups containing oxygen, sulphur and nitrogen multiple bonds can abstract hydrogen from a suitable donor inter or intramolecularly. The latter process is often followed by fragmentation or ring closures of synthetic importance as illustrated below for some classes of compounds.

2.1 Ketones

Photoreactions based on intramolecular hydrogen abstraction are efficient and synthetically useful if a six-membered transition state is involved (Quinkert *et al* 1980). However factors like framework rigidity (Paquette *et al* 1983), solvent-induced folding (Breslow *et al* 1973) and complexation (Gotschi and Eschenmoser 1973) can force



distant hydrogen atoms into a disposition favourable for abstraction resulting in 'remote functionalization'.

2.2 Amides







As compared to ketones carboxylic acid derivatives show less proclivity for hydrogen abstraction (Coyle 1978) but rich photochemistry is associated with imides (Kanaoka 1978). In α -ketoamides also hydrogen abstraction is facile and may be followed by a ring closure to a β -lactam (Henry-Logan and Chen 1973; Johansson *et al* 1976) or a α -lactone (Aoyama *et al* 1983).

The diradical initially formed from a α -ketoamide can also undergo a Norrish type-II fragmentation (Aoyama *et al* 1981) but in the case of tetrahydroisoquinoline derivatives further oxidation of the nitrogenous segment is observed (Kessar *et al* 1978). This aromatisation is of mechanistic and synthetic interest due to complications observed during dehydrogenation (Sainsbury *et al* 1969). The vinalogous benzamides with an ortho aldehydic group furnish, on irradiation, amino-lactones, presumably through a 1,5 hydrogen transfer (unpublished results from this laboratory).

2.3 Amines and imines

Many a claimed photoreductions of Schiff bases have been shown to be caused through 'chemical sensitisation' by carbonyl impurities (Padwa 1977; Pratt 1977).

The lack of photoinitiated hydrogen abstraction in imines has been variously attributed to energy dissipation through rapid *cis-trans* isomerisation, $\pi\pi^*$ character of the excited state or the lower strength of the incipient NH bond (Padwa 1977; Pratt 1977; Ohta and Tokumaru 1974). Whatever the reasons for the absence of this photoreactivity in imines, the methodology dependent on it is not available for heterocyclic synthesis. However, recently it has been shown that certain α -ketoimines do undergo phototransformations leading to spirobenzyl isoquinoline and berberine




alkaloids (Kessar *et al* 1978). The procedure has been extended to synthesis of indole alkaloids also (Hirai *et al* 1983).

The above reaction when used in conjunction with a known photodegradation of berberine derivatives (Wu *et al* 1978) can bring about an intriguing transposition of D ring substituents of these alkaloids (Kessar *et al* 1982). Mechanisms of both the involved photoreactions have been investigated by deuterium labelling experiments (Kessar *et al* 1982).

2.4 Nitrites and nitro compounds

Barton reaction (Barton and Akhtar 1962) is an outstanding example of the synthetic utility of photochemical reactions, although here the hydrogen abstraction step may not be occurring on the excited state surface.

Nitro compounds also serve as useful photolabile amino protecting groups (Rajashekhara Pillai 1980) as shown below.



3. Electron transfer reactions







In photoinduced electron transfer reactions, back electron transfer is an energy wasting process. Nature has its own mechanism for overcoming it and in artificial solar energy utilisation also ingenious procedures have been devised to minimise it (Whitten 1980). For a synthetically efficient reaction, electron transfer should be followed by a fast event like loss of a nucleofuge from the receptor or of a proton from the donor. A few such examples involving different functional groups are given below.

3.1 Ketones and amides

Even carbonyl triplets with a $\pi\pi^*$ character which are generally unreactive in hydrogen abstraction can be efficiently reduced by amines through electron transfer followed by proton transfer (Cohen *et al* 1973). Unfortunately, in the case of aminoketones intramolecular proton transfer is inefficient (Wagner and Ersfeld 1976) and not many reactions of synthetic import have been reported. However, electron transfer from a sulphur atom followed by intramolecular proton transfer has been invoked to explain the formation of medium and large ring systems on irradiation of certain imides (Sato *et al* 1976).

A mechanism involving electron transfer from the nitrogen atom has also been proposed for γ -hydrogen abstraction in α -ketoamides (Aoyama *et al* 1983). Evidence for bimolecular electron transfer in pyruvates has been adduced recently (Davidson *et al* 1983).

Light-promoted electron transfer from enolate anions to aryl halides has been invoked to explain the photoinduced $S_{RN}1$ reactions (Bunnet 1978). An intramolecular




version has been used in the synthesis of cephalotoxin (Semmelhack and Bargar 1980) and other medium-sized ring compounds (Semmelhack and Bargar 1977). Electron transfer-mediated cyclisation of metalated amides constitutes the key step in the synthesis of cherylline and corynoline-type alkaloids (Kessar *et al* 1981).


Witkop reaction which is very useful for synthesis of benzazepinones is claimed to entail electron transfer in the reverse direction *i.e.* from the aromatic moiety to the chlorocarbonyl function (Yonemitsu *et al* 1968; Hamada *et al* 1977).

3.2 Amines, imines and enamines

Metalated imines have been found to undergo photocyclisation (unpublished results from this laboratory) in a manner similar to that of carbonyl enolates. This reaction can open up convenient routes to many alkaloidal structures.

Although the nucleophilicity of the enamine carbon atom is well established, photocyclisation based on intramolecular electron transfer to aryl halides could not be induced in these compounds (unpublished results from this laboratory). In contrast, irradiation of the corresponding aromatic amines leads to ready ring closure (Mizuno *et al* 1973). Based on this observation, efficient syntheses of chelilutine and sanguilutine were carried out (Kessar *et al* 1977) to settle the controversy regarding the placement of alkoxy groups in this set of alkaloids (Ishii *et al* 1975). This route has been extended to





the synthesis of macarpine which carries six alkoxy groups on the benzophenanthridine nucleus (Takao *et al* 1981).

Finally, success of certain photoreactions may hinge on preventing the electron transfer process. For example, lack of aporphine formation on irradiation of ortho-halogenated benzylisoquinolines has been ascribed to a deleterious electron transfer from the nitrogen atom (Kessar *et al* 1981). A strategem for preventing this, and for locking the aryl moieties in close proximity, is based on introduction of a hydrogen bonding group. Synthesis of oliveridine and related alkaloids could thus be achieved readily (Kessar *et al* 1980; Kessar *et al* 1983).

Acknowledgements

Valuable participation of the author's research associates in the quoted work from this laboratory and assistance of Shri K R Agnihotri in the preparation of this manuscript are gratefully acknowledged.

References

- Aoyama H, Sakamoto M, Kuwabara K, Yoshida K and Qmote Y 1983 *J. Am. Chem. Soc.* **105** 1958
 Aoyama H, Sakamoto M and Qmote Y 1981 *J. Chem. Soc. Perkin I* 1357
 Barton D H R and Akhtar M 1962 *J. Am. Chem. Soc.* **84** 1496
 Breslow R, Baldwin S, Flechtner T, Kalicky P, Liu S and Washburn W 1973 *J. Am. Chem. Soc.* **95** 3251
 Bunnet J F 1978 *Acc. Chem. Res.* **11** 413
 Cohen S G, Davis G A and Clark W D K 1973 *Chem. Rev.* **73** 141
 Coyle J D 1978 *Chem. Rev.* **78** 115
 Davidson R S, Goodwin D and Pratt J L 1983 *J. Chem. Soc. Perkin II* 1729
 Gotschi E and Eschenmoser A 1973 *Angew. Chem. Int. Ed. Engl.* **12** 912
 Hamada T, Ohmori M and Yonemitsu O 1977 *Tetrahedron Lett.* 1519
 Henry-Logan K R and Chen C G 1973 *Tetrahedron Lett.* 1103
 Hirai Y, Egawa H and Yamazaki T 1983 Abstracts of the ninth International Congress of Heterocyclic Chemistry, p. 151
 Ishii H, Harada K, Ishida T, Veda E and Nakajima K 1975 *Tetrahedron Lett.* 319
 Johansson N G, Akermark B and Sjoberg B 1976 *Acta Chem. Scand.* **B30** 383
 Kanaoka Y 1978 *Acc. Chem. Res.* **11** 407
 Kessar S V, Gupta Y P and Mohammad T 1983 *Indian J. Chem.* **B22** 321
 Kessar S V, Gupta Y P, Singh T V, Nanda A, Sood A and Agnihotri K R 1982 *Tetrahedron Lett.* 3619
 Kessar S V, Singh P, Chawla R and Kumar P 1981a *J. Chem. Soc. Chem. Commun.* 1074

- Kessar S V, Gupta Y P and Mohammad T 1981b *Indian J. Chem.* **B20** 984
Kessar S V, Gupta Y P, Yadav V S, Narula M and Mohammad T 1980 *Tetrahedron Lett.* 3307
Kessar S V, Gupta Y P, Singh P, Gupta S K and Nanda A K 1978a *J. Chem. Soc. Chem. Commun.* 885
Kessar S V, Gupta Y P, Gainda A and Varsha A 1978b *Indian J. Chem.* **B16** 319
Kessar S V, Gupta Y P, Dhingra K, Sharma G S and Narula S 1977 *Tetrahedron Lett.* 1459
Mizuno K, Pac C and Sakurai H 1973 *Bull. Chem. Soc. Jpn* **46** 3316
Ninomiya I and Naito T 1981 *Heterocycles* **15** 1433
Ohta H and Tokumaru K 1974 *Tetrahedron Lett.* 2965
Oppolzer W and Godel T 1978 *J. Am. Chem. Soc.* **100** 2583
Padwa A 1977 *Chem. Rev.* **77** 37
Paquette L A, Ternasky R J, Balogh D W and Kentgen G 1983 *J. Am. Chem. Soc.* **105** 5446
Pratt A C 1977 *Chem. Soc. Rev.* **6** 63
Quinkert G, Weber W D, Schwartz U and Durner G 1980 *Angew. Chem. Int. Ed. Engl.* **19** 1027
Rajashekharan Pillai V N 1980 *Synthesis I*
Sainsbury M, Brown D W, Dyke S F and Hardy G 1969 *Tetrahedron* **25** 1882
Sato Y, Nakai H, Mizoguchi T, Hatanaka Y and Kanaoka Y 1976 *J. Am. Chem. Soc.* **98** 2349
Schulzy A G, Yee Y K and Berger M H 1977 *J. Am. Chem. Soc.* **99** 8065
Semmelhack M F and Bargar T 1980 *J. Am. Chem. Soc.* **102** 7765
Semmelhack M F and Bargar T 1977 *J. Org. Chem.* **42** 1481
Takao M, Kamaguchi M, Sagiura M and Ninomiya I 1981 *Heterocycles* **16** 223
Wagner P J and Ersfeld D A 1976 *J. Am. Chem. Soc.* **98** 4515
Wender P A and Howbert J J 1982 *Tetrahedron Lett.* 3983
Whitten D G 1980 *Acc. Chem. Res.* **13** 83
Wu T T, Moniot J L and Shamma M 1978 *Tetrahedron Lett.* 3419
Yonemitsu O, Tokuyama T, Chaykovsky M and Witkop B 1968 *J. Am. Chem. Soc.* **90** 776