

Extended metal–organic solids based on benzenepolycarboxylic and aminobenzoic acids

R MURUGAVEL*, G ANANTHARAMAN, D KRISHNAMURTHY,
M SATHIYENDIRAN and M G WALAWALKAR
Department of Chemistry, Indian Institute of Technology, Powai,
Mumbai 400 076, India
e-mail: rmv@ether.chem.iitb.ernet.in

Abstract. This article describes the recent results obtained in our laboratory on the interaction of polyfunctional ligands with divalent alkaline earth metal ions and a few divalent transition metal ions. Treatment of $MCl_2 \cdot nH_2O$ ($M = Mg, Ca, Sr$ or Ba) with 2-amino benzoic acid leads to the formation of complexes $[Mg(2-aba)_2]$ (**1**), $[Ca(2-aba)_2(OH_2)_3]_\infty$ (**2**), $[{Sr(2-aba)_2(OH_2)_2} \cdot H_2O]_\infty$ (**3**), $[Ba(2-aba)_2(OH_2)]_\infty$ (**4**), respectively. While the calcium ions in **2** are hepta-coordinated, the strontium and barium ions in **3** and **4** reveal a coordination number of nine apart from additional metal–metal interactions. Apart from the carboxylate functionality, the amino group also binds to the metal centres in the case of strontium and barium complexes **3** and **4**. Complexes $[{Mg(H_2O)_6}(4-aba)_2 \cdot 2H_2O]$ (**5**), $[Ca(4-aba)_2(H_2O)_2]$ (**6**) prepared from 4-aminobenzoic acid reveal more open or layered structures. Interaction of 2-mercaptobenzoic acid with $MCl_2 \cdot 6H_2O$ ($M = Mg, Ca$), however, leads to the oxidation of the thiol group resulting in the disulphide 2,2'-dithiobis(benzoic acid). New metal–organic framework based hydrogen-bonded porous solids $[{M(btec)(OH_2)_4}]_n \cdot n(C_4H_{12}N_2) \cdot 4nH_2O$ ($btec = 1,2,4,5$ -benzene tetracarboxylate) ($M = Co$ **9**; Ni **10**; Zn **11**) have been synthesized from 1,2,4,5-benzene tetracarboxylic acid in the presence of piperazine. These compounds are made up of extensively hydrogen-bonded alternating layers of anionic M-btec co-ordination polymer and piperazinium cations. Compounds **2–11** described herein form polymeric networks in the solid-state with the aid of different coordinating capabilities of the carboxylate anions hydrogen bonding interactions.

Keywords. Coordination polymer; aminobenzoic acid; benzene tetracarboxylic acid; alkaline earth metals; supramolecular chemistry.

1. Introduction

Microporous inorganic materials like zeolites find widespread use in heterogeneous catalysis, adsorption and ion exchange processes. The rigidity and stability of such framework allow for shape and size selective inclusion of guest molecules. The design of supramolecular structures with interesting structural and magnetic properties has fuelled research in this relatively new area. Due to their interesting structural features as well as the anticipated properties of the extended solids, advances in these fields are growing rapidly.

The designed construction of extended, porous, co-ordination polymer frameworks from soluble molecular building blocks represents one of the most challenging issues in

*For correspondence


synthetic chemistry^{1,2}. Given the diversity of the ligands and metal-ligand co-ordination chemistry, there exists extensive opportunities for synthesizing co-ordination polymers with novel, porous architectures. The first challenge is to control the orientation and stereochemistry of the building blocks, in the solid state in order to achieve a target molecular topology and architecture. Therefore, the synthesis of co-ordination polymers which have structures analogous to zeolites, may yield a new generation of multidimensional network which contain channels or cavities of various sizes and shapes, unobserved as yet, in zeolites³. Polymers produced this way have been termed as *modular solids* since they are assembled from discrete molecules².

In the construction of such hybrid inorganic-organic polymer network, the choice of the ligand which forms a basic building block, plays a very important role in determining the pore size². This is especially true as demonstrated by recent reports on new, microporous, metal-organic frameworks that have been assembled with multifunctional organic molecules and several transition metal ions using a modular approach⁴. All these complexes exhibit supramolecular behaviour due to the heavy association through covalent and hydrogen bonds. It has now been well documented that by a judicious choice of building units, it is possible to produce extended cationic, anionic and neutral porous networks with unusual pore architecture and functions.


The ligands as well as the metal ions may be tailored to effect highly selective inclusion of incoming guests. Aromatic acids are expected to form rigid stable co-ordination polymers on account of their appropriately placed functional groups which are attached to a rigid template such as a benzene ring (scheme 1). Several research groups in the last five years have synthesized polymeric networks with zeolite-like morphologies using these acids as well as other types of ligands. A more detailed description of these newly unraveled structures can be found in recent reviews in this area¹. The objective of this mini-review is to mainly highlight our own research efforts in this area.

2. Results and discussion

Our interests in this area are in the use of multifunctional ligands whose donor atoms are located on a rigid phenyl group. For this purpose, we are particularly interested in



Scheme 1.



Scheme 2.


employing aromatic amino acids such as **a–c** shown in scheme 2 for two reasons. They contain carboxylate and amino functionalities, which are known to behave differently towards metal centres because of differences in their hard-soft donor characteristics. Secondly, by choosing the second substituent's position (namely, ortho, meta, or para; i.e. 60, 120 or 180° between the functional groups) one can easily vary the resultant supramolecular organization. For similar reasons, the well-known mercaptobenzoic acid (**d**) was also chosen as the building block in our initial studies.

The considerable success that has been achieved in recent years in using benzene dicarboxylic/tricarboxylic acids as building blocks has prompted us to study the suitability of 1,2,4,5-benzenetetracarboxylic acid (**e**), in conjunction with several amines for this purpose. Moreover, the relative disposition of the four carboxylate ligands on the phenyl group is likely to offer several intriguing structures which are essentially based on a one-dimensional co-ordination polymer.

2.1 Coordination polymers derived from 2-aminobenzoic acid

The metal complexes **1–4** are best prepared by mixing group 2 metal chloride hydrates with 2-amino benzoic acid (2-abaH) under basic conditions ($pH \sim 10–12$) from an aqueous solution (scheme 3)⁵. In the case of magnesium, an insoluble precipitate is obtained instantaneously, whereas compounds **2–4** are fairly soluble in water and form X-ray quality single crystals after several days from the reaction mixtures. The 2-aba complexes **1–4** have been extensively characterized by both analytical and spectroscopic methods. Additionally, the solid state structures of **2–4** have been unambiguously established by single crystal X-ray diffraction studies.

The IR spectra of these complexes reveal that all the complexes, except for magnesium complex **1**, contain at least one coordinated or un-coordinated water molecule. This was further supported by subsequent thermal studies of these complexes and single X-ray crystal structure. Further, the IR spectral studies clearly establish the non-participation of the amino group in coordination to the metal in **2**, whereas the amino group co-ordinates to strontium and barium ions in complexes **3** and **4**, respectively. This observation could be attributed to the increase in the soft character of the metal ions on going down the group.



Scheme 3.

However, it is surprising to note that a simple 1:2 M:L complex, which has no coordinated water molecules is formed in the case of magnesium complex **1**. Synthesis of the compound using other bases such as NaOH or KOH, in place of aqueous ammonia, did not change the composition of the final product obtained. In all cases the resultant product yielded the same analytical data. This observation clearly contrasts several known magnesium complexes wherein Mg^{2+} ion preferentially binds to water molecules, often in the form of $[Mg(OH_2)_6]^{2+}$.

Crystal structure of 2: The calcium atoms in $[Ca(2-aba)_2(OH_2)_3]_\infty$ are hepta-coordinated. Four of the coordination sites in **2** are occupied by bridging carboxylate oxygen atoms. To complete the coordination sphere, each calcium atom is further complexed to three water molecules. The net result is the formation of an eight-membered ring ($Ca_2C_2O_4$) between every adjacent calcium atom which ultimately leads to the formation of a one-dimensional polymeric chain of calcium atoms (see figure 1). The amino group of the anthranilate ligand does not directly take part in the coordination to the metal, but forms a hydrogen bond with the oxygen atoms of a coordinated water molecule from a neighbouring polymeric chain. In addition, O–H...O hydrogen bonds exist in-between the polymeric chains which originate from either the coordinated water molecules or the carboxylate oxygen atoms. Thus, the one-dimensional polymeric chains are connected with each other resulting in the formation of a supramolecular metal–organic framework as is seen in figure 2. The distance of 4.699(1) Å between the neighbouring calcium atoms along the chain is much longer than the observed value of 4.05 Å between the calcium atoms in the case of the related complex $[Ca(4-aminosalicylate)(acetate)(H_2O)H_2O]^6$. The coordination geometry around the calcium atom is irregular and can, at best, be described as a heavily distorted pentagonal bipyramidal, with the angles around calcium varying from 73.7 to 163.9°.

Crystal structure of 3: In spite of identical elemental composition, the solid-state structures of calcium complex **2** and $[Sr(2-aba)_2(OH)_2] \cdot H_2O$ (**3**) completely differ from each other. For example, the strontium atoms in **3** are nine-coordinated with a weak metal–metal interaction. The two types of anthranilate ligands in **3** show an entirely different coordinating behaviour (figure 3). Like in **2**, strontium atoms in **3** also form a zigzag polymeric chain with the help of these two contrastingly different anthranilate ligands. The most important observation to note here is the ability of the amino group of one of the anthranilate ligands to take part in coordination with neighbouring strontium atom. Moreover, the hydrogen-bonding interaction in **3** becomes more complex due to the presence of an additional lattice water molecule apart from the three coordinated water molecules. There are as many as nine different O–H...O, N–H...O, C–H...O

hydrogen bonds which are responsible for the supramolecular assembly depicted in figure 4. In general, the coordination geometry, Sr–O distances, and O–Sr–O angles in **3** are




Figure 1. Illustration of the polymeric chain formed by $[\text{Ca}(\text{2-aba})_2(\text{OH})_2]_n$ (**2**) in solid-state.




Figure 2. Packing diagram showing hydrogen bonding interactions between polymeric chains in $[\text{Ca}(2\text{-aba})_2(\text{OH}_2)_3]_\infty$ (2).




Figure 3. Illustration of the polymeric chain formed by $[\{\text{Sr}(2\text{-aba})_2(\text{OH}_2)_2\} \cdot \text{H}_2\text{O}]_\infty$ (3) in solid-state.




Figure 4. Packing diagram showing hydrogen bonding interactions between polymeric chains in $[\{\text{Sr}(2\text{-aba})_2(\text{OH}_2)_2\}\cdot\text{H}_2\text{O}]_\infty$ (**3**).

comparable to those found in $[\text{Sr}(\text{L-glutamate})\cdot 6\text{H}_2\text{O}]$ complex reported by Schmidbauer *et al*⁷ and other Sr-carboxylate complexes^{8,9}, although no amino coordination was found in the glutamate complex. Another important structural feature in **3** is the rather unusually short Sr...Sr distance of 3.92 Å.

Crystal structure of 4: The barium atoms in **4** are nine-coordinated as in **3**, although the coordination environment around the barium atom varies significantly. Each barium atom is coordinated by six carboxylate oxygen atoms, two bridging water molecules and one amino group. Two chelate rings are found around each barium atom; the first is a four-membered chelate ring formed by the carboxylate group of one of the anthranilate ligands, while the second is a six-membered chelate ring that is formed by the carboxyl atom and the amino nitrogen. Apart from chelating, the two anthranilate ligands are also engaged in bridging the neighbouring barium atoms. The net result is the formation of a two-dimensional polymer, which is schematically represented in figure 5. The water molecule bridges the two adjacent barium atoms along the polymeric chain. Due to the presence of the coordinated water molecule and the coordinated and free amino groups, a few hydrogen bonds are found in the solid state.




Figure 5. Schematic diagram of the coordination network in $[\text{Ba}(2\text{-aba})_2(\text{OH}_2)]_\infty$ (4).

Thus, a rich coordination chemistry of the 2-aminobenzoic acid with group 2 metal ions, which are normally reluctant to form coordination compounds, has been unraveled. The observed modes of coordination of the 2-amino benzoate ligand, are illustrated in figure 6.

2.2 Coordination polymers derived from 4-aminobenzoic acid¹⁰

While the relative ortho positioning of the $-\text{NH}_2$ and COO^- groups in 2-aba led to the isolation of solids **1–4**, it was expected that the use of 4-amino benzoic acid (4-abaH) would lead to novel layered-type structures due to the presence of potentially hydrogen bond forming $-\text{NH}_2$ group in the para position. In order to verify this supposition, the synthesis of compounds **5** and **6** were accomplished by the reaction of $\text{MCl}_2 \cdot 6\text{H}_2\text{O}$ ($\text{M} = \text{Mg}$ or Ca) with 4-abaH¹⁰. The analytically pure products were characterized by their IR, ^1H NMR and UV-vis spectral data. The aqueous solutions of both **5** and **6** were found to be almost neutral ($\text{pH} = 7.1$ for **5** and 6.5 for **6** at 0.06 M). The IR spectra show several strong absorptions in the region $3200\text{--}3500\text{ cm}^{-1}$ indicating the presence of water molecules with varying degrees of hydrogen bonding. Further, in case of **1**, a fairly large change in the observed $\nu(\text{C}=\text{O})$ value ($\sim 50\text{ cm}^{-1}$) is suggestive of the presence of uncoordinated aminobenzoates which are involved in only a secondary interaction to the Mg^{2+} ion (as in the recently reported 1-methylcytosine complexes of Mg and Cd)¹¹.




Figure 6. Schematic depiction of the coordination behaviour of the 2-aba ligands in complexes 2, 3 and 4.




Figure 7. Packing diagram of $[\text{Mg}(\text{H}_2\text{O})_6][(\text{4-aba})_2]\cdot 2\text{H}_2\text{O}$ (5) in the solid state showing alternating organic and inorganic layers.

Crystal structure of 5: The structure of $[\text{Mg}(\text{H}_2\text{O})_6][(\text{4-aba})_2] \cdot 2\text{H}_2\text{O}$ **5** comprises of an hexa-aqua Mg^{2+} ion which has only secondary interactions with two 4-aminobenzoate anions through the coordinated H_2O molecules. The Mg^{2+} ion forms a nearly ideal octahedral hexa-aqua complex. The three different $\text{Mg}-\text{OH}_2$ distances observed in the molecules (2.070(3), 2.071(3), and 2.096(3) Å) is comparable to the corresponding distances observed for $[\text{Mg}(\text{salicylate})_2(\text{OH}_2)_4]$, $[\text{Mg}(\text{2-aminosalicylate})_2(\text{OH}_2)_4]$, and $[\text{Mg}(\text{2,6-pyridinedicarboxylate})(\text{OH}_2)_3](\text{OH}_2)$. The view down b axis, as shown in figure 7, illustrates the supramolecular assembly of alternating columns of cationic $[\text{Mg}(\text{H}_2\text{O})_6]^{2+}$ and anionic 4-aminobenzoate ligands. The extensive hydrogen bonding between the Mg-bound water molecules, 4-aminobenzoate anions and the lattice water molecules hold the cationic and the anionic part of the molecule together. In the small pores produced by the formation of these alternating layers of cationic and anionic columns, lattice water molecules are trapped inside.

Crystal structure of 6: The basic structural motif in $[\text{Ca}(\text{4-aba})_2(\text{H}_2\text{O})_2]$ (**6**) is made up of a coordination polymer of eight-coordinated calcium ions as against the hepta-coordinated calcium ions found in the 2-aminobenzoate complex $[\text{Ca}(\text{2-aba})_2(\text{OH}_2)_3]_\infty$. Each Ca^{2+} ion is bound to two 4-aminobenzoate anions and two water molecules. One of the aminobenzoate ligands, apart from acting as a chelating ligand through its carboxylate group, also bridges the neighbouring Ca^{2+} ions on either side of the polymeric chain (figure 8). As a net result, the Ca^{2+} ions form infinite zigzag chains. A view down the axis




Figure 8. Illustration of the polymeric chain formed by $[\text{Ca}(\text{4-aba})_2(\text{OH}_2)_2]_\infty$ (**6**) in solid-state.




Figure 9. Packing diagram showing hydrogen bonding interactions between polymeric chains in $[\text{Ca}(4\text{-aba})_2(\text{OH}_2)_2]_\infty$ (**6**).

of the polymer propagation reveals that each of these chains is surrounded by six similar chains which are held together by hydrogen bonds (figure 9). Further, the cooperative hydrogen bonding between the $-\text{NH}_2$ group of the organic ligand and the coordinated H_2O molecule of the adjacent Ca^{2+} ions manifests into the formation of small π - π stacks of the aryl rings throughout the lattice.

It should be noted that, apart from the change in the coordination number, there are principal differences between **2** and the corresponding 2-aminobenzoate complex $[\text{Ca}(2\text{-aba})_2(\text{OH}_2)_3]_\infty$. For example, while the 2-aminobenzoate ligands in $[\text{Ca}(2\text{-aba})_2(\text{OH}_2)_3]_\infty$ exclusively function as bridging ligands between the adjacent calcium ions, both the 4-aminobenzoate ligands in **2** chelate to the metal apart from one these ligands also bridging the adjacent calcium atoms.




Figure 10. Packing of 2,2'-dithiobis-benzoic acid and mono-ammonium salt (adduct 7) in the solid-state.

(expected value 180°) with respect to each other. The observed values rather indicate an approximate perpendicular disposition of the aryl rings with respect to each other (see figure 10). The formation of these five-membered rings through the non-bonded contacts probably influences the observed C–S–S–C dihedral angles and hence the conformation of the molecule. The presence of the NH_4^+ ion is the source of extensive hydrogen bonds in the lattice. Apart from the O–H...O hydrogen bonding between the adjacent neighbours, additional N–H...O and N–H...S hydrogen bonds are formed to eventually result in a supramolecular array of disulphide adducts.

Crystal structure of 8: The calcium complex of 2,2'-dithiobis(benzoic acid) $\text{Ca}(\text{OOC}-\text{C}_6\text{H}_4-\text{S}-\text{S}-\text{C}_6\text{H}_4-\text{COO})(\text{H}_2\text{O})_2)_n$ forms a perfect coordination polymer (figure 11). The two carboxylate anions of the 2,2'-dithiobis(benzoate) ligand act as a bridge between polymeric chains of calcium ions. Each of these carboxylate anions is involved both in chelating and bridging modes of coordination within the same calcium chain. Thus, a three-dimensional polymeric structure as shown in figure 11b is formed in the solid state with the formation of channels inside the lattice. The difference map is suggestive of the presence of some disordered ethanol molecules being trapped in these channels. Further characterization of this product by thermal analysis will clearly reveal the exact nature of the occluded species.





Figure 11. (a) A fragment of the polymeric chain formed by $[\text{Ca}(\text{OOC}-\text{C}_6\text{H}_4-\text{S}-\text{S}-\text{C}_6\text{H}_4-\text{COO})(\text{H}_2\text{O})_2]_n$ (**8**) in the solid-state; (b) its packing diagram.



Scheme 5.

2.4 Coordination polymers of 1,2,4,5-benzene tetracarboxylic acid¹⁵

As earlier described the use of divalent transition metal ions and benzene di- and tricarboxylic acids (scheme 1) as modular precursors in designing a number of new metal-organic polymeric solids prompted us to investigate the supramolecular structure forming capabilities of the 1,2,4,5-benzene tetracarboxylic acid (H_4BTEC). Although it may be argued that, due to steric reasons, it is unlikely that all four carboxyl groups of H_4BTEC would take part in co-ordination to the metal, even the presence of free $-\text{COOH}$ groups would lead to formation of new and novel types of porous solids aided by weak

hydrogen bonding. Keeping this in mind, we have studied the extended solid forming ability of H₄BTEC towards divalent transition metal ions such as Co²⁺, Ni²⁺, and Zn²⁺ in the presence of an added donor amine.




Figure 12. Illustration of the repeating unit in the coordination polymer $[\text{Zn}(\text{btec})(\text{OH}_2)_4]_n \cdot n(\text{C}_4\text{H}_{12}\text{N}_2) \cdot 4n\text{H}_2\text{O}$ (**11**).




Figure 13. Diagram depicting the metal–organic polymer and the organic amine layers alternation in $[\text{Ni}(\text{btec})(\text{OH}_2)_4]_n \cdot n(\text{C}_4\text{H}_{12}\text{N}_2) \cdot 4n\text{H}_2\text{O}$ (**10**).

The compounds $[M(\text{BTEC})(\text{OH})_2]_n \cdot n(\text{C}_4\text{H}_{12}\text{N}_2) \cdot 4n\text{H}_2\text{O}$ ($M = \text{Co}$, **9**, Ni , **10**, Zn , **11**) are obtained in good yields by a slow diffusion controlled reaction between an aqueous solution of a suitable transition metal salt and H_4BTEC , in the presence of piperazine hexahydrate (scheme 5)¹⁵. The single crystals (pink, Co ; green Ni ; colourless Zn), whose size is dependent on the rate of crystallization and the concentration of the initial solution, are like most organic solvents, completely insoluble in water and once again forcing us to employ only solid-state techniques to characterize the products (elemental analysis, IR and DR UV-vis spectral studies, thermal analysis and single crystal X-ray diffraction studies). The difference between the asymmetric (1618 cm^{-1}) and symmetric (1361 cm^{-1}) stretching frequencies in the IR spectrum is consistent with the unidentate terminal mode of co-ordination of the carboxylate groups. The diffuse reflectance UV-visible spectrum of the compounds are consistent with the expected values for the octahedral complexes of these metal ions¹⁶.

All the three compounds are isomorphous and crystallize in the triclinic centrosymmetric space group. The molecular structure depicting the repeating unit in the




Figure 14. Three-dimensional packing of $[\text{Co}(\text{btec})(\text{OH})_2]_n \cdot n(\text{C}_4\text{H}_{12}\text{N}_2) \cdot 4n\text{H}_2\text{O}$ (**9**) in the solid-state resulting in the formation of micro-channels. Lattice water molecules omitted for clarity.




Figure 15. Supramolecular architecture in $2[\text{C}_6\text{H}_3(\text{COOH})_2(\text{COO})][\text{piperazine}\cdot\text{H}_2\text{O}]$ (**12**).

lattice of the zinc derivative is seen in figure 12. The central metal ion is octahedrally coordinated. While four water molecules surround each metal ion in the form of a square plane, the axial positions are ligated by carboxylate anions coming from two different BTEC ligands (in other words, the BTEC ligands bridge adjacent metal ions through the 1,4-carboxylate ligands in a unidentate fashion), leading to the formation of a one-dimensional polymeric structure. These coordination polymeric chains stack one over the other with water molecules with the piperazine cations sandwiched between them yielding a layered structure (figure 13). The packing diagram of this layered structure shows the presence of micropores of the dimensions $\sim 4 \text{ \AA}$ within the crystal lattice (figure 14).

Thermal studies on the cobalt and nickel derivatives show a stepwise weight loss behaviour of the water molecules. The first weight loss occurring at $50\text{--}90^\circ\text{C}$ corresponds to the loss of two uncoordinated water molecules, while a further weight loss observed at $100\text{--}170^\circ\text{C}$ could be attributed to the loss of two more un-coordinated and two coordinated water molecules. In fact, heating pink coloured cobalt coordination polymer **9** at this temperature range results in the change of colour to blue, indicating the change of coordination geometry to tetrahedral. Diffuse reflectance visible spectrum of this heat-treated sample further testifies to the formation of a tetrahedral structure around cobalt ions. Exposing this dehydrated sample to moist air results in the tetrahedral complex reverting back to the original octahedral structure. A similar colour change was observed for the nickel case (green to greenish-yellow, and back).

Recently, our attempts to generate coordination polymers of BTC in the presence of piperazine resulted in the isolation of an all-organic polymeric compound $2[\text{C}_6\text{H}_3(\text{COOH})_2(\text{COO})][\text{piperazine}\cdot\text{H}_2]$ (**12**) with no metal ions, as the side product¹⁷. Figure 15 depicts the supramolecular architecture derived from H_3BTC and piperazine.

We are presently exploring the possibilities of extending this approach to incorporate other types of amines inside the channels formed by 1,3,5-benzenetetracarboxylic acid.

3. Conclusion

Generation of novel supramolecular structures based on coordination polymeric networks continues to be a fertile area of research and a number of new framework structures are being regularly reported in modern literature. By choosing amino carboxylic acids as organic templates, we have shown that a number of new structures with large structural variations can be realized by just changing the size and nature of the central metal ion. It has also been shown that the change of the position of the amino group (from ortho to para) results in large variations in the final structures. Similarly, our preliminary results on the use of H₄BTEC as an organic module is suggestive of the potential of this largely unexplored ligand in the synthesis of new coordination polymers. Our present efforts in this area centre around the incorporation of amines and other types of small guest molecules within the coordination polymeric structures formed by H₃BTC and H₄BTEC.

Acknowledgements

This work was supported by a research grant from the Council of Scientific and Industrial Research and Department of Science and Technology, New Delhi. RM acknowledges the Department of Atomic Energy, Mumbai, for a research award in Chemistry. We thank the DST-funded X-ray facility at IIT-Bombay, and Mr H G Schmidt, University of Göttingen for the single crystal X-ray diffraction data.

References

1. Hagrman P J, Hagrman D and Zubeita J 1999 *Angew. Chem., Int. Ed. Engl.* **38** 2638
2. Yaghi O M, Davis C E, Li H and Groy T L 1998 *Acc. Chem. Res.* **31** 474
3. Yaghi O M, Davis C E, Li G and Li H 1997 *J. Am. Chem. Soc.* **119** 2861
4. (a) Atwood J L, Groeneman R H and MacGillivray L R 1999 *Inorg. Chem.* **38** 208 (b) Yaghi O M, Li H and Groy T L 1996 *J. Am. Chem. Soc.* **118** 9096 (c) Choi H J and Suh M P 1998 *J. Am. Chem. Soc.* **120** 10622
5. Murugavel R, Karambelkar V V, Anantharaman G and Walawalkar M G 2000 *Inorg. Chem.* **39** 1381
6. Cole L B and Holt E M 1989 *Inorg. Chim. Acta* **160** 195
7. Schmidbaur H, Bach I, Wilkinson D L and Müller G 1989 *Chem. Ber.* **122** 1433
8. Poonia N S and Bajaj A V 1979 *Chem. Rev.* **79** 389
9. (a) Briaggman B and Oskarsom A 1977 *Acta Crystallogr.* **B33** 1900 (b) Palmer K J, Wong R Y and Lewis J C 1972 *Acta Crystallogr. Sect* **B28** 223
10. Murugavel R, Karambelkar V V, Anantharaman G and Walawalkar M G 2000 *Indian J. Chem. Section A* (in press)
11. Westerhausen H 1991 *Inorg. Chem.* **30** 90
12. Murugavel R, Anantharaman G and Walawalkar M G 1998 (unpublished results)
13. Block E, Eswarakrishnan V, Gernon M, Ofori-okai G, Saha C, Tang K and Zubieta J 1989 *J. Am. Chem. Soc.* **111** 658
14. Hogg D R 1979 In *Organic compounds of sulphur, selenium and tellurium* (Cambridge: University Press) **5** 153
15. Murugavel R, Krishnamurthy D and Sathiyendiran M 1999 (manuscript in preparation)
16. Lever A B P 1984 In *Inorganic electronic spectroscopy* (Amsterdam: Elsevier Science) p 480
17. Murugavel R and Krishnamurthy D 1999 (unpublished results)