

Mechanism of oxidation of L-methionine by iron(III)-1,10-phenanthroline complex – A kinetic study

P VANI*¹, K KRISHNA KISHORE¹, R RAMBABU² and
L S A DIKSHITULU²

¹Department of Inorganic and Analytical Chemistry, School of Chemistry,
Andhra University, Visakhapatnam 530 003, India

²Gayatri Engineering College, Madurawada, Visakhapatnam 530 041, India
e-mail: vani_chem.@yahoo.co.in

MS received 19 May 2001; revised 2 July 2001

Abstract. Kinetics and mechanism of oxidation of L-methionine by iron(III)-1,10-phenanthroline complex have been studied in perchloric acid medium. The reaction is first order each in iron(III) and methionine. Increase in [phenanthroline] increases the rate while increase in [HClO₄] decreases it. While the reactive species of the substrate is the zwitterionic form, that of the oxidant is [Fe(phen)₂(H₂O)₂]³⁺. The proposed mechanism leads to the rate law

$$\frac{d[\text{Fe}(\text{phen})_3^{2+}]}{dt} = \frac{k_2 K_4 K_3 K_2^2 [\text{Fe}^{\text{III}}] [\text{phen}]^2 [\text{Met}]}{(1 + K_1 [\text{H}^+]) ([\text{H}^+]^2 + K_4 K_3 K_2^2 [\text{phen}]^2)}$$

Keywords. Oxidation of methionine; iron(III)-phenanthroline complex.

1. Introduction

In continuation of our studies on the oxidation of methionine¹⁻³, one of the sulphur-containing essential amino acids, we now report the mechanism of its oxidation in HClO₄ medium by iron(III)-1,10-phenanthroline complex. It is interesting to note that while the protonated form of methionine was observed to be the reactive species in the manganese (III)¹ and bromate² oxidations, the zwitterionic form was found to be the active species in the oxidations by iron(III) in presence of 2,2'-bipyridyl³ or 1,10-phenanthroline. While both phenanthroline and bipyridyl form similar complexes with iron(III), the oxidation of methionine with iron(III)-phenanthroline complex is much faster compared to iron(III)-2,2'-bipyridyl. Further, the kinetics of oxidation of methionine with iron(III)-phenanthroline complex is studied in aqueous perchloric acid medium, unlike its oxidation with iron(III)-2,2'-bipyridyl which was carried out in presence of methanol.

2. Experimental

2.1 Preparation of solutions

A 0.2 mol dm⁻³ solution of L-methionine was prepared by dissolving L-methionine (Sarabhai Merck) in distilled water and its strength determined by the iodimetric

*For correspondence

method⁴. Iron(III) perchlorate solution was prepared by heating iron wire (E Merck, Pro Analyti grade) with concentrated perchloric acid. The moist solid thus obtained was dissolved in distilled water and a sufficient quantity of HClO₄ was added to prevent the hydrolysis of iron(III). The iron(III) content was determined by titrating it with EDTA using variamine blue B as indicator. The free acid present in iron(III) stock solution was determined by titrating with NaOH using phenolphthalein as indicator according to the method of Milburn *et al*⁵. A fresh 0.015 mol dm⁻³ solution of 1,10-phenanthroline was prepared by dissolving 1,10-phenanthroline (E Merck) in distilled water.


2.2 Kinetics procedure

The kinetic measurements were carried out at 50 ± 0.1°C in 0.06 mol dm⁻³ perchloric acid medium under the conditions [methionine] [phen] [iron(III)]. Since methionine and phenanthroline get protonated in acid medium, [H⁺] does not remain constant as the concentrations of these two species are altered in the kinetic runs. [H⁺] of the reaction mixture is therefore adjusted to the desired value in a trial run using dilute perchloric acid. The actual runs are then carried out duplicating these conditions in the reaction mixtures. The progress of the reaction is followed by measuring the absorbance of *tris*-phenanthroline iron(II) formed at 510 nm, using a Shimadzu double beam spectrophotometer UV-140 (Japan) with 1 cm cells. The temperature was kept constant using a SEW Lab liquid circulatory bath S-36. Rate constants were found to be reproducible within ± 4%.

3. Results and discussion

3.1 Stoichiometry

Known amounts of methionine were reacted completely with a known excess of iron(III) at 50°C in 0.06 mol dm⁻³ HClO₄ in presence of excess of phenanthroline. The stoichiometry of the reaction was found to correspond to the equation.



3.2 Product analysis

Product analysis was carried out by taking the reaction mixture containing 5.0 × 10⁻² mol dm⁻³ methionine, 6.0 × 10⁻² mol dm⁻³ HClO₄, 6.0 × 10⁻³ mol dm⁻³ phenanthroline and 8.0 × 10⁻⁵ mol dm⁻³ iron(III), in 50 cm³ solution, and allowing it to stand for 2 h at 50°C. Thereafter 10 cm³ of 0.4 mol dm⁻³ sodium bicarbonate was added and the solution stirred vigorously, followed by dropwise addition of benzoyl chloride solution until precipitation was complete. The precipitate obtained was identified as N-benzoyl methionine sulfoxide (m.p. 183°C), a derivative of methionine sulfoxide⁶.

Table 1. Effect of $[\text{Fe}^{\text{III}}]$, $[\text{Met}]$, $[\text{phen}]$, $[\text{H}^+]$ and ionic strength (μ) on the pseudo-first-order rate constant, $k\zeta$ at $50 \pm 0.1^\circ\text{C}$.

| $[\text{Fe}^{\text{III}}] \times 10^5$ (mol dm ⁻³) | $[\text{Met}] \times 10^2$ (mol dm ⁻³) | $[\text{Phen}] \times 10^3$ (mol dm ⁻³) | $[\text{H}^+] \times 10^2$ (mol dm ⁻³) | μ (mol dm ⁻³) | $k\zeta \times 10^4$ (s ⁻¹) |
|---|---|--|---|----------------------------------|--|
| 5.0 | 5.0 | 6.0 | 6.0 | 0.12 | 1.54 |
| 6.0 | 5.0 | 6.0 | 6.0 | 0.12 | 1.58 |
| 8.0 | 5.0 | 6.0 | 6.0 | 0.12 | 1.60 |
| 9.0 | 5.0 | 6.0 | 6.0 | 0.12 | 1.54 |
| 10.0 | 5.0 | 6.0 | 6.0 | 0.12 | 1.49 |
| 8.0 | 2.0 | 6.0 | 6.0 | 0.12 | 0.82 |
| 8.0 | 3.0 | 6.0 | 6.0 | 0.12 | 1.15 |
| 8.0 | 4.0 | 6.0 | 6.0 | 0.12 | 1.48 |
| 8.0 | 5.0 | 6.0 | 6.0 | 0.12 | 1.63 |
| 8.0 | 6.0 | 6.0 | 6.0 | 0.12 | 2.20 |
| 8.0 | 5.0 | 6.0 | 1.0 | 0.12 | 11.26 |
| 8.0 | 5.0 | 6.0 | 3.0 | 0.12 | 5.88 |
| 8.0 | 5.0 | 6.0 | 4.0 | 0.12 | 3.26 |
| 8.0 | 5.0 | 6.0 | 5.0 | 0.12 | 2.30 |
| 8.0 | 5.0 | 6.0 | 6.0 | 0.12 | 1.63 |
| 8.0 | 5.0 | 6.0 | 6.0 | 0.15 | 1.65 |
| 8.0 | 5.0 | 6.0 | 6.0 | 0.17 | 1.60 |
| 8.0 | 5.0 | 6.0 | 6.0 | 0.20 | 1.58 |
| 8.0 | 5.0 | 6.0 | 6.0 | 0.22 | 1.62 |
| 8.0 | 5.0 | 2.0 | 6.0 | 0.12 | 0.25 |
| 8.0 | 5.0 | 3.0 | 6.0 | 0.12 | 0.52 |
| 8.0 | 5.0 | 4.0 | 6.0 | 0.12 | 0.85 |
| 8.0 | 5.0 | 8.0 | 6.0 | 0.12 | 2.27 |
| 8.0 | 5.0 | 9.0 | 6.0 | 0.12 | 2.77 |

3.3 Product dependence

Iron(II) was found to have no effect on the rate of the reaction.

3.4 Iron(III) dependence

When the kinetic runs were made with isolation of iron(III) by taking methionine in excess, plots of $\log(A_\infty - A_t)$ versus time were found to be linear up to 75% of the reaction, indicating unit order with respect to iron(III). Furthermore, when $[\text{Fe}^{\text{III}}]$ was varied from $5.0\text{--}10.0 \times 10^{-5}$ mol dm⁻³ (table 1) the pseudo-first order rate constants remained constant, confirming first order dependence on $[\text{Fe}^{\text{III}}]$.

3.5 Methionine dependence

Reaction order with respect to methionine was determined by carrying out kinetic runs in 0.06 mol dm⁻³ HClO₄ in the presence of 0.006 mol dm⁻³ phenanthroline and varying the methionine concentration over the $2.0\text{--}6.0 \times 10^{-2}$ mol dm⁻³ range. Results thus obtained (table 1) show that the pseudo-first order rate constant, $k\zeta$ increases with increasing [methionine]. Further, the plot of k' versus [methionine] is a straight line passing through the origin, confirming first-order dependence on [methionine].

3.6 Hydrogen ion dependence

In order to study the effect of $[H^+]$ on the pseudo-first order rate constant, k' kinetic runs were carried out at various $HClO_4$ concentrations over $1.0\text{--}6.0 \times 10^{-2} \text{ mol dm}^{-3}$ range, keeping ionic strength and other reactant concentrations constant. The rate constants were found to decrease with increase in acid concentration (table 1).

3.7 Ionic strength dependence

The effect of ionic strength on the pseudo-first order rate constant, k' , was studied over the $0.12\text{--}0.22 \text{ mol dm}^{-3}$ range using $NaClO_4$, keeping the concentrations of all other species in the reaction mixture constant. The results recorded in table 1 show that the rate is independent of ionic strength.

3.8 Dependence on [phenanthroline]

Effect of phenanthroline on the pseudo-first order rate constant, k' , was studied at 50°C by varying the 1,10-phenanthroline concentration from 2.0×10^{-3} to $9.0 \times 10^{-3} \text{ mol dm}^{-3}$ keeping all other reactant concentrations constant. Rate increases with increase in phenanthroline concentration. Furthermore, plots of $1/k'$ versus $1/[\text{phen}]^2$ were linear with positive intercepts on the $1/k'$ axis. This suggests a Michaelis–Menten type of behaviour, with the formation of a 1 : 2 complex between iron(III) and 1,10-phenanthroline.

3.9 Thermodynamic parameters

Activation parameters were determined from the k' values obtained at the temperatures 45, 50 and 55°C (table 2) keeping the concentrations of all other ions constant: $[\text{Met}] = 5.0 \times 10^{-2} \text{ mol dm}^{-3}$; $[\text{Fe}^{\text{III}}] = 8.0 \times 10^{-5} \text{ mol dm}^{-3}$; $[H^+] = 6.0 \times 10^{-2} \text{ mol dm}^{-3}$; $m = 0.12 \text{ mol dm}^{-3}$ ($NaClO_4$). The energy of activation E_a , and the entropy of activation ΔS^\ddagger were calculated employing the linear least squares method and were $106.4 \pm 9.2 \text{ kJ mol}^{-1}$ and $28.5 \pm 28.5 \text{ J K}^{-1} \text{ mol}^{-1}$ respectively.











Phenanthroline has a pK_a value of 4.641 at 50°C and in the $[H^+]$ range investigated, phenanthroline exists mainly in the form of $H\text{phen}^+$ (> 98%). Baxendale and Bridge⁷ reported that a yellow complex is formed when iron(III) is mixed with 1,10-phenanthroline or 2,2'-bipyridyl and from a photometric study of the complex, concluded that the species has the formula $[\text{Fe}(\text{phen})_n]^{3+}$, the maximum value of n being 2.5 at very high phenanthroline concentrations. Baxendale⁸ carried out a photometric study of the brown complex formed by direct mixing of iron(III) and phenanthroline in aqueous

Table 2. Effect of temperature on the pseudo-first order rate constant, k' . $[\text{Fe}^{\text{III}}] = 8.0 \times 10^{-5} \text{ mol dm}^{-3}$; $[\text{Met}] = 5.0 \times 10^{-2} \text{ mol dm}^{-3}$; $[\text{phen}] = 6.0 \times 10^{-3} \text{ mol dm}^{-3}$; $[H^+] = 6.0 \times 10^{-2} \text{ mol dm}^{-3}$; $m = 0.12 \text{ mol dm}^{-3}$ ($NaClO_4$).


| Temperature | 318 | 323 | 328 |
|--|------|------|------|
| $k' \times 10^4 \text{ (s}^{-1}\text{)}$ | 0.77 | 1.58 | 2.63 |

solution and stated that the formula of the complex is probably $[\text{Fe}(\text{phen})_2]^{3+}$. Subsequently, considerable attention was paid to the elucidation of the structures of these brown complexes of iron(III) with 2,2'-bipyridyl and 1,10-phenanthroline. David and de Mello⁹ established the formation of $[\text{FeL}_2(\text{H}_2\text{O})_2]^{3+}$ from the interaction of the binuclear species with H^+ ions. In view of these reports, the formula of the iron(III)-1,10-phenanthroline complex in the present study may be considered to be $[\text{Fe}(\text{phen})_2(\text{H}_2\text{O})_2]^{3+}$. This complex is different from *tris*-1,10-phenanthroline iron(III), $[\text{Fe}(\text{phen})_3]^{3+}$, which is a blue compound that can only be prepared by the oxidation of the corresponding iron(II) complex¹⁰. While the blue complex is substitution inert (spin-paired), the yellow-brown complex formed by direct mixing of iron(III) and phenanthroline is labile (spin free).

All amino acids possess minimum two pK values, one corresponding to the carboxylic acid group ($pK_1 = 2.0$) and the other to the amino group ($pK_2 = 10.0$). Since methionine is a sulphur-containing amino acid, under the present experimental conditions $\{[\text{H}^+] = 6.0 \times 10^{-2} \text{ mol dm}^{-3}\}$ it exists in the form of the protonated species $\text{H}_3\text{CSCH}_2\text{CH}_2\text{CH}(\overset{+}{\text{N}}\text{H}_3)\text{COOH}(\text{HMet})$ to the extent of 85%, the remaining 15% being present in the zwitterionic form, $\text{H}_3\text{CSCH}_2\text{CH}_2\text{CH}(\overset{+}{\text{N}}\text{H}_3)\text{COO}^-(\text{Met}_2)$. Based on these observations, the following mechanism has been proposed.



fast
products, (12)



where $\text{R} = -\text{CH}_2\text{CH}_2\text{CH}(\overset{+}{\text{N}}\text{H}_3)\text{COO}^-$ and HR is its protonated form, $-\text{CH}_2\text{CH}_2\text{CH}(\text{NH}_3)\text{COOH}$, Met_z is the zwitterionic form and HMet is its protonated form.

$$\text{Rate} = \frac{d[\text{Fe(phen)}_3^{3+}]}{dt} = k_1[\text{C}_1][\text{Met}_z] + k_2[\text{C}_2][\text{Met}_z] + k_3[\text{C}_1][\text{HMet}] + k_4[\text{C}_2][\text{HMet}] \quad (14)$$

$$= \{k_1[\text{Met}_z] + k_3[\text{HMet}]\}[\text{C}_1] + \{k_2[\text{Met}_z] + k_4[\text{HMet}]\}[\text{C}_2], \quad (15)$$

where

$$[\text{C}_1] = K_3K_2[\text{Fe}^{\text{III}}]_e[\text{Hphen}^+]_e/[\text{H}^+]_e, \quad (16)$$

$$[\text{C}_2] = K_4K_3K_2^2[\text{Fe}^{\text{III}}]_e[\text{Hphen}^+]_e^2/[\text{H}^+]_e^2, \quad (17)$$

and

$$[\text{Fe}^{\text{III}}]_t = [\text{Fe}^{\text{III}}]_e + [\text{C}_1] + [\text{C}_2]. \quad (18)$$

Substituting for $[\text{C}_1]$ and $[\text{C}_2]$ from (16) and (17) in (18)

$$[\text{Fe}^{\text{III}}]_e = [\text{H}^+]_e^2[\text{Fe}^{\text{III}}]_t/[\text{H}^+]_e^2 + K_3K_2[\text{Hphen}^+]_e[\text{H}^+]_e + K_4K_3K_2^2[\text{Hphen}^+]_e^2. \quad (19)$$

Substituting for $[\text{Fe}^{\text{III}}]_e$ from (19) in (16) and (17) leads to

$$[\text{C}_1] = K_3K_2[\text{H}^+]_e[\text{Fe}^{\text{III}}]_t/[\text{H}^+]_e^2 + K_3K_2[\text{Hphen}^+]_e[\text{H}^+]_e + K_4K_3K_2^2[\text{Hphen}^+]_e^2, \quad (20)$$

and

$$[\text{C}_2] = K_4K_3K_2^2[\text{Fe}^{\text{III}}]_t/[\text{H}^+]_e^2 + K_3K_2[\text{Hphen}^+]_e[\text{H}^+]_e + K_4K_3K_2^2[\text{Hphen}^+]_e^2. \quad (21)$$

Similarly,

$$[\text{Met}]_t = [\text{Met}_z] + [\text{HMet}], \quad (22)$$

and from (1),

$$[\text{Met}_2] = [\text{Met}]_t / 1 + K_1[\text{H}^+]_e, \quad (23)$$

$$\text{and } [\text{HMet}] = K_1[\text{H}^+]_e [\text{Met}]_t / 1 + K_1[\text{H}^+]_e. \quad (24)$$

Finally, substituting for $[\text{C}_1]$, $[\text{C}_2]$, $[\text{Met}_2]$ and $[\text{HMet}]$ from (16), (17) and (24) in (15) we get

$$\begin{aligned} \text{rate} &= \frac{-d[\text{Fe}^{\text{III}}]}{dt} \\ &= \frac{\{(k_1 + k_3K_1[\text{H}^+]_e)K_3K_2[\text{H}^+]_e + (k_2 + k_4K_1[\text{H}^+]_e)K_4K_3K_2^2[\text{Hphen}^+]\}[\text{Fe}^{\text{III}}]_t [\text{Hphen}^+][\text{Met}]_t}{\{1 + K_1[\text{H}^+]_e\}\{[\text{H}^+]_e^2 + K_3K_2[\text{Hphen}^+]_e[\text{H}^+]_e + K_4K_3K_2^2[\text{Hphen}^+]_e^2\}}. \end{aligned} \quad (25)$$

Since the concentrations of phenanthroline and hydrogen ion are large compared to that of iron(III), $[\text{Hphen}^+]_e = [\text{phen}]$, and $[\text{H}^+]_e = [\text{H}^+]_t$, Equation (25), therefore, reduces to the form,

$$\text{rate} = \frac{\{(k_1 + k_3K_1[\text{H}^+])K_3K_2[\text{H}^+] + (k_2 + k_4K_1[\text{H}^+])K_4K_3K_2^2[\text{phen}]\}[\text{Fe}^{\text{III}}]_t [\text{phen}]_t [\text{Met}]}{\{1 + K_1[\text{H}^+]\}\{[\text{H}^+]^2 + K_3K_2[\text{phen}][\text{H}^+] + K_4K_3K_2^2[\text{phen}]^2\}}. \quad (26)$$

Ionic strength has negligible effect on the rate of the reaction, but increase in $[\text{H}^+]$ has considerable retarding effect. Since the $[\text{phen}]_e$ is not significantly altered in the $[\text{H}^+]$ range studied, retardation by the $[\text{H}^+]$ may be mainly attributed to the conversion of the more reactive neutral species of methionine into the far less reactive protonated form. In view of this, k_3 and k_4 may be presumed to be far less than k_1 and k_2 respectively. Hence, $k_3K_1[\text{H}^+]$ and $k_4K_1[\text{H}^+]$ may be neglected in comparison with k_1 and k_2 respectively in the numerator of (26). Therefore, the rate equation reduces to

$$\text{rate} = \frac{K_3K_2[\text{Fe}^{\text{III}}][\text{phen}][\text{Met}](k_1[\text{H}^+] + k_2K_4K_2[\text{phen}])}{(1 + K_1[\text{H}^+])([\text{H}^+]^2 + K_3K_2[\text{phen}][\text{H}^+] + K_4K_3K_2^2[\text{phen}]^2)}. \quad (27)$$


The rate equation explains the observed first order with respect to iron(III) and methionine, as well as the increase in rate with $[\text{phen}]$ and decrease in rate with $[\text{H}^+]$. But this equation does not predict any simple graphical relationship between the observed rate constant, $k^1(\text{rate}/[\text{Fe}^{\text{III}}])$ and either $[\text{H}^+]$ or $[\text{phen}]$. However, our experimental results indicate that the plot of $1/k^1$ versus $1/[\text{phen}]^2$ is linear with a positive intercept on the rate axis. Such a plot may be expected from the rate equation if only the terms $k_1[\text{H}^+]$ in the numerator and $K_3K_2[\text{H}^+][\text{phen}]$ in the denominator are neglected. In such a case, the rate equation reduces to

$$\frac{-d[\text{Fe}(\text{phen})_3^{2+}]}{dt} = \frac{k_2K_4K_3K_2^2[\text{Fe}^{\text{III}}][\text{phen}]^2[\text{Met}]}{(1 + K_1[\text{H}^+])([\text{H}^+]^2 + K_4K_3K_2^2[\text{phen}]^2)}, \quad (28)$$

thereby implying negligible participation of the 1:1 complex of phenanthroline (C_1) in the reaction. The main reactive species in the reaction may therefore be regarded as $[\text{Fe}(\text{phen})_2(\text{H}_2\text{O})_2]^{3+}$ and the zwitterionic form of methionine (Met_z).

4. Conclusion

Methionine is a sulphur containing amino acid and it has three coordinating centres, oxygen, nitrogen and sulphur. The oxidation product of methionine depends on the nature of the oxidant. With some oxidants it is oxidized to the sulphoxide stage only, whereas with certain others it is oxidized to the sulphone stage. In all these oxidations electrophilic attack was reported to take place at the sulphur site (sulphur being more nucleophilic than nitrogen and oxygen). The protonated form of methionine is the reactive species in its oxidation by manganese(III)¹ and bromate². But the oxidation of methionine by iron(III)-2,2'-bipyridyl³ or 1,10-phenanthroline has been found to be extremely slow in acid media. However, the rate increases with decrease in $[\text{H}^+]$ thereby indicating that the zwitterionic form is far more reactive than the protonated form. This might be due to the ability of the zwitterionic form of methionine to coordinate through the negative carboxylic oxygen also, thereby stabilizing the transition state through the formation of a completely chelated iron(III) complex. While both iron(III)-phenanthroline and iron(III) 2,2'-bipyridyl oxidize methionine to sulphoxide, oxidation with the former oxidant is much faster compared to the latter. This is because of the higher oxidation potential of the iron(III)-phenanthroline complex compared to the iron(III)-2,2'-bipyridyl. Hence the intimate mechanism of oxidation of methionine by iron(III) in presence of phenanthroline may be represented as in scheme 1.



Scheme 1.

References

1. Vani P and Rajeswari T R 1994 *React. Kinet. Catal. Lett.* **52** 175
2. Vani P, Rajeswari T R and Dikshitulu L S A 1995 *J. Indian Chem. Soc.* **72** 867
3. Rajeswari T R, Vani P and Dikshitulu L S A 1995 *Transition Met. Chem.* **20** 170
4. Gensch K H and Higuchi T 1967 *J. Pharm. Sci.* **56** 177
5. Milburn R M and Vosburgh W C 1955 *J. Am. Chem. Soc.* **77** 1352
6. Goswami K B, Chandra G and Srivastava S N 1981 *J. Indian Chem. Soc.* **58** 252
7. Baxendale J H and Bridge N K 1955 *J. Phys. Chem.* **59** 783
8. Baxendale J H 1952 *Advances in catalysis and related subjects* (New York: Academic Press) vol. 4, p. 64
9. David P C and de Mello P C 1973 *Inorg. Chem.* **12** 2188
10. Brandt W W, Dwyer F P and Gyrfas E C 1954 *Chem. Rev.* **54** 959