

Chemistry Department, APS University, Rewa 486 003, India

[†]Present address: UP Public Service Commission, Allahabad 211 001, India

MS received 26 June 2001; revised 26 November 2001

Abstract. Speciation has been determined in aqueous oxovanadium, glycylvaline and imidazoles at $25 \pm 1^\circ\text{C}$ and $m = 0.1\text{M}$ NaClO_4 using a combination of potentiometry, and visible and EPR spectroscopy. Results of potentiometric and spectroscopic methods are consistent. Calculations of stability constants have been made using the SCOGS computer program.

Keywords. Oxovanadium (IV); glycylvaline; imidazoles.

1. Introduction

The aqueous chemistry of vanadium has been attracting interest for many years. Solution equilibria involving oxovanadium can be followed very conveniently using potentiometry and EPR spectroscopy. The importance of histidyl residues makes imidazole a relevant target for metal binding studies in biomolecules. Further, only limited information is available on oxovanadium(IV) ternary complexes¹⁻⁴ and oxovanadium (IV) complexes with peptides^{5,6}, aminoacids⁷⁻¹⁵ and imidazoles¹⁶. There are also examples¹⁷⁻²⁰ with other simple molecules containing this type of donor groups. With this view in mind, we have studied oxovanadium (IV) ternary systems using glycylvaline (A) (I) and three imidazoles (B) (II) viz., imidazole (ImH), 2-methylimidazole (M-ImH) and 2-ethylimidazole (E-ImH), using three different techniques in aqueous solution.

2. Experimental



2.1 Chemicals

Vanadyl sulphate (Aldrich), glycylvaline (Aldrich) and imidazoles (s.d. Fine Chem.) were used as received. All other chemicals used were of reagent grade. Solutions were prepared using double distilled water.

2.2 Potentiometric measurements

Values of *pH* were determined with a Systronics *pH* meter, model 335 (accuracy ± 0.01 *pH* units) using special glass electrodes (*pH* 1–14) and saturated calomel electrodes.

*For correspondence



- (a) R = H, Imidazole (ImH)
 (b) R = -CH₃, 2-methylimidazole (M-ImH)
 (c) R = -CH₂CH₃, 2-ethylimidazole (E-ImH)

Titration procedures were the same as described in our earlier work²¹⁻²⁶. The following metal–ligand mixtures were prepared for the *pH*-potentiometric study:

- (i) 0.03 M perchloric acid + 0.1 M NaClO₄;
- (ii) 0.03 M perchloric acid + 0.003 M A + 0.1 M NaClO₄;
- (iii) 0.03 M perchloric acid + 0.003 M B + 0.1 M NaClO₄;
- (iv) 0.03 M perchloric acid + 0.003 M VO²⁺ + 0.003 M A + 0.1 M NaClO₄;
- (v) 0.03 M perchloric acid + 0.003 M VO²⁺ + 0.003 M B + 0.1 M NaClO₄;
- (vi) 0.03 M perchloric acid + 0.003 M VO²⁺ + 0.003 M A + 0.003 M B + 0.1 M NaClO₄.

Here A = glycylvaline and B = imidazoles. The initial volume of all the solutions was 50 ml and the required amount of NaClO₄ was added to each solution to maintain the ionic strength (*m* = 0.1 M NaClO₄). All the solutions were allowed to attain equilibrium at 25 ± 1°C and were titrated with 1 M NaOH.

2.3 Spectrophotometric measurements

Visible absorption spectra were recorded as functions of *pH* in aqueous solution using a Systronics UV-Vis 117 spectrophotometer and 1 cm quartz cells.

2.4 EPR measurements

EPR spectra were recorded with a Varian E-line Century Series EPR spectrometer equipped with dual cavity and operating with X-band with 100 kHz modulation frequency. TCNE (*g_e* = 2.00277) was used as the field marker.

3. Results and discussion

Protonation constants of ligands were determined by Calvin–Bjerrum's^{27,28} technique as adopted by Irving and Rossotti²⁹. The formation constants of different binary and ternary complexes were evaluated using the SCOGS computer program³⁰. Protonation constants



The equilibrium constants of the above equations are given in table 1. These hydroxo species are not seen in the species distribution curves (figures 1 and 2) as their percentage is less than 1%.

Table 1. Proton–ligand and metal–ligand binary and ternary constants at $25 \pm 1^\circ\text{C}$ in aqueous solution, ionic strength $I = 0.1 \text{ M NaClO}_4$ (standard deviations are ± 0.02 in log unit).

(a) *Proton–ligand constants*

Ligand	pK_2^{H}	pK_1^{H}
Glycylvaline (AH_2)	3.15	8.25
Imidazole (BH^+)	–	7.10
2-Methylimidazole (BH^+)	–	8.00
2-Ethylimidazole (BH^+)	–	8.00

(b) *Hydrolytic constants of VO^{2+} aq. ions*

$\log b_{\text{VO}}^{\text{H}}$	–6.00
$\log b_{\text{VO}}^{2\text{H}}$	–12.88

(c) *Stability constants ($\log b$) of binary complexes*

	GlyVal	ImH	2-M-ImH	2-E-ImH
VO(AH)	7.82	–	–	–
VO(AH ₋₁)	3.70	–	–	–
VO(B)	–	5.77	6.66	6.66

(d) *Stability constants ($\log b_{\text{VO(A)(B)}}^{\text{VO}}$) of ternary complexes*

A	B	$\log b_{\text{VO(A)(B)}}^{\text{VO}}$	$\Delta \log K_{\text{VO}}$
GlyVal	ImH	15.47	+ 1.88
	2-M-ImH	15.02	+ 0.54
	2-E-ImH	15.00	+ 0.52




Figure 1. Species distribution curves of M:AH (1:1) binary system: (1) AH_2 , (2) AH , (3) $M(OH)^+$, (4) $M(OH)_2$, (5) MAH and (6) MAH_{-1} .




Figure 2. Species distribution curves of M:AH:BH (1:1:1) ternary system: (1) AH_2 , (2) AH , (3) BH^+ , (4) $M(OH)^+$, (5) $M(OH)_2$, (6) MAH , (7) MB , (8) MAB and (9) $MABH_{-1}$.

Stability constants (table 1) and speciation curves (figures 1 and 2) were obtained from the SCOGS (stability constants of generalized species) computer program. Complex formation equilibria have been illustrated on the basis of the concentration distributions of the complexes formed in the VO^{2+} : A binary and VO^{2+} : A:B ternary systems in aqueous solutions. Stability of the ternary $VO(A)(B)$ complex was generally characterized on the basis of $\Delta \log K_{VO}$ values (table 1) calculated using relation³²⁻³⁴,

$$\Delta \log K_{VO} = \log b_{VO(A)(B)}^{VO} - \log b_{VO(A)}^{VO} - \log b_{VO(B)}^{VO}. \quad (3)$$



3.1 Stability constants with respect to imidazoles

Stability constants of ternary VO(A)(B) complexes were found to be in the order: imidazole < 2-methylimidazole < 2-ethylimidazole. As we pass from imidazole to 2-ethylimidazole, an increase in $\log b$ value is observed, which is due to increase in the basicity of the alkyl-substituted imidazoles. From steric considerations, the $\log b$ values in case of complexes with substituted imidazoles should be smaller than that for imidazole complexes. The increase in basicity of substituted imidazoles thus compensates for the negative contribution of the steric effect and further enhances the values of $\log b$ in all the cases. Among the methyl- and ethylimidazoles, there is not much difference either in ligand basicities or in steric effects so that both methyl- and ethylimidazole give almost similar $\log \beta$ values.

3.2 EPR studies

X-band EPR spectra at liquid nitrogen temperature may be simulated as axial spectra³⁵. The field region⁷ corresponding to $A_{||}$ and $M_I = 5/2$ and $7/2$ gives more information about the type and number of species. We have recorded the EPR spectra of binary and ternary systems as functions of pH. The derived EPR parameters are collected in table 2. Some representative EPR spectra are shown in figure 3. In the case of binary systems, the spectrum at pH 2.50 is characteristic of aquo vanadium. When pH is increased from 2.50 to 3.50 the peaks shift slightly to lower fields, hence more than one species contributes to each spectrum. At higher pH values, distinct species are detected, and signal intensity weakens significantly. In the case of ternary systems, at the lowest pH (i.e. pH 2.50) studied the spectrum is not due to aquo VO^{2+} ion. It could be due to the binary species VO(AH), as its EPR parameters resemble that of the binary species detected in $\text{VO}^{2+} + \text{A}(1:1)$ systems. When pH is increased from 2.50 to 3.50 the peak again shifts to lower fields, hence more than one species contributes to each spectrum. At higher pH values, signal intensities differ from that of binary high pH spectra. Change in signal intensity could be due to ternary species formation.

By applying the additivity relationship³⁶ to the species assumed to be present at different pH values, the $A_{z,i}$ values for different coordinate sites were calculated and are given in table 2. These calculated values for the equatorial sites are almost identical to the experimental values. Holyk³⁷ plotted the relation (figure 4) between $A_{||}$ and $g_{||}$ values for vanadyl complexes and defined zones for complexes having different equatorial donor atom sites, such as VO(N₄), VO(N₂O₂) and VO(O₄). In this plot, our data points shift from the main domains. This kind of shifting relative to the reference data implies that

Table 2. EPR parameters, proposed equatorial coordination and λ_{\max} for different binary and ternary oxovanadium (IV) systems.

Composition	Complex	pH	g_{\parallel}	g_{\perp}	A_{\parallel} (10^{-4} cm^{-1})	A_{\perp} (10^{-4} cm^{-1})	Proposed equatorial coordination	A_{zi} (10^{-4} cm^{-1})		λ_{\max} (nm)
								Calc. value	Lit. value	
VO ²⁺ :A (1:1)	VO ²⁺ aq.	2.50	1.931	1.972	183	72	4H ₂ O	45.9	45.7	~ 772
	VO(AH)	3.50	1.928	1.976	174	62	R-NH ₂ R-CO ₂ 2H ₂ O	39.9 42.5 45.5	40.1 42.7 45.7	~ 760
VO ²⁺ :A:ImH (1:1:1)	VO(AH ₁)	5.50	1.931	1.978	160	49	R-NH ₂ R-CO ₂ H ₂ O	39.6 42.2 45.2	40.1 42.7 45.7	ill- defined
							CONH	31.5	32.0	
	VO(AH)	2.50	1.933	1.981	174	63	R-NH ₂ R-CO ₂ 2H ₂ O	39.9 42.5 45.5	40.1 42.7 45.7	~ 760
							R-NH ₂ R-CO ₂ =N-	42.9 38.5 45.9	40.1 38.3 45.7	~ 766
VO(A)(B)	5.00	1.935	1.984	153	41	H ₂ O R-NH ₂ R-CO ₂ =N-	40.0 42.6 38.2	40.1 42.7 38.3	ill- defined	
						CONH	31.9	32.0		
VO(A)(B)	6.00	1.932	1.980	153	40	R-NH ₂ R-CO ₂ =N-	40.0 42.6 38.2	40.1 42.7 38.3	580, -	
						CONH	31.9	32.0		

(continued)

Table 2. (Continued).

VO ²⁺ :A:M-ImH (1:1:1)	VO(AH)	2.50	1.928	1.980	174	64	R-NH ₂	39.9	40.1	632, 768
							R-CO ₂	42.5	42.7	
							2H ₂ O	45.5	45.7	
	VO(A)(B)(H ₂ O)	3.50	1.928	1.978	167	57	R-NH ₂	40.3	40.1	-, 760
							R-CO ₂	42.9	42.7	
							=N-	38.5	38.3	
							H ₂ O	45.9	45.7	
	VO(A)(B)	5.00	1.930	1.950	153	45	R-NH ₂	40.0	40.1	ill-
							R-CO ₂	42.6	42.7	defined
							=N-	38.2	38.3	
VO ²⁺ :A:E-ImH (1:1:1)	VO(A)(B)	6.00	1.928	1.967	153	42	CONH	31.9	32.0	
							R-NH ₂	40.0	40.1	ill-
							R-CO ₂	42.6	42.7	defined
							=N-	38.2	38.3	
	VO(AH)	2.50	1.928	1.967	174	64	CONH	31.9	32.0	
							R-NH ₂	39.9	40.1	635, 765
							R-CO ₂	42.5	42.7	
							2H ₂ O	45.5	45.7	
	VO(A)(B)(H ₂ O)	3.50	1.930	1.950	167	58	R-NH ₂	40.3	40.1	-, 762
							R-CO ₂	42.9	42.7	
						=N-	38.5	38.3		
						H ₂ O	45.9	45.7		
VO(A)(B)		5.00	1.928	1.980	153	41	R-NH ₂	40.0	40.1	ill-
							R-CO ₂	42.6	42.7	defined
							=N-	38.2	38.3	
							CONH	31.9	32.0	
	VO(A)(B)	6.00	1.928	1.978	153	42	R-NH ₂	40.0	40.1	ill-
							R-CO ₂	42.6	42.7	defined
							=N-	38.2	38.3	
							CONH	31.9	32.0	
							R-NH ₂	40.0	40.1	ill-
							R-CO ₂	42.6	42.7	defined
						=N-	38.2	38.3		
						CONH	31.9	32.0		




Figure 3. X-band EPR spectra of M:AH:ImH (1:1:1) ternary system as a function of pH.




Figure 4. A correlation plot of $A_{||}$ vs. $g_{||}$ for the VO^{2+} complexes: for (●) N_3O , N_2O_2 , NO_3 , O_4 (points of the present study) and (.) in various domains for a series of oxovanadium (IV) complexes with different donor atoms²².

the physical mechanism which determines the $A_{||}$ values in our complexes is not the same as in reference compounds.

3.3 Spectrophotometric study

Visible absorption spectra of the oxovanadium (IV) binary and ternary systems were recorded as a function of pH. Values of I_{\max} obtained are shown in table 2. Values of I_{\max} are not given for all binary and ternary species owing to ill-defined band positions. Band positions of binary and ternary species differ.

4. Conclusion

In biological systems, the available ligand atoms are S, N, and O. The oxovanadium (IV) is a powerful paramagnetic probe of protein structure and therefore it is important to ascertain the nature of its aminoacid, peptide and protein complexes. $A_{\lambda,i}$ values are identical to the literature values.

Acknowledgement

This work was carried out under financial assistance from the University Grant Commission, New Delhi.

References

1. Elvingson K, Keramidas A D, Crans D C and Pettersson L 1998 *Inorg. Chem.* **37** 6153
2. Elvingson K, Crans D C and Pettersson L 1997 *J. Am. Chem. Soc.* **119** 7005
3. Crans D C, Schelble S M and Theisen L A 1991 *J. Org. Chem.* **56** 1266
4. Galeffi B and Tracey A S 1989 *Inorg. Chem.* **28** 1726
5. Pessoa J C, Luz S M, Duarte R, Moura J J G and Gillard R D 1993 *Polyhedron* **12** 2857
6. Pessoa J C, Gajada T, Gillard R D, Kiss T, Luz S M, Moura J J G, Tornaz I, Telo J P and Torok I 1998 *J. Chem. Soc., Dalton Trans.* 3587
7. Pessoa J C, Boas L F V, Gillard R D and Lancashire R J 1988 *Polyhedron* **7** 1245
8. Pessoa J C, Luz S M, Cavaco I and Gillard R D 1994 *Polyhedron* **13** 3177
9. Pessoa J C, Boas L F V and Gillard R D 1990 *Polyhedron* **9** 2101
10. Pessoa J C, Marques R L, Boas L F V and Gillard R D 1990 *Polyhedron* **9** 81
11. Pessoa J C, Antunes J L, Boas L F V and Gillard R D 1992 *Polyhedron* **11** 1449
12. Pessoa J C, Boas L F V and Gillard R D 1989 *Polyhedron* **8** 1173
13. Pessoa J C, Boas L F V and Gillard R D 1989 *Polyhedron* **8** 1745
14. Tomiyasu H and Gordon G 1973 *J. Coord. Chem.* **3** 47
15. Fabian I and Nagypal I 1982 *Inorg. Chim. Acta* **62** 193
16. Sanna D, Micera G, Erre L S, Molinu M G and Garribba E 1996 *J. Chem. Res.* 40
17. Felceman J and Da Silva F J J R 1983 *Dalton* **8** 565
18. Alberico E, Micera G and Sanna D 1994 *Polyhedron* **13** 1763
19. Nagypal I and Fabian I 1982 *Inorg. Chim. Acta* **61** 109
20. Jain A K and Chaturvedi G K 1979 *J. Indian Chem. Soc.* **56** 850
21. Pandeya K B and Patel R N 1990 *Indian J. Chem.* **A29** 602
22. Pandeya K B and Patel R N 1991 *Indian J. Chem.* **A30** 193
23. Patel R N, Pandey H C and Pandeya K B 1999 *Indian J. Chem.* **A38** 850
24. Patel R N, Singh N, Shrivastava R P, Kumar S and Pandeya K B 2000 *J. Mol. Liq.* **89** 207
25. Patel R N, Shrivastava R P, Singh N and Pandeya K B 2000 *Proc. Natl. Acad. Sci., India* **70** 133
26. Patel R N, Shrivastava R P, Singh N and Pandeya K B 2001 *Indian J. Chem.* **A40** 361
27. Calvin H and Wilson K W 1945 *J. Am. Chem. Soc.* **67** 2003
28. Bjerrum J 1941 *Metal amine formation in aqueous solutions* (Copenhagen: P Hasse)
29. Irving H M and Rossotti H S 1954 *J. Chem. Soc.* 2904
30. Sayce I G 1968 *Talanta* **15** 1397
31. Rossotti F J C and Rossotti H S 1955 *Acta Chem. Scand.* **9** 1177

32. Sigel H 1975 *Angew Chem., Int. Ed. Engl.* **14** 394
33. Dewitt R and Watters J J 1954 *J. Am. Chem. Soc.* **76** 3810
34. Martin R B and Praclos R 1975 *J. Inorg. Nucl. Chem.* **36** 1665
35. Casella L, Gulloti M and Pintar A 1988 *Inorg. Chim. Acta.* **144** 89
36. Chasteen N D 1981 *Biological magnetic resonance* (eds) L Berliner J and Rueben (New York: Plenum) vol 3, p 53
37. Holyk N H 1979 *An electron paramagnetic resonance study of model oxovanadium(IV) complexes in aqueous solution: Correlation of magnetic properties with ligand type and metal chelate structure*, M S thesis, University of New Hampshire, Durham, New Hampshire