Oxidation reactions of 1,3-diphenylpropane-1,3-dione

MEDHA RELE¹, B S PATRO², S ADHIKARI¹, G P KALENA²,
S CHATTOPADHYAY² and T MUKHERJEE^{1*}

¹Radiation Chemistry and Chemical Dynamics Division, and

²Bio-Organic Division, Bhabha Atomic Research Centre, Mumbai 400 085,
India

e-mail: mukherji@magnum.barc.ernet.in

Abstract. The free radical scavenging properties and possible antioxidant activity of 1,3-diphenylpropane-1,3-dione (**1**) are reported. Pulse radiolysis technique was employed to study the one-electron oxidation of **1** with various radicals viz. $\text{CCl}_3\text{O}_2^\bullet$, $\text{N}_3^\bullet$ and $^\bullet\text{OH}$ in homogeneous aqueous solution. All these radicals reacted with **1** under ambient conditions at almost diffusion controlled rates producing transient species with an absorption maximum around 420 nm that decayed at first order rates. The transient absorption peak was shifted in the case of $\text{CCl}_3\text{OO}^\bullet$ radical reaction with **1** due to change in the polarity of the medium. Formation of a stable product with a broad absorption band starting from 400 nm and cut off at 230 nm was observed in the oxidation of **1** with $^\bullet\text{OH}$ and $^\bullet\text{N}_3$ radicals. In a biological system also, **1** showed significant inhibitory activity against Fe^{2+} -mediated lipid peroxidation. Based on these observations, a suitable mechanism for the oxidation of **1** has been proposed.

Keywords. *b*-Diketone; pulse radiolysis; oxidation; lipid peroxidation.

1. Introduction

A wide array of phenolic substances, particularly those present in dietary and medicinal plants, has been reported to possess substantial anti-oxidative and anti-inflammatory properties, examples being curcuminoids, gingeroids etc. Numerous studies have been conducted on the anti-cancer activity of curcumin.^{1–5} Recently, we have shown that in the iron-independent peroxidation, the ginger-derived curcuminoid [6]-dehydrogingerdione possesses comparable antioxidant activity⁶ to curcumin due to its greater-membrane binding capacity.

Of late, the mechanism of antioxidant action of the curcuminoids has been the matter of considerable debate due to the possible involvement of their constituent phenolic as well as *b*-diketone groups in scavenging the oxidizing radicals. It is speculated⁷ that a considerable percentage of their antioxidant activity is due to the facile methylenic hydrogen transfer from the *b*-diketone moiety. In contrast, using classical methods, Ross *et al*⁸ have put forward a phenolic chain-breaking antioxidant mechanism for curcumin. Very recently, based on evidence from kinetic techniques like laser flash photolysis and pulse radiolysis, it has been proposed⁹ that though a methylenic radical is produced initially at the *b*-diketone moiety, the radical centre shifts its location to the phenolic site via a hydrogen transfer. In our study, the participation of both the designated functional groups in the antioxidant action of [6]-dehydrogingerdione has been established.⁶

*For correspondence

However, formation of the carbon-centred methylenic radical could not be demonstrated unequivocally due to the overlap of its absorptions with the bleaching spectrum due to depletion of the parent compound.

The present study was undertaken for further confirmation of the role of the **b**diketone group. For this, 1,3-diphenylpropane-1,3-dione (**1**) which is devoid of any phenolic group was chosen as the model **b**diketone substrate and its reactions with $\text{CCl}_3\text{O}_2^\bullet$ (a model peroxy radical), $\text{N}_3^\bullet$ and the physiologically relevant $^\bullet\text{OH}$ radical have been studied, primarily using the pulse radiolysis technique. The transients produced in these reactions were followed and the rate constants for the reactions were determined. The transients were subsequently used in elucidating the detailed mechanism for oxidation reactions. The measured rate constants were not mere radiation chemical parameters but reflected on the efficiency of **1** in scavenging free radicals and helped in understanding the ease with which the possible competing reactions occur. In addition, the protective activity of **1** against Fe^{2+} -induced lipid peroxidation of rat brain homogenate was also carried out to test its antioxidant efficacy in a biological test system.

2. Materials and methods

2.1 Chemicals

Glutathione (Aldrich, USA) and trichloroacetic acid (TCA) (Thomas Baker, India) were used as received. 2-Thiobarbituric acid (TBA), vitamin C and ferrous ammonium sulfate were procured from Himedia Lab. Pvt. Ltd., India and used as received. All other chemicals were of AR grade. The experiments were conducted in freshly de-ionised 'nanopure' water (conductivity $< 0.06 \mu\text{S cm}^{-1}$, Barnstead nano-pure cartridge filtration system). Suitable controls were used for the bioassay. Compound **1** was synthesised by condensing equimolar amounts of acetophenone and ethyl benzoate in the presence of NaOEt as the base in ether. It was characterised by IR and ^1H NMR spectra. The purity of the compound was assessed by HPLC analysis (RP-18 column) which showed a single peak.

2.2 Methods

2.2a Pulse radiolysis: The pulse radiolysis system using 7 MeV electrons has been described earlier.¹⁰ The dosimetry was carried out using an air-saturated aqueous solution containing $5 \times 10^{-2} \text{ mol dm}^{-3}$ KSCN ($G_e = 23,889 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$ per 100 eV at 500 nm ¹¹). The kinetic spectrophotometric detection system covered the wavelength range from 250 to 800 nm. The optical path length of the cell was 1.0 cm. The width of the electron pulse was 50 or 500 ns as per requirement and the dose was 16 Gy or as specified. Equimolar ($5 \times 10^{-3} \text{ mol dm}^{-3}$) HPO_4^{2-} and H_2PO_4^- were used to prepare solutions of pH 6.8. Other pH values were obtained by adding NaOH or HClO_4 . Alkaline pH was obtained by adding NaOH only. High purity ($> 99.9\%$) N_2 and/or N_2O , both from British Oxygen Company (India) Pvt. Ltd. were used as per requirement.

Irradiation of the reaction medium leads to the following reactions.

The bimolecular rate constants were calculated by plotting the pseudo-first order rates of formation of the transients against the concerned solute concentrations. The uncertainty in the measurement of bimolecular rate constants was $\pm 10\%$.

2.2b Lipid peroxidation: Rat brain homogenate, prepared from the brains of freshly killed Wistar rats, was subjected to Fe^{2+} -induced lipid peroxidation as described earlier¹² with minor modifications. Briefly, the total reaction mixture (1.0 ml) contained pH 7.4 *tris*-HCl buffer ($1.25 \times 10^{-1} \text{ mol dm}^{-3}$), brain homogenate (0.5 mg protein/ml) with or without the test compound. The reaction was triggered by the addition of ferrous ammonium sulfate ($2.0 \times 10^{-5} \text{ mol dm}^{-3}$) and vitamin C ($2.0 \times 10^{-4} \text{ mol dm}^{-3}$) followed by incubation of the mixture at 37°C for 30 min. The reaction was terminated by addition of 2 ml of TBA-TCA-HCl (0.37% TBA, 2.8% TCA, 0.25 mol dm^{-3} HCl) solution and boiling the mixture at 100°C for 10 min. The extent of lipid peroxidation was assessed from the amount of thiobarbituric acid reactive substrates (TBARS) produced, which was spectrophotometrically read at 532 nm.

3. Results and discussion

3.1 Ground state absorption study

Figure 1 shows the change in absorption of **1** in aqueous solution as a function of pH in the pH range of 2 to 12. With increasing pH of the solution the intensity of the absorption at $\sim 365 \text{ nm}$ increased while that at $\sim 320 \text{ nm}$ decreased and an isosbestic point at 340 nm was observed. From this, existence of **1** in two tautomeric (keto and enol) forms (scheme 1) at the two extreme pH levels was inferred.

Figure 1. Ground state absorption spectra of a solution of $1.0 \times 10^{-4} \text{ mol dm}^{-3}$ of **1** at different pH levels in 40% propanol-2-ol/10% acetone/50% water.

Scheme 1.

Figure 2. Transient absorption spectrum obtained from N_2O -saturated aqueous solution containing **1** ($1 \times 10^{-4} \text{ mol dm}^{-3}$) at pH 9 after the electron pulse at (a) 20 μs , (b) 50 μs , (c) 200 μs and (d) 400 μs . Insets: The absorbance (OD) vs time plot for the formation (inset A) and decay (inset B) of the radicals produced in the reaction of **1** with $\cdot\text{OH}$ radical.

3.2 Reaction with the hydroxyl radical

Hydroxyl radical was found to react with **1** to give absorption maximum at around 420 nm at 20 μs after the electron pulse with a small peak at 340 nm and a bleaching in the ground state absorption around 375 nm (figure 2). When the reaction was followed at a higher time scale, a recovery in the bleaching was also observed (not shown in the figure). The bimolecular rate constants for the formation of the peak at 420 nm at pH 9 was $1 \times 10^{10} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$. Though the formation rate constants of the other peak (at 340 nm) was similar at the very early stage, it was observed that the absorption peak at 340 nm started decaying after 6 μs , while that at 410 nm continued to build up to 20 μs . This suggested that in the initial step, $\cdot\text{OH}$ radical might possibly be forming (inset A of

figure 2) an adduct ($I_{\max}$ 340 nm) as well as a radical ($I_{\max}$ 420 nm) simultaneously. Later, a part of the adduct gets transformed into the radical. Both these events happen at the very initial stage which is followed by their decay (inset B of figure 2) almost at the same rate over a time scale of 1 ms. Based on the structure of **1**, we envisaged that the species absorbing at 420 nm would be the carbon-centered radical formed by hydrogen abstraction from the methylenic group of its *b*-diketone moiety. Similar carbon-centered radical from curcumin was reported to appear at around 490 nm.⁷ The blue shift of the observed absorption peak (420 nm) in the present case might be due to the fact that compound **1** is a relatively smaller molecule and is less conjugated. The reaction of **1** with $\cdot\text{OH}$ radical at pH 8 also showed similar absorption spectra with a negligible reduction in the rate constant. Thus, within this pH range, the site of $\cdot\text{OH}$ radical attack to the molecule appeared to remain same.

3.3 Reaction with the azide radical

The azide radical is known to react selectively with organic molecules to give the corresponding one-electron oxidized radical species. Compound **1** was found to react with this radical at pH 8 to give a spectrum showing transient absorption maximum at 410 nm with a shoulder at around 310 nm (figure 3). The bimolecular rate constant for the formation of the absorption peak at 410 nm was $1.7 \times 10^9 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$. From this, it can be inferred that the species responsible for the absorption was the same methylenic radical as observed in the reaction between **1** and the $\cdot\text{OH}$ radical ($I_{\max}$ 420 nm). It is worth mentioning that the bleaching due to depletion of the parent (inset of figure 3) at 360 nm recovered during the time period of 5 ms. One explanation of the recovery might be the most unlikely possibility of disproportionation of two methylenic radicals regenerating one molecule of **1**. For better understanding, we carried out *g*-radiolysis of **1** employing $\cdot\text{OH}$ radical as the reactant and measured the change in its absorption spectrum

Figure 3. Transient absorption spectrum obtained from N_2O -saturated aqueous solution containing **1** ($1.0 \times 10^{-4} \text{ mol dm}^{-3}$) and NaN_3 ($1.0 \times 10^{-1} \text{ mol dm}^{-3}$) at pH 8 after the electron pulse at (a) 20 μs , (b) 50 μs and (c) 200 μs . Inset: Typical oscilloscope traces for the bleach recovery at 360 nm obtained in the above reaction.

Figure 4. Steady-state absorption spectra obtained from an N_2O -saturated aqueous solution containing **1** ($5.0 \times 10^{-5} \text{ mol dm}^{-3}$) at pH 7 before and after γ -irradiation at a dose rate of 10 Gy/min (a) before irradiation, (b) after 5 min irradiation and (c) after 40 min irradiation.

(figure 4). As evident from the figure, a new product with a strong absorbance around 360 nm was formed in the reaction which contributed significantly to the apparent bleach recovery.

3.3a Oxygen effect: The carbon-centred radicals are known to react at a very fast rate with O_2 to form the corresponding peroxy radical.⁹ Thus, to confirm that the reactive species (I_{max} 410–420 nm) observed in the present study was indeed a carbon-centred radical, we examined its decay in the presence of O_2 (figure not shown). A noticeable enhancement in the rate of decay of the species was evident, thereby confirming it to be the **b**diketone methylenic radical as proposed earlier.

3.4 Reaction with trichloromethylperoxyl radical

The trichloromethylperoxyl radical ($\text{Cl}_3\text{COO}^\bullet$) has been extensively used as a representative peroxy radical for the inherent simplicity in performing the experiments and indeed, has been used earlier to study the free radical interaction between vitamins E and C¹³. The radical can be generated in aerated water/propan-2-ol/acetone (50:40:10 v/v) mixtures containing carbon tetrachloride by the following reactions.

Figure 5. Transient absorption spectra obtained from an air-saturated aqueous solution (40% propanol-2-ol, 10% acetone, 50% water) of pH 9.0, containing **1** (3×10^{-4} mol dm $^{-3}$) and CCl $_4$ (1×10^{-2} mol dm $^{-3}$), 20 (a), 50 (b) and 200 ns (c) after the electron pulse. Inset: Typical oscilloscope traces recorded at 410 nm for the radical formed in the reaction of **1** (3×10^{-4} mol dm $^{-3}$) and CCl $_3$ OO $\cdot$ radical at pH 9.0 with (2.5×10^{-5} mol dm $^{-3}$) and without ascorbic acid.

Compound **1** was found to react with CCl $_3$ O $_2\cdot$ radical at pH 9 to give transient absorptions at 450 nm (figure 5) with a bimolecular rate constant in the order of 10^9 dm 3 mol $^{-1}$ s $^{-1}$. As the absorbance at 450 nm in figure 5 is low, the error in the measurement of rate constant may be greater than mentioned in the experimental section. Hence we feel it is reasonable to present the order of magnitude of the rate constant rather than exact numerical value. The species produced is probably the same carbon-centred radical obtained in both $\cdot$ N $_3$ and $\cdot$ OH radical induced oxidation of **1**. The shift in its absorption spectrum in the present case might be caused by the change in the reaction medium. Scheme 1 represents the oxidation reactions. It must be noticed (inset of figure 5) that there was no change in the bleaching signal in the presence and absence of ascorbic acid. Hence, the radical was not repaired by ascorbic acid, in agreement with the results of our earlier study on [6]-dehydrogingerdione.⁶ It must be noticed that the bleaching recovery that was evident in reaction with $\cdot$ N $_3$ radical was not observed in this case. Possible reason is that the presence of oxygen in the case of CCl $_3$ OO $\cdot$ radical-induced oxidation changes the fate of the radical produced.

3.5 Inhibition of Fe $^{2+}$ -induced lipid peroxidation

In unstimulated control experiments, little TBARS was formed ($A_{532} = 0.06 \pm 0.01$) in the rat brain homogenate. However, addition of ferrous iron and ascorbate triggered the lipid peroxidation leading to an increase in TBARS content ($A_{532} = 0.32 \pm 0.01$). The control experiments showed that **1** itself did not interfere with the absorption at 532 nm as its addition to a terminated reaction mixture did not change the TBARS absorption at 532 nm. However, compound **1** inhibited lipid peroxidation in a concentration-dependent manner (figure 6) showing > 90% protection at a concentration of 5×10^{-4} mol dm $^{-3}$.

Figure 6. Inhibition of Fe²⁺-induced lipid peroxidation of rat brain homogenate by compound **1**.

This result clearly demonstrates that **1** can scavenge not only CCl₃O₂• radicals having high redox potentials¹⁴ but also the biologically relevant lipid peroxyl radicals very efficiently.

4. Conclusions

The present study has demonstrated that **β**-diketones such as **1** can act as effective free radical scavengers in spite of being devoid of any phenolic group. The mode of their scavenging action for the free radicals is via hydrogen abstraction from the active methylene group of **β**-diketone moiety. As revealed from the pulse radiolysis experiment, **1**, a model substrate of this class of compounds, can scavenge various oxidizing radicals efficiently. In addition, its biological efficacy as a non-phenolic antioxidant is also evident from its inhibitory activity against lipid peroxidation. Thus, a compound having a phenolic group as well as the above structural feature is anticipated to possess enhanced antioxidant activity. This may be the case with the curcuminoids.

Acknowledgements

The authors are grateful to Dr J P Mittal for his constant encouragement and support. We are also thankful to Shri V N Rao and his colleagues for valuable technical help.

References

1. Singh S V, Hu X, Srivastava S K, Singh M, Xia H, Orchard J L and Zaren H A 1998 *Carcinogenesis* **19** 1357
2. Khafif A, Schantz S P, Chou T C, Edelstein D and Sacks P G 1998 *Carcinogenesis* **19** 419
3. Reddy B S, Kawamori T, Rao C V, Lubet R A, Steele V E and Kelloff G J 1979 *Proc. Am. Assoc. Cancer Res.* **39** 126
4. Kawamori T, Lubet R, Steele V, Kelloff G J, Kaskey R B, Rao C V and Reddy B S 1999 *Cancer Res.* **59** 597

5. Samaha H S, Kelloff G J, Steele V, Rao C V and Reddy B S 1997 *Cancer Res.* **57** 1301
6. Patro B S, Rele S, Chintalwar G J, Chattopadhyay S, Adhikari S and Mukherjee T 2002 *Chem. Biochem.* **3** 364
7. Jovanovic S V, Steenken S, Boone C W and Simic M G 1999 *J. Am. Chem. Soc.* **121** 9677
8. Ross L, Berclay C, Vinqvist M R, Mukai K, Goto H, Hashimoto Y, Tokunaga A and Uno H 2000 *Org. Lett.* **2** 2841
9. Jovanovic S V, Boone C W, Steenken S, Trinoga M and Kaskey R B 2001 *J. Am. Chem. Soc.* **123** 3064
10. Mukherjee T 1997 *Atomic, molecular and cluster physics* (ed.) S A Ahmad (New Delhi: Narosa) pp. 299–316
11. Buxton G V and Stuart C R 1995 *J. Chem. Soc. Faraday Trans.* **91** 279
12. Joshi R, Adhikari S, Patro B S, Chattopadhyay S and Mukherjee T 2001 *Free Radical Biol. Med.* **30** 1390
13. Packer J E, Slater T F and Willson R L 1979 *Nature (London)* **278** 737
14. Das T N, Dhanasekaran T, Alfassi Z B and Neta P 1998 *J. Phys. Chem.* **A102** 280