Main group chemistry of 9-hydroxophenalenone: Syntheses and structural characterization of the alkaline earth and zinc complexes

ARUP MUKHERJEE^a, PRINSON P SAMUEL^b, CAROLA SCHULZKE^c and SWADHIN K MANDAL^{a,*}

^aDepartment of Chemical Sciences, Indian Institute of Science Education and Research Kolkata, Mohanpur 741 252, India

^bInstitut für Anorganische Chemie, Universität Göttingen, Tammannstrasse 4, 37077 Göttingen, Germany

^cInstitut für Biochemie, Ernst-Moritz-Arndt Universität Greifswald, Felix-Hausdorff-Straße 4, Greifswald, Germany

e-mail: swadhin.mandal@iiserkol.ac.in

MS received 13 March 2014; accepted 21 August 2014

Abstract. Herein, we report the synthesis and characterization of 9-hydroxophenalenone based alkaline earth and zinc complexes. The reaction of 9-hydroxophenalenone (HO,O-PLY (**1**)) with one equivalent of $\text{KN}(\text{SiMe}_3)_2$ and MI_2 in THF yields heteroleptic complexes $[(\text{O},\text{O}-\text{PLY})\text{M}(\text{THF})_n]\text{I}$ [$\text{M} = \text{Mg}$ (**2**), Ca (**3**), Sr (**4**), Ba (**5**); $n = 1-4$], while use of two equivalents of $\text{KN}(\text{SiMe}_3)_2$ in THF (with respect to PLY) produces homoleptic complex $(\text{O},\text{O}-\text{PLY})_2\text{Mg}(\text{THF})_2$ (**6**). Moreover, reaction between two equivalents of **1** with one equivalent of ZnMe_2 in THF produces complex $(\text{O},\text{O}-\text{PLY})_2\text{Zn}(\text{THF})_2$ (**7**). All these complexes were characterized by NMR spectroscopy and elemental analyses. The solid state structures of complexes **2**, **6** and **7** were established by single crystal X-ray diffraction analysis.

Keywords. Main group chemistry; alkaline earth complex; zinc; 9-hydroxophenalenone; X-ray crystallography.

1. Introduction

O-coordinating chelating ligands provide stabilizing scaffolds for the synthesis and isolation of highly reactive monomeric organometallic reagents.¹ Importantly, main group and transition metal complexes supported by such ancillary ligands emerged as highly efficient catalysts for the synthesis of fine chemicals.² In particular, alkaline earth and zinc based metal complexes of acetyl acetonato (acac) and tropolone (chart 1) have been extensively used in hydroamination catalysis.^{3,4} In the recent past several groups have also developed the alkaline earth metal complexes using amido- and amidophosphine-borane ligands to explore the interesting chemistry of these heavier alkaline earth metals.⁵ These ligand systems provided suitable stabilization effect to the alkaline earth metal complexes. While the metal chemistry has been developed quite exhaustively using these ancillary ligand systems,⁶ very few reports have been published dealing with closely related phenalenyl (PLY) based ligand systems.⁷ Phenalenyl is a well-known odd alternant hydrocarbon with high symmetry (D_{3h}) comprising three fused benzene rings (chart 1).⁸ Despite the phenalenyl ligand unit appearing quite promising in terms of development of metal

complexes, this ligand system has been neglected and only one early report was available in the literature dealing with the synthesis of metal complexes.^{7a} Most of the early reports considered the materials aspect of phenalenyl unit in which phenalenyl moiety acted as an organic unit capable of trapping free electron in its nonbonding orbital leading to the intriguing materials with highest room temperature conductivity among any neutral organic solids and bistable magnetic behaviour.⁹ Recently, we have started a program to design and develop metal chemistry with the PLY based ligands for different homogeneous catalytic transformations.¹⁰ Furthermore, we have recently shown that the O,O-PLY ligand coordinated to zinc ion can be used as a non-innocent building block for the construction of spin memory devices.¹¹ Herein, we report the syntheses of a series of phenalenyl ligand based heteroleptic metal complexes $(\text{O},\text{O}-\text{PLY})\text{Mg}(\text{THF})_4\text{I}$ (**2**), $(\text{O},\text{O}-\text{PLY})\text{Ca}(\text{THF})_n\text{I}$ ($n=1-4$, **3**), $(\text{O},\text{O}-\text{PLY})\text{Sr}(\text{THF})_n\text{I}$ ($n = 1-4$, **4**), $(\text{O},\text{O}-\text{PLY})\text{Ba}(\text{THF})_n\text{I}$ ($n = 1-4$, **5**) and homoleptic metal complexes $(\text{O},\text{O}-\text{PLY})_2\text{Mg}(\text{THF})_2$ (**6**), as well as $(\text{O},\text{O}-\text{PLY})_2\text{Zn}(\text{THF})_2$ (**7**). The solid state structures of complexes **2**, **6**, and **7** were established unambiguously by single crystal X-ray diffraction analysis. Although, complexes **3-5** were isolated in pure form, suitable crystals could not be isolated from the solution

*For correspondence

Acetyl acetone (ACAC)

O,O-Tropolone system

9-Hydroxophenalenone

Chart 1. Frequently used O,O-donor based ligand systems and O,O-donor based phenalenyl ligand.

despite several crystallization attempts. These complexes (**3–5**) were characterized by ^1H and ^{13}C NMR spectroscopy and elemental analysis.

2. Experimental

2.1 General

All manipulations were performed under a dry and oxygen free atmosphere (under N_2) using standard Schlenk techniques or inside a MBraun glovebox maintained at below 0.1 ppm of O_2 and H_2O level, utilizing glasswares that were oven-dried (130°C) and evacuated while hot prior to use. All solvents were distilled from Na/benzophenone prior to use. Other chemicals were purchased commercially and used as received. THF- d_8 was purchased from Cambridge Isotope Laboratories and dried by sodium potassium alloy and stored over 4 Å molecular sieves prior to use. ^1H and ^{13}C NMR spectra were recorded with a JEOL-ECS 400 MHz and Bruker AV-500 MHz spectrometer at 25°C unless otherwise stated. Elemental analyses were performed by the Analytisches Labor des Instituts für Anorganische Chemie der Universität Göttingen. Melting points were measured in sealed glass tubes with a Büchi melting point B-540 instrument. Phenalenyl ligand was synthesized according to the literature procedure.¹² Anhydrous MgI_2 , CaI_2 , SrI_2 , MgI_2 , and $\text{KN}(\text{SiMe}_3)_2$ were purchased from Sigma-Aldrich and used as received. Dimethyl zinc (1.2 M in toluene) solution was purchased from Acros Organics and used as received.

2.2 Syntheses of metal complexes

2.2a (O,O-PLY) $\text{Mg}(\text{THF})_4\text{I}$ (2**):** In a nitrogen filled glovebox, a Schlenk flask was loaded with (O,O-PLY-H) (0.196 g, 1.0 mmol), $\text{KN}(\text{SiMe}_3)_2$ (0.199 g, 1.0 mmol) and anhydrous MgI_2 (0.278 g, 1.0 mmol). The flask was sealed and removed from the glove box and attached to the vacuum line. Then 30 mL of dry THF was added to it at 25°C and the mixture was stirred at the same temperature for further 24 h. The reaction mixture was filtered through an activated celite pad and the filtrate was concentrated to approximately 10 mL under reduced

pressure and kept at -20°C for crystallization. After 1 day, suitable crystals developed from the reaction mixture. Yield: 0.31 g (74%). M.p. $255\text{--}265^\circ\text{C}$ (decomposes to black mass). ^1H NMR (THF- d_8 , 400 MHz, 25°C): δ 7.96 (2H, d, $J = 9.1$ Hz, Ar-H), 7.89 (2H, d, $J = 7.3$ Hz, Ar-H), 7.40 (1H, t, $J = 7.32$ Hz, Ar-H), 6.95 (2H, d, $J = 9.2$ Hz, Ar-H), 3.62 (m, 4H, THF), 1.77 (m, 4H, THF) ppm. ^{13}C NMR (THF- d_8 , 100 MHz, 298K): δ 182.4 (s, Ar), 140.6 (s, Ar), 132.7 (s, Ar), 130.8 (s, Ar), 129.9 (s, Ar), 126.1 (s, Ar), 122.6 (s, Ar), 115.7 (s, Ar), 68.1 (s, THF), 25.9 (s, THF) ppm. Elemental analysis: Anal. Calcd. for $\text{C}_{33}\text{H}_{47}\text{IMgO}_7$: C, 48.80%; H, 3.59%; Found: C, 48.75%; H, 3.52%. Samples for NMR spectroscopy and elemental analysis were prepared by evacuating the sample under reduced pressure (10^{-3} mbar) overnight.

2.2b (O,O-PLY) $\text{Ca}(\text{THF})_n\text{I}$ ($n = 1\text{--}4$, **3):** In a nitrogen filled glove box, a Schlenk flask was loaded with (O,O-PLY-H) (0.196 g, 1.0 mmol), $\text{KN}(\text{SiMe}_3)_2$ (0.199 g, 1.0 mmol) and anhydrous CaI_2 (0.293 g, 1.0 mmol). The flask was sealed and removed from the glove box and attached to the vacuum line. Then 30 mL of dry THF was added to it at 25°C and it was stirred at the same temperature for further 24 h. The reaction mixture was filtered through an activated celite pad and the filtrate was concentrated to approximately 10 mL under reduced pressure and kept at -20°C . After 1 day, crystals were developed from the reaction mixture, but the crystals were too thin to carry out single crystal X-ray measurements. Yield: 0.350 g (60%). M.p. $220\text{--}230^\circ\text{C}$ (decomposes to yellowish brown mass). ^1H NMR (THF- d_8 , 400 MHz, 25°C): δ 7.87 (2H, d, $J = 9.2$ Hz, Ar-H), 7.81 (2H, d, $J = 7.3$ Hz, Ar-H), 7.33 (1H, t, $J = 7.8$ Hz, Ar-H), 6.89 (2H, d, $J = 9.2$ Hz, Ar-H), 3.63 (m, 12H, THF), 1.78 (m, 12H, THF) ppm. ^{13}C NMR (THF- d_8 , 100 MHz, 298K): δ 182.4 (s, Ar), 140.3 (s, Ar), 132.6 (s, Ar), 131.9 (s, Ar), 130.9 (s, Ar), 126.6 (s, Ar), 122.5 (s, Ar), 116.9 (s, Ar), 68.5 (s, THF), 26.4 (s, THF) ppm. Elemental analysis: Anal. Calcd. for $\text{C}_{25}\text{H}_{31}\text{ICaO}_5$: C, 51.91%; H, 5.36%; Found: C, 51.53%; H, 5.73%. Samples for NMR spectroscopy and elemental analysis were prepared by evacuating the sample under reduced pressure (10^{-3} mbar) overnight.

2.2c (O,O-PLY) $\text{Sr}(\text{THF})_n\text{I}$ ($n = 1\text{--}4$, **4):** In a nitrogen filled glovebox, a Schlenk flask was loaded with (O,O-PLY-H) (0.196 g, 1.0 mmol), $\text{KN}(\text{SiMe}_3)_2$ (0.199 g, 1.0 mmol) and anhydrous SrI_2 (0.341 g, 1.0 mmol). The flask was sealed and removed from the glovebox and attached to the vacuum line. Then 30 mL of dry THF was added to it at 25°C and it was

stirred at the same temperature for further 24 h. The reaction mixture was filtered through an activated celite pad and the filtrate was concentrated to approximately 20 mL under reduced pressure and kept at -20°C . After 1 day, micro crystals were developed from the reaction mixture. Yield: 0.30 g (62%). M.p. $280-285^{\circ}\text{C}$ (decomposes to brown mass). ^1H NMR (THF-d_8 , 400 MHz, 25°C): δ 7.81 (2H, d, $J = 9.1$ Hz, Ar-H), 7.76 (2H, d, $J = 8.2$ Hz, Ar-H), 7.27 (1H, t, $J = 6.4$ Hz, Ar-H), 6.95 (2H, br d, $J = 9.1$ Hz, Ar-H), 3.63 (m, 4H, THF), 1.79 (m, 4H, THF) ppm. ^{13}C NMR (THF-d_8 , 100 MHz, 298K): δ 182.1 (s, Ar), 139.4 (s, Ar), 132.4 (s, Ar), 132.1 (s, Ar), 131.4 (s, Ar), 126.4 (s, Ar), 121.8 (s, Ar), 117.1 (s, Ar), 68.5 (s, THF), 26.4 (s, THF) ppm. Elemental analysis: Anal. Calcd. for $\text{C}_{17}\text{H}_{15}\text{ISrO}_3$: C, 42.41%; H, 3.12%; Found: C, 42.57%; H, 3.21%. Samples for NMR spectroscopy and elemental analysis were prepared by evacuating the sample under reduced pressure (10^{-3} mbar) overnight.

2.2d (*O,O*-PLY) $\text{Ba}(\text{THF})_n$ (*n* = 1-4, 5): In a nitrogen filled glove box, a Schlenk flask was loaded with (*O,O*-PLY-H) (0.196 g, 1.0 mmol), $\text{KN}(\text{SiMe}_3)_2$ (0.199 g, 1.0 mmol) and anhydrous BaI_2 (0.391 g, 1.0 mmol). The flask was sealed and removed from the glove box and attached to the vacuum line. Then 30 mL of dry THF was added to it at 25°C and it was stirred at the same temperature for further 24 h. The reaction mixture was filtered through an activated celite pad and the filtrate was concentrated to approximately 8 mL under reduced pressure and kept at 25°C . After a few days, very thin crystals were developed from the reaction mixture. Yield: 0.285 g (54%). M.p. $290-295^{\circ}\text{C}$ (decomposes to brown mass). ^1H NMR (THF-d_8 , 400 MHz, 25°C): δ 7.76 (2H, d, $J = 9.5$ Hz, Ar-H), 7.72 (2H, d, $J = 7.3$ Hz, Ar-H), 7.23 (1H, t, $J = 7.3$ Hz, Ar-H), 6.78 (2H, d, $J = 8.8$ Hz, Ar-H), 3.62 (m, 4H, THF), 1.78 (m, 4H, THF) ppm. ^{13}C NMR (THF-d_8 , 125.7 MHz, 298K): δ 180.9 (s, Ar), 138.5 (s, Ar), 132.1 (s, Ar), 131.2 (s, Ar), 130.9 (s, Ar), 125.8 (s, Ar), 120.9 (s, Ar), 116.5 (s, Ar), 68.3 (s, THF), 26.2 (s, THF) ppm. Elemental analysis: Anal. Calcd. for $\text{C}_{17}\text{H}_{15}\text{IBaO}_3$: C, 38.42%; H, 2.83%; Found: C, 38.38%; H, 2.79%. Samples for NMR spectroscopy and elemental analysis were prepared by evacuating the sample under reduced pressure (10^{-3} mbar) overnight.

2.2e (*O,O*-PLY) $_2\text{Mg}(\text{THF})_2$ (**6**): In a nitrogen filled glove box, a Schlenk flask was loaded with (*O,O*-PLY-H) (0.196 g, 1.0 mmol), $\text{KN}(\text{SiMe}_3)_2$ (0.398 g, 2.0 mmol) and anhydrous MgI_2 (0.278 g, 1.0 mmol). The flask was sealed and removed from the glovebox and attached to the vacuum line. Then 30 mL of dry THF was added

to it at 25°C and it was stirred at the same temperature for further 24 h. The reaction mixture was filtered through an activated celite pad and the filtrate was concentrated to approximately 20 mL under reduced pressure and kept at -20°C for crystallization. After 1 day, suitable crystals were developed from the reaction mixture. Yield: 0.390 g (70%). M.p. $305-310^{\circ}\text{C}$ (decomposes to brown mass). ^1H NMR (THF-d_8 , 400 MHz, 25°C): δ 7.84 (4H, d, $J = 9.1$ Hz, Ar-H), 7.78 (4H, d, $J = 7.3$ Hz, Ar-H), 7.29 (2H, t, $J = 7.32$ Hz, Ar-H), 6.93 (4H, d, $J = 9.2$ Hz, Ar-H), 3.58 (m, 8H, THF), 1.74 (m, 8H, THF) ppm. ^{13}C NMR (THF-d_8 , 125.7 MHz, 298K): δ 183.9 (s, Ar), 140.2 (s, Ar), 132.7 (s, Ar), 130.8 (s, Ar), 130.2 (s, Ar), 125.3 (s, Ar), 122.8 (s, Ar), 114.7 (s, Ar), 66.3 (s, THF), 26.2 (s, THF) ppm. Elemental analysis: Anal. Calcd. for $\text{C}_{42}\text{H}_{46}\text{MgO}_8$: C, 73.25%; H, 5.21%; Found: C, 73.53%; H, 5.73%. Samples for NMR spectroscopy and elemental analysis were prepared by evacuating the sample under reduced pressure (10^{-3} mbar) for overnight.

2.2f (*O,O*-PLY) $_2\text{Zn}(\text{THF})_2$ (**7**): Initially, (*O,O*-PLY-H) (0.392 g, 2.0 mmol) was dissolved in dry THF (15 mL) and then ZnMe_2 (1.2 M in toluene; 0.8 mL, 1.2 mmol) was added drop-wise to it at 25°C . The resulting solution was stirred at 25°C for another 4 h and then passed through an activated celite pad, concentrated to approximately 5 mL under reduced pressure and kept at -20°C . After 1 day, suitable crystals were developed from the reaction mixture. Yield: 0.420 g (80%). M.p. $220-225^{\circ}\text{C}$ (decomposes to yellowish brown mass). ^1H NMR (THF-d_8 , 400 MHz, 25°C): δ 7.96 (4H, d, $J = 9.2$ Hz, Ar-H), 7.90 (4H, d, $J = 7.3$ Hz, Ar-H), 7.42 (2H, t, $J = 8.2$ Hz, Ar-H), 7.05 (4H, d, $J = 9.1$ Hz, Ar-H), 3.58 (4H, br, THF), 1.73 (4H, br, THF) ppm. ^{13}C NMR (THF-d_8 , 100 MHz, 298 K): δ 183.9 (s, Ar), 140.3 (s, Ar), 132.7 (s, Ar), 130.7 (s, Ar), 130.2 (s, Ar), 126.3 (s, Ar), 122.7 (s, Ar), 114.6 (s, Ar), 68.1 (s, THF), 25.9 (s, THF) ppm. Elemental analysis: Anal. Calcd. for $\text{C}_{42}\text{H}_{46}\text{ZnO}_8$: C, 68.32%; H, 4.18%; Found: C, 68.39%; H, 4.15%. Samples for NMR spectroscopy and elemental analysis were prepared by evacuating the sample under reduced pressure (10^{-3} mbar) for overnight.

2.3 X-ray crystallographic details

Suitable crystals of **2** and **6** were mounted on a glass fibre, and data were collected on an IPDS II Stoe image-plate diffractometer (graphite monochromated Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å) at 133(2) K (table 2). The data was integrated with X-area. While, the data of **7** was collected from a shock-cooled crystal at 100(2) K

on a Bruker SMART-APEX II diffractometer with a D8 goniometer equipped with a fine focus INCOATEC Mo-microsource (table 2).¹³ The structures were solved by Direct Methods (SHELXS-97)¹⁴ and refined by full-matrix least-square methods against F^2 (SHELXL-97). All non-hydrogen atoms were refined with anisotropic displacement parameters. The hydrogen atoms were refined isotropically on calculated positions using a riding model. No restraints were used. CCDC 941715 (**2**), 941716 (**6**), and 941717 (**7**), respectively contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

3. Results and Discussion

The syntheses of complexes **2-7** were accomplished by reacting **1** with the suitable anhydrous metal precursors MI_2 , $M = Mg, Ca, Sr, Ba$ in the presence of $KN(SiMe_3)_2$ and $ZnMe_2$ at 25°C (scheme 1). Complexes **2-5** were synthesized by reacting the phenalenyl ligand, 9-hydroxyphenalenone in a 1:1:1 ratio with $KN(SiMe_3)_2$ and alkaline earth metal iodides (Mg, Ca, Sr and Ba) in THF at 25°C (scheme 1). In contrast, when the reaction was carried in 1:2:1 ratio of 9-hydroxyphenalenone, $KN(SiMe_3)_2$ and MgI_2 , it resulted in the formation of the homoleptic complex $(O,O-PLY)_2Mg(THF)_2$ (**6**) (scheme 1). Complex **7** was synthesized by adding a $ZnMe_2$ solution in toluene drop-by-drop to a solution of **1** in THF in 1:2 stoichiometric ratio at 25°C over a period of 4h. The 1H NMR spectra of all the reaction mixtures reveal clean and

nearly quantitative conversions of the reactants to products as exhibited by the absence of any characteristic O-H resonance at δ 18.1 ppm in C_6D_6 . Complexes **2-7** were characterized by 1H and ^{13}C NMR spectroscopy and elemental analyses. The solid state structures of complexes **2, 6** and **7** were determined by single crystal X-ray diffraction technique. X-ray quality crystals of **2, 6** and **7** were grown from their saturated THF solutions at $-20^\circ C$. All the phenalenyl based metal complexes were synthesized in reasonable to good isolated yield (54–80%). The colour of complex **2** is deep yellow, while that of **3** is pale yellow, the colour of **4** is yellowish orange and that of **5** is light yellow. The 1H NMR spectra of the phenalenyl unit of complexes **2-5** are listed in table 1.

The 1H NMR spectra of **2** shifted to most downfield, while that of **5** was shifted most upfield (table 1). This can be attributed to the fact that on going down to a group from Mg to Ba , the effective nuclear charge (charge/radius) on the metal centre decreases, as a result the charge transfer from attached phenalenyl unit to the metal centre decreases from **2** to **5** resulting a most downfield shift of the phenalenyl unit in case of **2**.

3.1 X-ray crystallographic studies

X-ray quality crystals of **2, 6**, and **7** were grown from suitable solvents at specific temperatures and the structural refinement parameters of these compounds are summarized in table 2. Selected bond lengths ($\text{\AA}$) and angles ($^\circ$) of these compounds are given in tables 3–4 and structures are depicted in figures 1–3.

Scheme 1. Syntheses of phenalenyl ligand based metal complexes. Reaction conditions: (a) **1** $KN(SiMe_3)_2$, MI_2 (1:1:1) in the THF at 25°C; (b) **1**, $KN(SiMe_3)_2$, MgI_2 (1:2:1) in THF at 25°C; and (c) **1**, $ZnMe_2$ (2:1) in THF at 25°C. Uncoordinated THF molecules found during the crystallization have been omitted for clarity.

Table 1. ^1H NMR spectra of the phenalenyl unit observed in complexes **2-5**.

Complex	^1H NMR spectra of the phenalenyl unit in THF- d_8 (δppm)
2	7.96 (d, 2H), 7.89 (2H, d), 7.40 (1H, t), 6.95 (2H, d)
3	7.87 (d, 2H), 7.81 (2H, d), 7.33 (1H, t), 6.89 (2H, d)
4	7.81 (d, 2H), 7.76 (2H, d), 7.27 (1H, t), 6.95 (2H, d)
5	7.76 (d, 2H), 7.72 (2H, d), 7.23 (1H, t), 6.78 (2H, d)

Suitable crystals of **2** were obtained from THF solution at -20°C and the molecular structure of **2** was determined unambiguously by single crystal X-ray diffraction analysis (figure 1). Complex **2** crystallizes in the monoclinic space group $P2_1/n$ with one molecule in the asymmetric unit. The X-ray structure of **2** reveals a distorted octahedral geometry around the magnesium centre with two oxygen atoms of the phenalenyl ligand and four oxygen atoms from coordinated THF molecules. Inspection of figure 1 reveals that there is an iodide ion present, which does not show a direct bond with the magnesium, rather it stays as counter anion. The $\text{Mg(1)-O}_{\text{PLY}}$ bond distances of **2** [Mg(1)-O(1) , 2.014(3) Å; and Mg(1)-O(2) , 2.011(3) Å] (table 3) are comparable with the one found in the complex $\{\text{Mg}[\text{OC}(\text{Mes})]_2[\text{Mes} = 2,4,6\text{-Me}_3(\text{C}_6\text{H}_2)]\}$ (2.042 Å),¹⁵ but longer than the Mg-O_L bond distances found in the complex $\text{L}(\text{Me})\text{Al-O-Mg}(\text{THF})_2\text{-O-Al}(\text{Me})\text{L}$ [$\text{L} = \text{CH}\{(\text{CMe})(2,6-$

$i\text{Pr}_2\text{C}_6\text{H}_3\text{N})\}_2$] (1.865 Å).¹⁶ The bond angle of O(2)-Mg(1)-O(4) is nearly linear [$175.53(12)^\circ$] and it is wider than the other bond angles $\text{O}_{\text{THF}}\text{-Mg-O}_{\text{THF}}$ observed in **2** [O(5)-Mg(1)-O(6) $169.12(13)^\circ$, O(5)-Mg(1)-O(3) $172.58(13)^\circ$] (table 3). The axial sites around the magnesium centre of **2** are occupied by two more oxygen atoms of coordinated THF molecules exhibiting a O(6)-Mg(1)-O(5) bond angle of $169.12(13)^\circ$ (table 3), which is deviated from the ideal octahedral angle.

Suitable crystals of **6** were obtained from THF solution at -20°C and the molecular structure of **6** was determined by single crystal X-ray diffraction analysis (figure 2). Complex **6** crystallizes in the monoclinic space group $P2_1/n$ with one molecule in the asymmetric unit (figure 2). The X-ray structure of **6** reveals a distorted octahedral geometry around the magnesium centre with four oxygen atoms of the phenalenyl ligands and two oxygen atoms from coordinated THF

Table 2. Crystal and structure refinement parameters for phenalenyl based metal complexes **2**, **6** and **7**.

	2	6	7
CCDC no.	941715	941716	941717
Fw	706.92	703.10	744.16
Cryst size/mm	$0.19 \times 0.16 \times 0.11$	$0.45 \times 0.20 \times 0.13$	$0.12 \times 0.09 \times 0.06$
Cryst syst	Monoclinic	Monoclinic	Monoclinic
Space group	$P2_1/n$	$P2_1/n$	$P2_1/n$
$a/\text{\AA}$	16.510(5)	9.4844(19)	9.4716(2)
$b/\text{\AA}$	10.078(5)	15.956(3)	16.0292(4)
$c/\text{\AA}$	20.055(5)	11.784(2)	11.7172(3)
β/deg	90.219(5)	96.08(3)	96.1280(10)
$V/\text{\AA}^3$	3337(2)	1773.3(6)	1768.77(7)
$D_{\text{calcd}}/\text{g cm}^{-3}$	1.407	1.317	1.282
Z	4	2	2
Abs coeff/ mm^{-1}	1.023	0.106	1.397
θ range/deg	1.60–27.06	2.16–27.05	2.16–28.28
Reflns collected/indep reflns	20462 / 7241	17450 / 3852	33694 / 4393
Max. and min. Transmn.	0.9232 and 0.8565	0.8796 and 0.7970	0.9606 and 0.8919
Final R indices [$I > 2\sigma(I)$]	$R1 = 0.0547$, $wR2 = 0.0972$	$R1 = 0.0509$, $wR2 = 0.1104$	$R1 = 0.0368$, $wR2 = 0.0765$
R indices (all data)	$R1 = 0.0884$, $wR2 = 0.1067$	$R1 = 0.0902$, $wR2 = 0.1201$	$R1 = 0.0608$, $wR2 = 0.0857$
Largest diff peak and hole/ $e \text{\AA}^{-3}$	1.154 and -0.551	0.435 and -0.303	0.396 and -0.406

Figure 1. Molecular structure of complex **2**. Thermal ellipsoids are drawn with 50% probability. Hydrogen atoms and uncoordinated THF molecule have been omitted for the sake of clarity.

molecules. Due to symmetry, the Mg(1)–O(1) and Mg(1)–O(1A) bond distances of the two phenalenyl ligands in **6** are the same [2.0127(14) Å] as that of Mg(1)–O(2) and Mg(1)–O(2A) bond distances [2.0121(15) Å] (table 4). The bond distances of Mg(1)–O_{PLY} found in **6** are comparable with the one found in **2** (2.011 Å). The bond distances of the Mg–O_{THF} [Mg(1)–O(3) 2.1431(14) and Mg(1)–O(3A) 2.1431(14)] found in **6** are comparatively longer than those found in **2** (average Mg–O_{THF} bond distance 2.10 Å) (table 4). The bond angles of O(3)–Mg(1)–O(3A), O(2)–Mg(1)–O(2A) and O(1)–Mg(1)–O(1A) observed in **6** are all 180° based on the crystal's symmetry. The O_{PLY}–Mg–O_{PLY} bond angle found in **6** [O(1)–Mg(1)–O(2) and O(1A)–Mg(1)–O(2A) 85.99(6)°] is marginally larger than the one found in **2** [O(2)–Mg(1)–O(1) 85.71(12)°] (table 4).

Complex **7** crystallizes in the monoclinic space group $P2_1/n$ with one molecule in the asymmetric unit

(figure 3). The X-ray structure of **7** reveals a distorted octahedral geometry around the zinc centre. The zinc atom is six-coordinated by the two bi-dentate chelating phenalenyl (O,O-PLY) ligands, and two molecules of coordinated THF. The (O,O-PLY) ligand is almost symmetrically attached to the metal centre resulting in equal Zn–O_{PLY} bond lengths of Zn(1)–O(1) 2.0185(12) Å and Zn(1)–O(2) 2.0137(13) Å (table 4). These Zn–O_{PLY} bond distances found in **7** are slightly longer than the Zn–O_{PLY} bond distances found in [(O,O-PLY)ZnEt(THF)] [Zn(2)–O(1) 1.95(15) Å and Zn(2)–O(2) 1.97(15) Å].^{10d} Moreover, the Zn–O_{THF} bond distance [Zn(1)–O(3) 2.2075(13) Å] found in **7** is longer than the Zn–O_{PLY} bond distances (2.02 Å) found in the same complex (table 4). The O(1)–Zn(1)–O(2) angle [92.00(5)°] falls within the range of the bond angle observed [91.78(6)°] in previously reported complex [(O,O-PLY)ZnEt(THF)] (table 4).^{10d}

Table 3. Selected bond distances (Å) and angles (°) for **2**.

Bond	distances (Å)	Bonds	angles (°)
Mg(1)–O(1)	2.014(3)	O(2)–Mg(1)–O(1)	85.71(12)
Mg(1)–O(2)	2.011(3)	O(2)–Mg(1)–O(5)	95.15(12)
Mg(1)–O(3)	2.106(3)	O(2)–Mg(1)–O(6)	94.35(12)
Mg(1)–O(4)	2.097(3)	O(2)–Mg(1)–O(3)	87.90(12)
Mg(1)–O(5)	2.094(3)	O(2)–Mg(1)–O(4)	175.53(12)
Mg(1)–O(6)	2.104(3)	O(1)–Mg(1)–O(3)	172.58(13)
C(1)–O(1)	1.274(5)	O(1)–Mg(1)–O(6)	91.23(12)
C(11)–O(2)	1.278(4)	O(1)–Mg(1)–O(4)	89.86(11)
C(1)–C(2)	1.446(5)	O(1)–Mg(1)–O(5)	94.76(13)
C(10)–C(11)	1.439(5)	O(5)–Mg(1)–O(6)	169.12(13)
C(1)–C(12)	1.449(5)	O(5)–Mg(1)–O(3)	89.55(12)
C(11)–C(12)	1.439(5)	O(4)–Mg(1)–O(6)	85.19(12)

Figure 2. Molecular structure of complex **6**. Thermal ellipsoids are drawn with 50% probability. Hydrogen atoms and uncoordinated THF molecule have been omitted for the sake of clarity.

Table 4. Selected bond distances (Å) and angles (°) for **6** and **7**.

Bond	distances (Å), 6 (M = Mg)	distances (Å), 7 (M = Zn)
M(1)-O(1)	2.0127(14)	2.0185(12)
M(1)-O(1A)	2.0127(14)	2.0185(12)
M(1)-O(2)	2.0121(15)	2.0137(13)
M(1)-O(2A)	2.0121(15)	2.0138(13)
M(1)-O(3)	2.1431(14)	2.2075(13)
M(1)-O(3A)	2.1431(14)	2.2075(13)
Bond	angles (°)	angles (°)
O(3)-M(1)-O(3A)	180.00(7)	179.999(1)
O(2)-M(1)-O(2A)	180.00(9)	180.0
O(1)-M(1)-O(1A)	180.00(7)	180.0
O(1)-M(1)-O(2)	85.99(6)	92.00(5)
O(1A)-M(1)-O(2A)	85.99(6)	92.00(5)
O(1)-M(1)-O(3)	90.31(6)	89.51(5)
O(1)-M(1)-O(3A)	89.69(6)	90.49(5)
O(2)-M(1)-O(3)	89.55(6)	89.09(5)
O(2)-M(1)-O(3A)	90.45(6)	90.91(5)

Figure 3. Molecular structure of complex **7**. Thermal ellipsoids are drawn with 50% probability. Hydrogen atoms and uncoordinated THF molecule have been omitted for the sake of clarity.

4. Conclusions

We have successfully synthesized and characterized the phenalenyl ligand based alkaline earth metal complexes as well as the zinc complexes. The reaction of the phenalenyl ligand with one equivalent of $\text{KN}(\text{SiMe}_3)_2$ and anhydrous alkaline earth metal iodides in THF yielded heteroleptic complexes, while use of two equivalents of $\text{KN}(\text{SiMe}_3)_2$ in THF (with respect to PLY) yielded homoleptic complex. Moreover, reaction between two equivalents of the phenalenyl ligand with one equivalent of ZnMe_2 in THF provided homoleptic zinc complex. The solid state structures of two magnesium and a zinc based complexes were established by single crystal X-ray diffraction analysis. All the complexes have been characterized by NMR-spectroscopy and elemental analysis technique.

Supplementary Information

Supplementary information contains the NMR characterization data of all the complexes synthesized herein, CIF files and check CIF files of complexes **2**, **6** and **7**, and atomic coordinates of these complexes determined by X-ray crystallography. Electronic supplementary information can be seen at www.ias.ac.in/chemsci.

Acknowledgements

A.M. is thankful to IISER-Kolkata for a research fellowship. S.K.M. thanks CSIR (Sanction no. 01 (2369)/10/EMR-II), India for financial support.

References

- (a) Edelmann F T 2009 *Chem. Soc. Rev.* **38** 2253; (b) Edelmann F T, Freckmann D M M and Schumann H 2002 *Chem. Rev.* **102** 1851; (c) Gade L H 2002 *Acc. Chem. Res.* **35** 575; (d) Kempe R 2000 *Angew. Chem. Int. Ed.* **39** 468
- (a) Zeimentz P M, Arndt S, Elvidge B R and Okuda J 2006 *Chem. Rev.* **106** 2404; (b) Park S J, Han Y Y, Kim S K, Lee J S, Kim H K and Do Y K 2004 *J. Organomet. Chem.* **689** 4263; (c) Gibson V C and Spitzmesser S K 2003 *Chem. Rev.* **103** 283
- (a) Crimmin M R, Casely I J and Hill M S 2005 *J. Am. Chem. Soc.* **127** 2042; (b) Crimmin M R, Arrowsmith M, Barrett A G M, Casely I J, Hill M S and Procopiou P A 2009 *J. Am. Chem. Soc.* **131** 9670; (c) Barrett A G M, Brinkmann C, Crimmin M R, Hill M S, Hunt P and Procopiou P A 2009 *J. Am. Chem. Soc.* **131** 12906; (d) Datta S and Roesky P W 2007 *Organometallics* **26** 4392
- (a) Mukherjee A, Nembenna S, Sen T K, Sarish S P, Ghorai P K, Ott H, Stalke D, Mandal S K and Roesky H W 2011 *Angew. Chem. Int. Ed.* **50** 3968; (b) Zhang X, Emge T J and Hultsch K C 2012 *Angew. Chem. Int. Ed.* **51** 394; (c) Whitehorne T J J and Schaper F 2012 *Chem. Commun.* **48** 10334; (d) Mukherjee A, Sen T K, Mandal S K, Maity B and Koley D 2013 *RSC Adv.* **3** 1255
- (a) Yang D, Ding Y, Wu H and Zheng W 2011 *Inorg. Chem.* **50** 7698; (b) Sarazin Y, Roşca D, Poirier V, Roisnel T, Silvestru A, Maron L and Carpentier J 2010 *Organometallics* **29** 6569; (c) Kottalanka R K, Anga S, Naktode K, Laskar P, Nayek H P and Panda T K 2013 *Organometallics* **32** 4473
- (a) Bourget-Merle L, Lappert M F and Severn J R 2002 *Chem. Rev.* **102** 3031; (b) Roesky P W 2000 *Chem. Soc. Rev.* **29** 335
- (a) Demura Y, Kawato T, Kanatomi H and Murase I 1975 *Bull. Chem. Soc. Jpn.* **48** 2820; (b) Reid D H 1965 *Quart. Rev.* **19** 274; (c) Das A, Scherer T M, Mobin S M, Kaim W and Lahiri G K 2012 *Inorg. Chem.* **51** 4390; (d) Das A, Scherer T M, Mondal P, Mobin S M, Kaim W and Lahiri G K 2012 *Chem. Eur. J.* **18** 14434; (e) Dey S K, Honecker A, Mitra P, Mandal S K and Mukherjee A 2012 *Eur. J. Inorg. Chem.* 5814
- Morita Y, Suzuki S, Sato K and Takui T 2011 *Nat. Chem.* **3** 197
- (a) Mandal S K, Samanta S, Itkis M E, Jensen D W, Reed R W, Oakley R T, Tham F S, Donnadiou B and Haddon R C 2006 *J. Am. Chem. Soc.* **128** 1982; (b) Pal S K, Itkis M E, Tham F S, Reed R W, Oakley R T and Haddon R C 2005 *Science* **309** 281; (c) Mandal S K, Itkis M E, Chi X, Samanta S, Lidsky D, Reed R W, Oakley R T, Tham F S and Haddon R C 2005 *J. Am. Chem. Soc.* **127** 8185; (d) Itkis M E, Chi X, Cordes A W and Haddon R C 2002 *Science* **296** 1443
- (a) Mukherjee A, Sen T K, Ghorai P K, Samuel P P, Schulzke C and Mandal S K 2012 *Chem. Eur. J.* **18** 10530; (b) Mukherjee A, Sen T K, Mandal S K, Kratzert D, Stalke D, Döring A and Schulzke C 2011 *J. Chem. Sci.* **123** 139; (c) Sen T K, Mukherjee A, Modak A, Ghorai P K, Kratzert D, Granitzka M, Stalke D and Mandal S K 2012 *Chem. Eur. J.* **18** 54; (d) Sen T K, Mukherjee A, Modak A, Mandal S K and Koley D 2013 *Dalton Trans.* **42** 1893; (e) Mukherjee A, Sen T K, Ghorai P K and Mandal S K 2013 *Sci. Rep.* **3** 2821; (f) Mukherjee A, Sen T K, Ghorai P K and Mandal S K 2013 *Organometallics* **32** 7213
- Raman K V, Kamerbeek A M, Mukherjee A, Atodiresei N, Sen T K, Lazić P, Caciuc V, Michel R, Stalke D, Mandal S K, Blügel S, Münzenberg M and Moodera J S 2013 *Nature* **493** 509
- Haddon R C, Rayford R and Hirani A M 1981 *J. Org. Chem.* **46** 4587
- Schulz T, Meindl K, Leusser D, Stern D, Graf J, Michaelsen C, Ruf M, Sheldrick G M and Stalke D 2009 *J. Appl. Crystallogr.* **42** 885
- Sheldrick G M 2008 *Acta Crystallogr. Sect. A* **64** 112
- Zhang D and Kawaguchi H 2006 *Organometallics* **25** 5506
- Nembenna S, Roesky H W, Mandal S K, Oswald R B, Pal A, Herbst-Irmer R, Noltemeyer M and Schmidt H-G 2006 *J. Am. Chem. Soc.* **128** 13056