



REGULAR ARTICLE

Sulfated polyborate as an eco-compatible solid acid catalyst for efficient and facile solvent-free synthesis of polyhydroquinolines

DILIP AUTE^{a,b}, AKSHAY KSHIRSAGAR^a, BHAGWAT UPHADE^a and ANIL GADHAVE^{a,*}

^aDepartment of Chemistry and Research Centre, Padmashri Vikhe Patil College (Affiliated to Savitribai Phule Pune University, Pune), Pravaranagar District, Ahmednagar, Maharashtra 413 713, India

^bDepartment of Chemistry, Arts Commerce & Science College, Satral, Maharashtra, India

E-mail: anilgadhave@gmail.com

MS received 16 April 2020; revised 29 August 2020; accepted 2 September 2020

Abstract. An environmentally viable, green and efficient protocol for the synthesis of polyhydroquinolines using sulfated polyborate catalyst have been developed by Hantzsch four-component condensation between aromatic aldehydes, dimedone, ammonium acetate and ethylacetoacetate or ethylcyanoacetate at 100 °C under solvent-free condition. The sulfated polyborate catalyst is characterized by FT-IR, XRD, SEM and EDAX techniques. The noticeable features of this methodology are mild Brønsted acidic nature of the catalyst, good to excellent product yields (85–94%), short reaction time (18–30 min), solvent-free condition, operational simplicity and applicability to a wide range of substrates (29 examples).

Keywords. Sulfated polyborate; polyhydroquinolines; Hantzsch condensation; solvent-free; solid acid catalyst.

1. Introduction

The solid acid catalyst plays an important role to develop economically and environmentally sustainable processes.¹ Boric acid is a readily available inexpensive mild catalyst used in a variety of organic transformations.² It is known that above 100 °C it undergoes polymerization and liberates water molecule which may hamper the rate of reactions.³ This led to the development of sulfated polyborate with enhanced catalytic activity.⁴ Over the past years, the sulfated polyborate has emerged as a potential solid acid catalyst used in several MCRs with excellent yields, short reaction time and operational simplicity.^{5–7} The interesting feature of this catalyst is, it possesses Lewis acidic property due to polymeric borate backbone and Brønsted acidic nature due to support of -SO₃H group which makes it more applicable.

Polyhydroquinolines (PHQs) are privileged scaffolds due to the presence of 1,4-dihydropyridine (DHP) moiety which is a core part of calcium channel functioning as well as hypertension preventive drugs

viz., nimodipine, lacidipine, felodipine, amlodipine, nicardipine, nifedipine, etc.⁸ The polyhydroquinolines show diverse biological functions such as antimalarial, bronchodilatory, vasodilator, antidiabetic and antiasthmatic activities.⁹ In 1881, synthesis of substituted 1,4-dihydropyridines was first reported by Arthur Hantzsch and further Safak's group synthesized 1,4-DHP-annulated novel heterocycle known as polyhydroquinoline (Figure 1).¹⁰

The straightforward synthesis of polyhydroquinoline involves one-pot four-component condensation reaction between aromatic aldehydes, dimedone, ammonium acetate and ethylacetoacetate or ethylcyanoacetate using a variety of catalytic systems. Among them synthesis of polyhydroquinolines using ethylacetoacetate involves spinel FeAl₂O₄,¹¹ Cu(II)-DCC-CMK-3,¹² AFGO nanosheets,¹³ TEDETA@BNPs,¹⁴ MCM-41@serine@Cu(II),¹⁵ ascorbic acid,¹⁶ Fe₃O₄@SiO₂/ZnCl₂,¹⁷ SBA-15@Glycine-M,¹⁸ PMOICSPrSO₃H,¹⁹ KH₂PO₄,²⁰ [PyridineSO₃H]Cl,²¹ CdS nanowires,²² Fe₃O₄/SiO₂-OSO₃H,²³ ZnO[DABCO(C₂COOH)₂]²⁺[Br]₂,²⁴ nano VTiO₂,²⁵ H₃BW₁₂O₄₀²⁶ and nicotinic acid.²⁷ The condensation reaction using ethylcyanoacetate utilizes

*For correspondence

Electronic supplementary material: The online version of this article (<https://doi.org/10.1007/s12039-020-01850-w>) contains supplementary material, which is available to authorized users.




Figure 1. Some bioactive molecules containing 1,4-dihydropyridine nucleus.

systems such as MCM-41,²⁸ Pd-Schiff-Base@Fe₃O₄ MNPs,²⁹ ZnO-nanoparticle,³⁰ FSM-16-SO₃H nanoparticles,³¹ microwave irradiation,³² ClO₄/Zr-MCM-41,³³ nano ionic liquid 1-methylimidazolium trinitromethanide,³⁴ Pd(0) nanoparticles,³⁵ Fe₃O₄@FSM-16-SO₃H³⁶ and Fe₃O₄@B-MCM-41.³⁷

Despite the merits of reported methods, they may also suffer from one or more shortcomings like harsh conditions, expensive catalyst/reagents, tedious workup, use of toxic catalyst/reagents, limited scope, lengthy reaction time and poor product yields. Hence, there is a need for a general, efficient, feasible and environment-friendly protocol for the synthesis of polyhydroquinolines. With this view and in continuation to our research work^{38–42} herein, we wish to explore a catalytic application of sulfated polyborate for efficient and facile one-pot four-component Hantzsch condensation between differently substituted aromatic aldehydes, dimedone, ethylacetoacetate/ethylcyanoacetate and ammonium acetate at 100 °C under the solvent-free condition to give polyhydroquinolines with excellent yields in a short time.

2. Experimental

2.1 General information

The solvents and reagents used in this work were of AR grade and used without further purification. Fourier transform infrared (FT-IR) spectrum was recorded on Perkin-Elmer, 400 FT-IR spectrometer. The X-ray diffraction pattern was obtained on Panalyticals X Pert Pro, X-Ray Diffractometer. The SEM images were obtained from JSM6100, Jeol Scanning Electron Microscope. The ¹H NMR spectrum was recorded on Bruker Avance 500MHz NMR spectrometer. Mass spectrum was recorded on Waters Q-ToF Micromass spectrometer. The TLC (silica G60 F254 plates, Merck) was used to monitor the progress of the reaction. The melting points were recorded in an open capillary tube and are uncorrected.

2.2 Preparation of sulfated polyborate acid catalyst

The readily available and inexpensive boric acid and chlorosulfonic acid were used for the preparation of sulfated polyborate following the literature reported method.⁴ The catalyst obtained was solid and crystalline in nature. It was characterized by FT-IR, XRD, SEM and EDAX techniques.

2.3 General procedure for the synthesis of substituted polyhydroquinolines


A round bottom flask equipped with water condenser was charged with a mixture of substituted aromatic aldehydes (**1**, 1 mmol), dimedone (**2**, 1 mmol), ammonium acetate (**3**, 1.5 mmol) and ethyl acetoacetate (**4**, 1 mmol) or ethylcyanoacetate (**5**, 1 mmol). To this, 80 mg of sulfated polyborate catalyst was added and the resulting reaction mixture was magnetically stirred in oil-bath at 100–110 °C under solvent-free conditions till completion of the reaction (Table 3). The progress of the reaction was monitored by TLC using ethyl acetate: n-hexane (3:7) as an eluent. The hot ethanol was added to the reaction mixture and filtered to remove insoluble material. The solid appeared in filtrate was filtered to furnish polyhydroquinolines (**6a-p** and **7a-m**). The physical and spectral data of synthesized compounds were compared with the data reported in the literature which supports the formation of polyhydroquinolines (Scheme 1).

3. Results and Discussion

3.1 Characterization of aluminized polyborate

The catalyst was synthesized by the following procedure reported in the literature.⁴ After preparation of sulfated polyborate its formation was confirmed by FT-IR, XRD, SEM and EDAX techniques.

3.1a FT-IR analysis: The FT-IR spectra show characteristic broad band between 3100–3600 cm⁻¹ is due to O-H stretching of -SO₃H and B-O-H groups.



Scheme 1. Synthesis of polyhydroquinolines using aldehydes (1 mmol), dimedone (1 mmol), ethylacetoacetate or ethylcyanoacetate (1 mmol) and ammonium acetate (1.5 mmol) catalysed by sulfated polyborate.




Figure 2. XRD pattern of sulfated polyborate.

The broad band between $1400\text{--}1650\text{ cm}^{-1}$ includes peaks due to O–H bending of $-\text{SO}_3\text{H}$ group and B–O stretching. The peaks at 1192 cm^{-1} , 1057 cm^{-1} , 1000 cm^{-1} and 546 cm^{-1} corresponds to O=S=O asymmetric stretching, O=S=O symmetric stretching, S=O stretching and O–B–O stretching of catalyst, respectively (Supplementary Information).^{43, 44}

3.1b XRD analysis: The formation of sulfated polyborate was confirmed by its powder X-ray

diffraction pattern which shows a significant peak at $2\theta = 28.11^\circ$ is due to B–O bonds in the crystalline structure of the catalyst (Figure 2).⁴⁵

3.1c SEM analysis: The SEM image shows that catalyst is crystalline in nature with particles of different shapes and sizes (Figure 3).

3.1d EDAX analysis: The energy dispersive spectrum shows the presence of oxygen/sulfur/boron




Figure 3. SEM image of sulfated polyborate.





Figure 4. EDAX analysis of sulfated polyborate.

with a ratio of 81.82: 2.47: 15.58 wt % over different areas which indicate the formation of sulfated polyborate (Figure 4).

3.2 Study of catalytic activity of sulfated polyborate in the synthesis of polyhydroquinolines

Inspired by catalytic applications of sulfated polyborate reported in the literature⁴⁻⁷ we explored its

catalytic utility in the multicomponent synthesis of polyhydroquinolines. In order to set the best reaction conditions, we choose a model reaction of benzaldehyde (**1a**, 1 mmol), dimedone (**2**, 1 mmol), ammonium acetate (**3**, 1.5 mmol) and ethylacetoacetate (**5**, 1 mmol) or ethylcyanoacetate (**6**, 1 mmol). We first performed a series of trial experiments to optimize the influence of different solvents and temperature for the synthesis of polyhydroquinolines (**6a/7a**) with 80 mg of fixed amount of sulfated polyborate catalyst. The


Table 1. Optimization of solvent and temperature for the synthesis of polyhydroquinolines^a.

Entry	Solvent	Temperature (°C)	Time (min)	Yield ^b (%)	Yield ^c (%)
1	H ₂ O	r.t.	60	–	–
2	EtOH	r.t.	60	–	–
3	H ₂ O:EtOH	r.t.	60	–	–
4	H ₂ O	reflux	60	37	41
5	EtOH	reflux	60	85	79
6	CH ₃ CN	reflux	60	60	56
7	THF	reflux	60	45	47
8	DMF	80 °C	60	52	51
9	solvent-free	r.t.	60	45	41
10	solvent-free	80 °C	30	85	80
11	solvent-free	100 °C	20	93	92
12	solvent-free	110 °C	20	92	92

^aReaction conditions: benzaldehyde (1 mmol), dimedone (1 mmol), ammonium acetate (1.5 mmol) and ethylcyanoacetate (1 mmol) using sulfated polyborate catalyst (80 mg).

^bIsolated yield of **6a**.

^cIsolated yield of **7a**.


Table 2. Optimization of the amount of catalyst.

Entry	Catalyst amount (mg)	Time (min)	Yield ^b (%)	Yield ^c (%)
1	none	30	trace	trace
2	20	30	48	40
3	40	30	70	66
4	60	25	82	83
5	80	20	93	92
6	100	20	91	90

^aReaction conditions: **1** (1 mmol), **2** (1 mmol), **4** or **5** (1 mmol) and **3** (1.5 mmol) at 100–110 °C under solvent-free condition.

^bIsolated yield of **6a**.

^cIsolated yield of **7a**.

Table 3. Synthesis of polyhydroquinoline derivatives.^a


Entry	Substituents ('R')	'X'	Product	Time (min)	Yield ^b (%)	M. P. (°C)		Ref.
						Observed	Reported	
1	-H	-CH ₃	6a	20	93	217–219	218–220	[14]
2	2-Cl	-CH ₃	6b	28	85	206–207	207–209	[22]
3	2-NO ₂	-CH ₃	6c	23	89	206–208	206–207	[12]
4	3-NO ₂	-CH ₃	6d	25	92	183–184	182–184	[21]
5	3-Cl	-CH ₃	6e	27	90	204–206	205–207	[21]
6	3-OMe	-CH ₃	6f	30	88	199–201	199–201	[21]
7	4-OMe	-CH ₃	6g	25	92	258–260	257–260	[22]
8	4-Me	-CH ₃	6h	26	91	260–262	259–262	[22]
9	4-Br	-CH ₃	6i	20	90	254–256	256–257	[22]
10	4-NO ₂	-CH ₃	6j	18	94	240–242	240–242	[22]
11	4-Cl	-CH ₃	6k	20	93	244–246	244–246	[22]
12	4-OH	-CH ₃	6l	30	86	231–233	232–234	[22]
13	4-F	-CH ₃	6m	24	91	186–187	184–186	[14]
14	3,4 (OMe) ₂	-CH ₃	6n	30	87	196–197	196–198	[15]
15	2-Thiophene	-CH ₃	6o	26	89	240–241	240–241	[22]
16	2-Furan	-CH ₃	6p	28	85	247–249	248–250	[22]
17	-H	-NH ₂	7a	20	92	158–160	158–161	[33]
18	2-Cl	-NH ₂	7b	25	90	180–182	182–183	[33]
19	2-OH	-NH ₂	7c	28	85	113–114	114–116	[37]
20	3-OMe	-NH ₂	7d	30	88	157–159	155–158	[33]
21	3-NO ₂	-NH ₂	7e	20	90	182–183	183–184	[36]
22	3-Br	-NH ₂	7f	20	89	136–137	135–136	[36]
23	3-OH	-NH ₂	7g	25	91	166–167	165–168	[36]
24	4-Me	-NH ₂	7h	20	92	158–162	158–161	[33]
25	4-F	-NH ₂	7i	20	91	156–157	155–158	[36]
26	4-NO ₂	-NH ₂	7j	19	93	188–190	189–190	[33]
27	4-OMe	-NH ₂	7k	20	92	134–136	134–136	[33]
28	4-Cl	-NH ₂	7l	20	93	161–162	160–162	[33]
29	4-(NMe) ₂	-NH ₂	7m	25	86	122–125	123–124	[36]

^aSynthesized under optimized conditions.^bIsolated yield.


model reaction was performed in solvents like H₂O, EtOH, H₂O:EtOH (1:1), DMF, THF and CH₃CN as well as a solvent-free condition at different temperature (Table 1). From these experiments, a reaction under the solvent-free condition at 100–110 °C is found to be the most suitable for the synthesis of polyhydroquinolines (**6a/7a**) with excellent yields in a short time.

With this optimized condition, our next aim was to find the role and amount of sulfated polyborate

catalyst towards the synthesis of polyhydroquinolines (**6a/7a**). The model reaction was performed with different amount of catalyst and results obtained are summarized in Table 2. The outcome of this study indicates that reaction does not occur in absence of the catalyst. The increase in the amount of catalyst in order of 20 mg, 40 mg, 60 mg and 80 mg increases the yield of the product and decreases the time required for conversion into the product. Further increase in the amount of catalyst has no significant effect on the

Table 4. The comparison of obtained results with literature reported methods for the synthesis of polyhydroquinolines.

Entry	Catalyst and solvent	Catalyst amount	Temp. (°C)	Time (min)	Yield (%)	Ref.
1	spinel FeAl ₂ O ₄ , EtOH	40 mg	80 °C	30–80	96	[11]
2	Cu(II)-DCC-CMK-3, solvent-free	30 mg	80 °C	20–90	98	[12]
3	AFGO nanosheets, EtOH	25 mg	r.t.	120–240	87–91	[13]
4	MCM-41@serine@Cu(II), EtOH	5 mg	80 °C	170–215	96	[15]
5	Ascorbic acid, solvent-free	5% mol	80 °C	90–300	70–99	[16]
6	Fe ₃ O ₄ @SiO ₂ /ZnCl ₂ , solvent-free	5 mg	110 °C	30	92	[17]
7	SBA-15@Glycine-M, EtOH	80 mg	60 °C	110–220	92	[18]
8	Sulfated polyborate, solvent-free	80 mg	100 °C	20	93	Present work




Entry	Catalyst and solvent	Catalyst amount	Temp. (°C)	Time (min)	Yield (%)	Ref.
1	MCM-41-OBF ₂ , EtOH	50 mg	reflux	15–110	75–95	[28]
2	Pd-Schiff-Base@Fe ₃ O ₄ MNPs, solvent-free	3 mg	70 °C	15	99	[29]
3	ZnO-NPs, solvent-free	10 mol%	r.t.	20–25	91	[30]
4	FSM-16-SO ₃ H NPs, EtOH	40 mg	reflux	5–60	80–94	[31]
5	MWI/EtOH	–	MWI	5–7	89–92	[32]
6	ClO ₄ ⁻ /Zr-MCM-41, EtOH	10 mg	reflux	10–60	80–94	[33]
7	Fe ₃ O ₄ @FSM-16-SO ₃ H, EtOH	40 mg	reflux	10–50	85–96	[36]
8	Fe ₃ O ₄ @B-MCM-41, EtOH	50 mg	reflux	15–130	75–90	[37]
9	Sulfated polyborate, solvent-free	80 mg	100 °C	20	92	Present work

yield and rate of reaction. From all these investigations, we report that optimal results were obtained with 80 mg of sulfated polyborate catalyst under the solvent-free condition at 100–110 °C for the synthesis of polyhydroquinolines (**6a/7a**).

To explore the generality, effectiveness, versatility and scope of the present protocol, we carried out reactions between dimedone, ammonium acetate, ethylacetoacetate or ethylcyanoacetate with

aromatic aldehydes having –OMe, –Me, –Cl, –Br, –F, –NO₂ at the ortho, meta or para positions under the optimized conditions (Table 3). It is worth noting that all reactions with aldehydes having electron-donating as well electron-withdrawing groups occurred with no side reaction with good to excellent yields (85–94%) within a short time (18–30 min). This method is also suitable for heteroaromatic aldehydes such as furan-2-carbaldehyde (**60**)



Scheme 2. Plausible mechanism for the synthesis of polyhydroquinolines.

and thiophene-2-carbaldehyde (**6p**). The aldehydes having ortho substituents also give products with good yields (**6b**, **6c**, **7b**, **7c**).

To see the merits of the present protocol, we compared results obtained with literature reported methods for the synthesis of polyhydroquinolines (**6/7**). The data summarized (Table 4) indicates that sulfated polyborate shows excellent catalytic efficacy for the

synthesis of **6** and **7** under environmentally benign conditions.

3.3 Plausible mechanism

A reaction mechanism was proposed for the four-component synthesis of polyhydroquinolines as

depicted in Scheme 2. The electrophilicity of carbonyl group of aldehyde, dimedone and ethylacetoacetate or ethylcyanoacetate is activated due to the acidic nature of sulfated polyborate catalyst. In presence of a catalyst, the dimedone reacts with ammonia to give enamine. The subsequent Knoevenagel condensation of aromatic aldehydes with ethylacetoacetate or ethylcyanoacetate gives intermediate. The enamine and the intermediate undergo Michael addition followed by sequential cyclization and dehydration reaction gives polyhydroquinolines (6/7).

4. Conclusions

This study reports an application of sulfated polyborate as a mild solid acid catalyst in a cost-effective, efficient and facile protocol for the synthesis of polyhydroquinolines using aromatic aldehydes, dimedone, ammonium acetate and ethylacetoacetate or ethylcyanoacetate at 100-110 °C under solvent-free condition. The key features of this protocol are, use of inexpensive and non-toxic sulfated polyborate catalyst, eco-sustainable and green approach, short reaction times, good to excellent product yields, synthetic simplicity, safe and clean reaction profile and simple purification of the product.

Supplementary Information (SI)

Spectral data is available at www.ias.ac.in/chemsci.

Acknowledgements

The authors are thankful to the Management and Principal (P.V.P. College, Pravaranagar & A.C.S. College, Satral) for providing all the necessary facilities. We express our gratitude towards Principal Dr. B. K. Karale for his constant encouragement. We express our sincere thanks to SAIF, Punjab University (Chandigarh) for providing all necessary spectral analysis facility.

References

1. Tanemura K and Suzuki T 2013 Aniline-terephthalaldehyde resin p-toluenesulfonic acid (ATRT) salt as efficient mild polymeric solid acid catalyst *Tetrahedron Lett.* **54** 6740
2. Pal R 2018 Boric acid in organic synthesis: scope and recent developments *Arkivoc* **1** 346
3. Chandrasekhar S and Gopalaiah K 2002 Beckmann rearrangement of ketoximes on solid metaboric acid: a simple and effective procedure *Tetrahedron Lett.* **43** 2455
4. Khatri C K, Rekunge D S and Chaturbhuj G U 2016 Sulfated polyborate: a new and eco-friendly catalyst for one-pot multi-component synthesis of 3,4-dihydropyrimidin-2(1H)-ones/thiones via Biginelli reaction *New J. Chem.* **40** 10412
5. Patil M S, Palav A V, Khatri C K and Chaturbhuj G U 2017 Rapid, efficient and solvent-free synthesis of (un)symmetrical xanthenes catalyzed by recyclable sulfated polyborate *Tetrahedron Lett.* **58** 2859
6. Rekunge D S, Khatri C K and Chaturbhuj G U 2017 Sulfated polyborate-catalyzed efficient and expeditious synthesis of (un)symmetrical ureas and benzimidazolones *Tetrahedron Lett.* **58** 4304
7. Jejurkar V P, Khatri C K, Chaturbhuj G U and Saha S 2017 Environmentally benign, highly efficient and expeditious solvent-free synthesis of trisubstituted methanes catalyzed by sulfated polyborate *ChemistrySelect* **2** 11693
8. Safak C and Simsek R 2006 Fused 1, 4-dihydropyridines as potential calcium modulatory compounds *Mini-Rev. Med. Chem.* **6** 747
9. Godfraind T R, Miller R and Wibo M 1986 Calcium antagonism and calcium entry blockade *Pharmacol. Rev.* **38** 321
10. Hantzsch A 1881 Condensationsprodukte aus aldehydammoniak und ketonartigen Verbindungen *Ber. Dtsch. Chem. Ges.* **14** 1637
11. Ghorbani-Choghamarani A, Ghadermazi M, Shiri L and Taherinia Z 2019 Synthesis and characterization of spinel FeAl₂O₄ (hercynite) magnetic nanoparticles and their application in multicomponent reactions *Res. Chem. Intermed.* **45** 5705
12. Ghafouri-Nejad R and Hajjami M 2020 Synthesis and characterization of Cu–N, N'-dicyclohexylcarbodiimide supported on CMK-3 as a novel, efficient and recoverable nanocatalyst for synthesis of tetrazole, polyhydroquinoline, and sulfoxide derivatives *React. Kinet. Mech. Cat.* **129** 371
13. Choudhury P, Ghosh P and Basu B 2020 Amine-functionalized graphene oxide nanosheets (AFGONs): an efficient bifunctional catalyst for selective formation of 1,4-dihydropyridines, acridinediones and polyhydroquinolines *Mol. Divers.* **24** 283
14. Ghorbani-Choghamarani A, Heidarneshad Z, Tahmasbi B and Azadi G 2018 TEDETA@BNPs as a basic and metal free nanocatalyst for Knoevenagel condensation and Hantzsch reaction *J. Iran. Chem. Soc.* **15** 2281
15. Tamoradi T, Ghadermazi M and Ghorbani-Choghamarani A 2018 Synthesis of polyhydroquinoline, 2, 3-dihydroquinazolin-4(1H)-one, sulfide and sulfoxide derivatives catalyzed by new copper complex supported on MCM-41 *Catal. Lett.* **148** 857
16. Sheout I, Boulcina R, Boudjemaa B, Boumoud T and Debache A, Solvent-free Synthesis of polyhydroquinoline and 1, 8-dioxodecahydroacridine derivatives via the Hantzsch reaction catalyzed by a natural organic acid: a green method *Synth. Commun.* **47** 1185
17. Maleki B, Alinezhad H, Atharifar H, Tayebee R and Mofrad A V 2019 One-pot synthesis of polyhydroquinolines catalyzed by ZnCl₂ supported on nano Fe₃O₄@SiO₂ *Org. Prep. Proced. Int.* **51** 301
18. Tamoradi T, Ghorbani-Choghamarani A, Ghadermazi M and Veisi H 2019 SBA-15@Glycine-M (M= Ni and Cu): two green, novel and efficient catalysts for the one-pot synthesis of 5-substituted tetrazole and polyhydroquinoline derivative *Solid. State Sci.* **91** 96

19. Yaghoubi A, Dekamin M G and Karimi B 2017 Propylsulfonic acid-anchored isocyanurate-based periodic mesoporous organosilica (PMO-ICS-PrSO₃H): A highly efficient and recoverable nanoporous catalyst for the one-pot synthesis of substituted polyhydroquinolines *Catal. Lett.* **147** 2656
20. Yü S, Wu S, Zhao X and Lu C 2017 Green and efficient synthesis of acridine-1,8-diones and hexahydroquinolines via a KH₂PO₄ catalyzed Hantzsch-type reaction in aqueous ethanol *Res. Chem. Intermed.* **43** 3121
21. Sakram B, Sonyanaik B, Ashok K and Rambabu S 2016 Polyhydroquinolines: 1-sulfonypyridinium chloride catalyzed an efficient one-pot multicomponent synthesis via Hantzsch condensation under solvent-free conditions *Res. Chem. Intermed.* **42** 7651
22. Harale R R, Shitre P V, Sathe B R and Shingare M S 2017 Visible light motivated synthesis of polyhydroquinoline derivatives using CdS nanowires *Res. Chem. Intermed.* **43** 3237
23. Maleki A, Akbarzade A R and Bhat A R 2017 Green synthesis of polyhydroquinolines via MCR using Fe₃O₄/SiO₂-OSO₃H nanostructure catalyst and prediction of their pharmacological and biological activities by pass *J. Nanostruct. Chem.* **7** 309
24. Mondal P, Chatterjee S, Sarkar P, Bhaumik A and Mukhopadhyay C 2019 Preparation of DABCO-based acidic-ionic-liquid-supported ZnO nanoparticles and their application for ecofriendly synthesis of *N*-aryl polyhydroquinoline derivatives *ChemistrySelect* **4** 11701
25. Dharma Rao G B, Nagakalyan S and Prasad G K 2017 Solvent-free synthesis of polyhydroquinoline derivatives employing mesoporous vanadium ion doped titania nanoparticles as a robust heterogeneous catalyst via the Hantzsch reaction *RSC Adv.* **7** 3611
26. Momeni T, Heravi M M, Hosseinejad T, Mirzaei M and Zadsirjan V 2020 H₅BW₁₂O₄₀-catalyzed syntheses of 1, 4-dihydropyridines and polyhydroquinolines via Hantzsch reaction: joint experimental and computational studies *J. Mol. Struct.* **1199** 127011
27. Davarpanah J, Ghahremani M and Najafi O 2019 Synthesis of 1, 4-dihydropyridine and polyhydroquinoline derivatives via Hantzsch reaction using nicotinic acid as a green and reusable catalyst *J. Mol. Struct.* **1177** 525
28. Abdollahi-Alibeik M and Rezaeipoor-Anari A 2015 MCM-41 functionalized with B-F bond: preparation, characterization and catalytic application in the synthesis of polyhydroquinolines *Lett. Org. Chem.* **12** 651
29. Karimi Zarchi M A and Darbandizadeh Mohammad Abadi S S A 2019 Facile and efficient protocols for C-C and C-N bond formation reactions using a superparamagnetic palladium complex as reusable catalyst *Res. Chem. Intermed.* **45** 2605
30. Kassaei M Z, Masroui H and Movahedi F 2010 ZnO-nanoparticle-promoted synthesis of polyhydroquinoline derivatives via multicomponent Hantzsch reaction *Monatsh. Chem.* **141** 317
31. Hashemi-Uderji S, Abdollahi-Alibeik M and Ranjbar-Karimi R 2018 FSM-16-SO₃H nanoparticles as a novel heterogeneous catalyst: preparation, characterization, and catalytic application in the synthesis of polyhydroquinolines *Main. Group. Met. Chem.* **41** 91
32. Saha M, Luireingam T S, Merry T and Pal A K 2013 Catalyst-free, Knoevenagel-Michael addition reaction of dimedone under microwave irradiation: An efficient one-pot synthesis of polyhydroquinoline derivatives *J. Heterocycl. Chem.* **50** 941
33. Abdollahi-Alibeik M and Hoseinikhah S S 2016 ClO₄⁻/Zr-MCM-41 Nanoparticles prepared at mild conditions: a novel solid acid catalyst for the synthesis of polyhydroquinolines *J. Iran. Chem. Soc.* **13** 1339
34. Zolfigol M A, Bagheri S, Moosavi-Zare A R, Alinezhad H, Norouzi M and Vahdat S M 2014 Synthesis of the first nano ionic liquid 1-methylimidazolium trinitromethanide {[HMIM]C(NO₂)₃} and its catalytic use for Hantzsch four-component condensation *RSC Adv.* **4** 57662
35. Saha M and Pal A K 2011 Palladium (0) nanoparticles: an efficient catalyst for the one-pot synthesis of polyhydroquinolines *Tetrahedron Lett.* **52** 4872
36. Hashemi-Uderji S, Abdollahi-Alibeik M and Ranjbar-Karimi R 2019 Fe₃O₄@FSM-16-SO₃H as a novel magnetically recoverable nanostructured catalyst: preparation, characterization and catalytic application *J. Porous Mater.* **26** 467
37. Abdollahi-Alibeik M and Rezaeipoor-Anari A 2016 Fe₃O₄@B-MCM-41: a new magnetically recoverable nanostructured catalyst for the synthesis of polyhydroquinolines *J. Magn. Magn. Mater.* **398** 205
38. Borhade A V, Uphade B K and Gadhave A G 2016 Calcined eggshell: an environmentally benign green catalyst for synthesis of 2-arylbenzothiazole derivatives *Res. Chem. Intermed.* **42** 6301
39. Gadhave A G, Gaikar R B, Kuchekar S R and Karale B K 2014 Synthesis and antimicrobial activity of some novel [4-(1,2,3-thiadiazol-4-yl) phenoxy] methylene anchored 1,3,4-triazoles and 1,3,4-thiadiazoles *J. Heterocycl. Chem.* **51** 1849
40. Borhade A V, Uphade B K and Gadhave A G 2015 Efficient, solvent free synthesis of acridinediones catalysed by CdO nanoparticles *Res. Chem. Intermed.* **41** 1447
41. Aute D S, Kshirsagar A S, Uphade B K and Gadhave A G 2020 Aluminized polyborate-catalysed green and efficient synthesis of polyhydroquinolines under solvent-free conditions *Res. Chem. Intermed.* **46** 3491
42. Aute D S, Kshirsagar A S, Uphade B K and Gadhave A G 2020 Ultrasound assisted and aluminized polyborate prompted green and efficient one pot protocol for the synthesis of hexahydroquinolines *Polycycl. Aromat. Comp.* <https://doi.org/10.1080/10406638.2020.1781206>
43. Yeo J S 2013 Enhancement of the rate capability of graphite via the introduction of boron-oxygen functional groups *Int. J. Electrochem. Sci.* **8** 1308
44. Siqueira R L, Yoshida I V P, Pardini L C and Schiavon M A 2007 Poly(borosiloxanes) as precursors for carbon fiber ceramic matrix composites *Mat. Res.* **10** 147
45. Khalafi-Nezhad A, Foroughi H O, Doroodmand M M and Panahi F 2011 Silica boron-sulfuric acid nanoparticles (SBSANs): preparation, characterization and their catalytic application in the Ritter reaction for the synthesis of amide derivatives *J. Mater. Chem.* **21** 12842