



REGULAR ARTICLE

Reactivity of ynamides with AlCl_3 and ICl : Ready access to (*E*)- α -chloroenamides and (*E/Z*)- α -chloro- β -iodo-enamides

ALLA SIVA REDDY, MANDALA ANITHA, SURAJ and K C KUMARA SWAMY*

School of Chemistry, University of Hyderabad, Hyderabad 500 046, Telangana, India

E-mail: kckssc@uohyd.ac.in; kckssc@yahoo.com

MS received 10 November 2020; accepted 29 November 2020

Abstract. AlCl_3 acts as a chlorinating agent for ynamides in the presence of stoichiometric amount of water in the environmentally benign solvent dimethylcarbonate, affording efficient access to (*E*)- α -chloroenamides via hydrochlorination, with water as a protic source. The role of water in the reaction was proven by deuterium labelling experiment. Epoxy-ynamides undergo iodochlorination in addition to the cleavage of the epoxy ring to afford (*E/Z*)- α -chloro- β -iodo-enamides. Regio- and stereochemical assignments for the products are based on X-ray crystallographic studies.

Keywords. Ynamide; hydrochlorination; iodochlorination; enamide; epoxy-ynamide; stereoselective; X-ray structure.

1. Introduction

Ynamides are reactive alkynes and hence are useful precursors that can be utilized in different organic transformations due to the presence of the triple bond directly attached to the nitrogen atom.¹ The divergence in the reactivity or the regio- and stereo-selective transformations of these precursors when compared to simple alkynes, is presumably due to the formation of a reactive keteniminium ionic species (Scheme 1).²


Ynamides are well-explored in several organic transformations, including those involved in the formation of carbocycles, heterocycles and enamides.^{3–5} Especially during the last decade, several protocols for the synthesis of enamides from ynamide substrates have been developed, illustrating the versatility of these precursors.^{5–9} Significantly, the reports mainly included halogenation,⁷ reaction with carbonyl compounds⁸ and addition of phosphite/alkyne/hydrostannane.⁹ However the α -halogenation of ynamides is a simple approach to prepare enamides; the products can be readily derivatized further.^{7,10–13} Hsung and co-workers accomplished stereoselective access to enamides by the hydrohalogenation of ynamides using the magnesium halides in wet DCM (Scheme 2a).^{10a}

Another report from Matsuo *et al.*, involved the titanium tetrachloride mediated addition of carbonyl compounds to terminal ynamides (Scheme 2b).^{10b} Sahoo and coworkers developed a triphenylphosphine promoted stereoselective hydrohalogenation of ynamides using carbon tetrahalides as a halogen source (Scheme 2c).^{10c} The report from Zhu *et al.*, involved the chloro-allylation of ynamides under palladium catalysis (Scheme 2d).^{10d} The most recent paper by Tang *et al.*, utilized aq. hydrohalic acids in DCM (Scheme 2e).^{10e} Thus there is a significant interest in these reactions- in particular with respect to regio- and stereoselectivity. Additionally, it should be noted that enamides themselves are versatile building blocks in organic synthesis.^{11,12}


Initially, our idea was to use AlCl_3 which may readily activate the alkyne because of its strong Lewis acidic character.¹³ It should be noted that even though AlCl_3 has been used in some transformations of ynamides either in catalytic or stoichiometric amounts, they did not include the involvement of chloride ion transfer from the AlCl_3 .¹⁴ To our knowledge, the use of AlCl_3 as a chlorinating agent, in particular for ynamides, has not been explored. Consequently, our interest to develop new synthetic strategies by using

*For correspondence

Electronic supplementary material: The online version of this article (<https://doi.org/10.1007/s12039-020-01880-4>) contains supplementary material, which is available to authorized users.



Scheme 1. Special Reactivity of Ynamide Precursors.



Scheme 2. Reports on α -chlorination of ynamides.

ynamides¹⁵ prompted us to explore this aspect. Herein, we report an elegant approach to (*E*)- α -chloroenamides by the chlorination of ynamides using AlCl_3 as a chlorinating agent in a regio- and stereoselective manner; we have used dimethyl carbonate, a non-halogenated solvent in the present work.

In addition to the above, iodochlorination of ynamides using the inexpensive ICl could lead to functionalized enamides, with regiospecific iodination/chlorination. Iodochlorination of only alkynyl esters has been reported earlier by Ogilvie *et al.*, although iodobromination is known¹⁶ This aspect has not been explored with ynamides so far (although the products could be useful synthons) and hence we briefly dwell upon this reaction also in this paper. Use of epoxyynamides led to epoxide ring-opening in addition to iodochlorination.

2. Experimental

General experimental set up with the details of equipment used, including X-ray structural analysis are given in the Supplementary Information. The known precursors used in the present study are prepared by methods reported in the literature.^{15,17}

(i) Synthesis of enamide compounds **2a-2w**

Representative procedure for the synthesis of compound 2a. To an oven-dried Schlenk tube was added 4-*N*-dimethyl-*N*-phenylethynyl-benzenesulfonamide **1a** (0.20 mmol), AlCl_3 (0.22 mmol), dimethyl carbonate (1 mL) and H_2O (0.20 mmol). The tube was sealed under a nitrogen atmosphere and heated at 80 °C (oil bath temperature) overnight. After completion of the reaction as monitored by TLC, the crude mixture was cooled to room temperature. The mixture was then passed through celite, washed with ethyl acetate (20 mL) and concentrated in vacuum. The residue was then purified by using silica gel column chromatography using hexane-ethyl acetate (9:1) as the eluent to afford (*E*)-*N*,4-dimethyl-*N*-styrylbenzenesulfonamide **2a**. Compounds **2b-w** were prepared following the same procedure and the same molar quantities.

(*E*)-*N*-(1-chloro-2-phenylvinyl)-*N*,4-dimethylbenzenesulfonamide (2a**):** White solid. Yield 0.059 g (92%); M.p. 104–106 °C; $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.80 (d, $J = 8.5$ Hz, 2H), 7.63–7.61 (m, 2H), 7.40–7.36 (m, 2H), 7.35–7.32 (m, 3H), 6.68 (s, 1H), 3.06 (s, 3H), 2.46 (s, 3H); $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ 144.5, 134.2, 133.1, 132.4, 129.8, 129.5, 128.9, 128.8₁, 128.7₈, 128.7, 35.7, 21.6; IR (KBr) 3060, 2926, 1634, 1593, 1443, 1361, 1164, 1092, 963, 808, 756, 689 cm^{-1} ; HRMS (ESI): Calcd. for $\text{C}_{16}\text{H}_{16}\text{ClNO}_2\text{SNH}_4$ ($\text{M}^+ + \text{NH}_4$): m/z 339.0934. Found: 339.0935.

(ii) Synthesis of compounds **6a-c**

General procedure for 6a. To an oven dried 10 mL RBF, 4-methyl-*N*-(oxiran-2-ylmethyl)-*N*-(phenylethynyl)benzenesulfonamide (**5a**; 0.100 g, 0.3 mmol), 1M ICl in acetonitrile (1 mL) solution was added at 0 °C. The mixture was stirred at 0 °C to rt (25 °C) for 1 h. After completion of the reaction as monitored by TLC, the solvent was removed under reduced pressure and purification by column chromatography (hexane/ethyl acetate 9:1) afforded compound **6a**. Compounds **6b-6c** were prepared following the same procedure and by using the same molar quantities.

(*E*)-*N*-(3-chloro-2-hydroxypropyl)-*N*-(1-chloro-2-iodo-2-phenylvinyl)-4-methylbenzenesulfonamide (6a**):** Yield 0.143 g (89% with *E*:*Z* in 53:47 ($^1\text{H NMR}$), M.p. 124–126 °C; IR (KBr): 3534, 2926, 1595, 1490, 1441, 1353, 1167, 1090, 1025, 816, 701, 658 cm^{-1} . $^1\text{H NMR}$ (400 MHz, CDCl_3): Isomer 1: δ 7.89–7.86 (m, 4H), 7.41–7.40 (m, 5H), 4.22–4.18 (m, 1H), 3.76–3.73 (m, 1H), 3.69–3.66 (m, 1H), 3.59–3.54 (m, 1H), 3.44–3.40 (m, 1H), 2.47 (s, 3H). Isomer 2:

7.39–7.32 (m, 9H), 4.29–4.25 (m, 1H), 4.05–4.02 (m, 1H), 3.85–3.81 (m, 1H), 3.66–3.62 (m, 1H), 3.37–3.33 (m, 1H), 2.47 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): Isomer 1: δ 145.3, 141.3, 134.3, 129.9, 129.2, 129.0, 128.5, 128.5, 128.4, 128.2, 103.6, 68.6, 52.2, 46.1, 21.7. Isomer 2: 145.2, 141.1, 134.3, 129.8, 129.1, 128.9, 128.5, 128.4, 127.7, 102.9, 69.3, 51.3, 47.9, 21.7. HRMS (ESI): Calcd. for $\text{C}_{18}\text{H}_{19}^{35,37}\text{Cl}_2\text{INO}_3\text{S}$ ($\text{M}^+ + \text{H}$): m/z 525.9507, 527.9477, 529.9447. Found: 525.9506, 527.9475, 529.9429. This compound was crystallized from DCM/ethyl acetate (2:1) mixture at 25 °C. X-ray structure was determined for *E*-isomer.

Full characterization data for the products **2a–2w** (compounds **2j**, **2k** and **2w** are known^{7j, 10e}) and **6a–6c** are given in the Supplementary Information.

3. Results and Discussion


Initially, we performed the reaction of *N*-alkynyl benzenesulfonamide **1a** with AlCl_3 (0.4 equiv) using H_2O (2 equiv) as a proton source in DMF solvent at 80 °C for 12 h (Table 1, entry 1). Delightfully, the α -chloroamide **2a** was isolated in 36% yield in addition to the water addition by-product **3** in 52% yield. To further improve the yield and stereoselectivity of the desired product **2a** (Scheme 3), we then moved to the optimization of the reaction conditions (Table 1).

The reaction using 1 equiv. of water also led to the formation of both the products **2a** and **3**. However, the use of stoichiometric amounts of AlCl_3 (1.1 equiv.) furnished the desired product **2a** in good yield. It is noteworthy that, no over-chlorination was observed even with an excess of AlCl_3 (3 equiv.). Variation of solvents from DMF to THF and PEG-400 improved the stereoselectivity. Unexpectedly, the reaction was not clean in the ethanol solvent. Both the yield and stereoselectivity decreased when we performed the reaction in toluene. Interestingly, the product was obtained in good yield and stereoselectivity by using acetonitrile as the solvent. Further, to our surprise, the reaction proceeded cleanly by the use of dimethyl carbonate as a solvent affording the product **2a** in 92% isolated yield in a regio- and stereo-selective manner. We noticed that the reaction also proceeded at the room temperature (25 °C) but with a slower reaction rate, reduction in the yield of the product with some unreacted starting material. More importantly, no product formation was observed in the absence of AlCl_3 . We also checked other Lewis acids. While TiCl_4 gave a mixture of products, SnCl_4 led predominantly to the water addition product **3** with only traces of the desired product. In fact, $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ is also suitable as a chlorinating agent, though with decreased reaction rate. Hence the optimized condition for the regio- and stereo-specific hydro-chlorination of yn-amides was **1a** (0.20 mmol), AlCl_3 (0.22 mmol) in

Table 1. Optimization study for the synthesis of (*E*)-*N*-(1-chloro-2-phenylvinyl)-*N*,4-dimethylbenzenesulfonamide **2a** from ynamide **1a**^a.

Entry	AlCl_3 (equiv.)	Solvent	Yield (%) ^b 2a (<i>E</i> : <i>Z</i>)
1 ^{c,d}	AlCl_3 (0.4)	DMF	36 (94:6)
2 ^d	AlCl_3 (0.4)	DMF	42
3	AlCl_3 (1.1)	DMF	85
4	AlCl_3 (3)	DMF	86
5	AlCl_3 (1.1)	THF	90 (96:4)
6 ^d	AlCl_3 (1.1)	PEG-400	82 (99:1)
7	AlCl_3 (1.1)	EtOH	traces
8	AlCl_3 (1.1)	toluene	76 (92:8)
9	AlCl_3 (1.1)	CH_3CN	92 (96:4)
10	AlCl_3 (1.1)	$(\text{MeO})_2\text{CO}$	92 ('99:traces)
11 ^c	AlCl_3 (1.1)	$(\text{MeO})_2\text{CO}$	46
12	–	$(\text{MeO})_2\text{CO}$	No reaction
13	TiCl_4 (1.1)	$(\text{MeO})_2\text{CO}$	traces
14 ^d	SnCl_4 (1.1)	$(\text{MeO})_2\text{CO}$	traces

^aYnamide (0.20 mmol), AlCl_3 (x equiv.), solvent (1 mL) and H_2O (1 equiv.) at 80 °C for 12 h. ^bYield of the isolated product and stereoisomeric ratio in parenthesis was based on ^1H NMR. ^c H_2O (2 equiv.) used. ^d**3** was observed. ^eperformed at rt (25 °C).



Scheme 3. Reaction of ynamide **1a** with AlCl_3 affording products **2a** and **3**.

dimethyl carbonate (1 mL) and H_2O (0.20 mmol) at 80 °C for 12 h [Table 1, entry 10].

After having the above-optimized reaction conditions in hand, we then sought to explore the substrate scope for this chloride ion transfer from AlCl_3 to ynamides. As illustrated in Scheme 4, various types of ynamides are transformed into the corresponding (*E*)-enamides in good to excellent yields in a regio- and stereo-selective manner using AlCl_3 as the hydrochlorinating source. Precursors with substituents on the aryl moiety attached to sulfonyl group afforded products **2a–c** in excellent yields. Ynamides with bulky, bicyclic or heterocyclic moiety also offered the corresponding products **2d–i** in good yields. Ynamides with different substituents on the nitrogen atom of ynamides too produced the enamides **2j–l**. The reaction proceeded well with cyclic ynamides to provide the related products **2m–n**. In addition, this transformation worked well with precursor having an aliphatic substituent on the alkyne functionality, delivering the compound **2o** in good yield and selectivity. Functionalities such as halo, nitro and free sulfonamide functionalities on the ynamides were compatible with this reaction and produced the enamides **2p–t** in good yields. Interestingly, bis-hydrochlorination could be effected to yield the enamide **2u** in good yield by using 2.2 equiv of AlCl_3 . Furthermore, the present method could be extended by varying the electron-withdrawing groups. Thus replacing the sulfonyl group with the phosphoryl group or carbonyl group did not affect the regio- and stereo-selectivity, and afforded the corresponding products **2v–w** in excellent yields. The structures of compounds **2l**, **2q** and **2w** were confirmed by X-ray crystallographic analysis (Figure 1 and Figure S3, Supplementary Information). X-ray structures of these compounds showed strong hydrogen bonding between the chlorine and the alkenyl hydrogen atom which may account for the observed stereo-specificity.


To elucidate the probable pathway for this chloride ion transfer from AlCl_3 to ynamides, we have done the following control experiments. Thus the reaction of ynamide **1a** with AlCl_3 (1.1 equiv) and D_2O (1 equiv) in dimethyl carbonate solvent delivers the compound **2a'**

with about 76% deuteration at one of the olefinic positions (Scheme 5a). The fact that the isolation of mono-deuterated α -chloroenamide **2a'** suggested the crucial role of water as a proton source in the reaction. No reaction occurred when we treated *N*-propargylated sulfonamide **4** under similar reaction conditions due to the nonexistence of reactive keteniminium ion (Scheme 5b).

It is known that AlCl_3 reacts vigorously with water (excess) to liberate HCl. Thus in the above reaction, it is possible that hydrolysis of AlCl_3 takes place leading to the liberation of HCl which then reacts with the ynamide. However, it can be noted that $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ is commercially available and hence it is difficult to say whether the reaction takes place with the liberated HCl or not. We have not done further experiments since the idea was to get the chloroenamides. Also, as pointed above, for reactions of ynamides in which AlCl_3 is used as a catalyst,^{14a} traces of moisture could lead to hydrochlorination and one has to be careful.

3.1 ICl addition reactions of epoxy ynamides

In principle, the interhalogen compound $\text{I}^{\delta+}\text{Cl}^{\delta-}$ can also be used for iodochlorination of ynamides, but it is possible that it can act as an oxidizing agent. To test this, initially, we conducted the reaction with epoxy ynamides. We treated epoxy-ynamides **5a–c** with a 1M solution of ICl in CH_3CN (0.5 mL) at 0 °C for 10 min and obtained highly regioselective *E/Z*-isomeric mixture (1:1) of α -chloro- β -iodoenamide derivatives **6a–c** in good yields. In this transformation, along with addition reaction, intermolecular $\text{S}_{\text{N}}2$ nucleophilic ring-opening of epoxide part of the ynamide occurs simultaneously (Scheme 6). A band due to OH group is observed at $\sim 3534 \text{ cm}^{-1}$ in the IR spectra as expected. Here, electrophilic addition reactions are highly favored in contrast to the corresponding 5-*exo-dig* and/or 6-*endo-dig* cyclization reactions of epoxy-ynamides.^{15d–e} Although *E/Z* isomeric mixture was obtained, the *E*-isomer preferentially crystallized. The structures of the addition products **6a** and **6c** (Figure 2) were confirmed by single-crystal X-ray



Scheme 4. AlCl_3 Mediated Synthesis of (*E*)- α -chloroenamides from the Ynamides^a. ^a Conditions: **1** (0.20 mmol), AlCl_3 (0.22 mmol) in dimethyl carbonate (1 mL) and H_2O (0.20 mmol) at $80\text{ }^\circ\text{C}$ for 12 h. Isolated yields after column chromatography are given in parenthesis. For the preparation of compound **2u** we used AlCl_3 (0.44 mmol), H_2O (0.40 mmol).

diffraction; although the data was not great for **6c** due to low quality of the crystals, the structure could be readily ascertained. In compound **6a**, the C8-C9 distance [1.340(10) Å] proved the presence of a double bond between these two atoms. Three single bonds Cl(1)-C(8), C(12)-C(18) and I(1)-C(9) are newly formed. Similar structural features are observed in

compound **6c** also. Crystal structures of the compounds **6a** and **6c** show $\text{I}\cdots\text{O}$ halogen bonding interactions (Figure 2).¹⁸ We did perform a similar reaction of **1a** with ICl but ended up in obtaining a mixture that contained the hydrolyzed product. We did not succeed in obtaining a product without opening of the epoxide ring. Hence we did not pursue this further.






Figure 1. X-ray structures of compounds **2l** (left; CCDC 2033589) and **2q** (right; CCDC 2033590). Selected bond parameters: **2l** C11-C7 1.758(2), S1-N1 1.6520(15), N1-C7 1.404(2), C7-C8 1.326(3), C8-C9 1.468(3), C9-C10 1.396(3), C9-C14 1.390(3) (Å). **2q** C11-C7 1.770(3), S1-N1 1.662(2), N1-C7 1.414(3), C7-C8 1.316(4), C8-C9 1.484(4), C9-C10 1.385(4), C9-C14 1.388(4) (Å).



Scheme 5. Control Experiments.



Scheme 6. Reaction of epoxy ynamides with ICl.

4. Conclusions

An operationally simple approach for the regio- and stereo-selective synthesis of (*E*)- α -chloroenamides by the hydrochlorination of ynamides using AlCl₃ (as the source for hydrochlorination) is developed. This is complementary to other approaches reported in the literature, but emphasizes the point that AlCl₃ can also act as a hydrochlorinating agent (cf. ref. 14a) in the presence of even one equivalent of water; we also point out that a non-halogenated solvent, dimethyl carbonate, has been used in the present work. This




Figure 2. Molecular structures of compounds **6a** (Top; CCDC 2033592) and **6c**·CH₂Cl₂ (Bottom, solvent molecule is omitted; CCDC 2033593). Selected bond lengths [Å] with esds in parentheses: **6a** I1-C9 2.099(7), C9-C8 1.340(10), C11-C8 1.754(8), C16-C17 1.494(15), C18-C17 1.44(2), O3-C17 1.550(18), C12-C18 1.908(15), I...O 3.137(71). **6c**·CH₂Cl₂ I1-C12 2.116(10), C11-C12 1.315(13), C12-C13 1.447(13), C11-C11 1.747(10), C12-C10 1.789(15), O5-C9 1.464(16), N1-C11 1.411(12), I...O 3.011(15). Data quality for **6c** was only moderate but structure could be readily refined. The I...O distances are considerably shorter than the sum of the corresponding van der Waals radii of O (1.52 Å) and I (1.98 Å) i.e. they are <3.50 Å [Note: Average I-O covalent bond distance: 2.14 Å].

methodology is also applicable to phosphoramidate and carbamate derived ynamide substrate illustrating its utility. Deuterium labelling experiment demonstrated the essential role of water as a proton source in the reaction.

Supplementary Information

Experimental details; X-ray crystallographic data (cif files; CCDC nos. 2033589-2033593); ORTEPS of **2l**, **2q**, **2w**, **6a** and **6c**; ¹H / ¹³C / ³¹P NMR spectra. This material is available free of charge at www.ias.ac.in/chemsci.

Acknowledgements

We thank the Department of Science and Technology (DST, New Delhi) and University Grants Commission (UGC, New Delhi) for support. ASR, MA and Suraj thank UGC (New

Delhi) for fellowships. KCK thanks SERB for the J. C. Bose fellowship.

References

- Selected recent reviews: (a) Dekorver K A, Li H, Lohse A G, Hayashi R, Lu Z, Zhang Y and Hsung R P 2010 Ynamides: A Modern Functional Group for the New Millennium *Chem. Rev.* **110** 5064; (b) Evano G, Coste A and Jouvin K 2010 Ynamides: Versatile Tools in Organic Synthesis *Angew. Chem., Int. Ed.* **49** 2840; (c) Wang X-N, Yeom H-S, Fang L-C, He S, Ma Z-X, Kedrowski B L and Hsung R P 2014 Ynamides in Ring Forming Transformations *Acc. Chem. Res.* **47** 560; (d) Lu T and Hsung R P 2014 Novel ynamide structural analogues and their synthetic transformations *Arhivoc i* 127; (e) Evano G, Jouvin K and Coste A 2013 General Amination Reactions for the Synthesis of Ynamides *Synthesis* **45** 0017; (f) Pan F, Shu C and Ye L-W 2016 Recent progress towards gold-catalyzed synthesis of N-containing tricyclic compounds based on ynamides *Org. Biomol. Chem.* **14** 9456; (g) Prabagar B, Ghosh N and Sahoo A K 2017 Cyclization and Cycloisomerization of π -tethered ynamides: An expedient synthetic method to construct carbo- and heterocycles *Synlett* **28** 2539; (h) Evano G, Michelet B and Zhang C 2017 The anionic chemistry of ynamides: A review *C. R. Chimie* **20** 648; (i) Zhou B, Tan T-D, Zhu X-Q, Shang M and Ye L-W 2019 Reversal of Regioselectivity in Ynamide Chemistry *ACS Catal.* **9** 6393
- Lecomte M and Evano G 2016 Harnessing the Electrophilicity of Keteniminium Ions: A Simple and Straightforward Entry to Tetrahydropyridines and Piperidines from Ynamides *Angew. Chem., Int. Ed.* **55** 4547
- Selected examples: (a) Rettenmeier E, Schuster A M, Rudolph M, Rominger F, Gade C A and Hashmi A S K 2013 Gold Catalysis: Highly Functionalized Cyclopentadienes Prepared by Intermolecular Cyclization of Ynamides and Propargylic Carboxylates *Angew. Chem., Int. Ed.* **52** 5880; (b) Kiruthika S E, Nandakumar A and Perumal P T 2014 Synthesis of Pyrrolo-/Indolo[1,2-a]quinolines and Naphtho[2,1-b]thiophenes from gem-Dibromovinyls and Sulphonamides *Org. Lett.* **16** 4424 (c) Chen L, Cao J, Xu Z, Zheng Z-J, Cui Y-M, Xu L-W 2016 Lewis acid catalyzed [2+2] cycloaddition of ynamides and propargyl silyl ethers: synthesis of alkylidene-cyclobutenones and their reactivity in ring-opening and ring expansion *Chem. Commun.* **52** 9574
- Selected recent examples: (a) Fugino D, Yorimitsu H and Osuka A 2014 Regiocontrolled Palladium-Catalyzed Arylative Cyclizations of Alkynols *J. Am. Chem. Soc.* **136** 6255; (b) Karad S N and Liu R-S 2014 Regiocontrolled Gold-Catalyzed [2+2+2] Cycloadditions of Ynamides with Two Discrete Nitriles to Construct 4-Aminopyrimidine Cores *Angew. Chem., Int. Ed.* **53** 9072; (c) Briocche J, Meyer C and Cossy J 2015 *Org. Lett.* **17** 2800; (d) Adcock H V, Chatzopoulou E and Davies P W 2105 Divergent C \equiv H Insertion–Cyclization Cascades of N-Allyl Ynamides *Angew. Chem., Int. Ed.* **54** 15525; (e) Straker R N, Peng Q, Mekareeya A, Paton R S and Anderson E A 2016 Computational ligand design in enantio- and diastereoselective ynamide [5+2] cycloisomerization *Nat. Commun.* **7** 10109; (f) Cao Z, Zhu J, Liu L, Pang Y, Tian L, Sun X and Meng X 2019 AgNTf₂-catalyzed formal [3 + 2] cycloaddition of ynamides with unprotected isoxazol-5-amines: efficient access to functionalized 5-amino-1H-pyrrole-3-carboxamide derivatives *Beilstein J. Org. Chem.* **15** 2623; (g) Dutta S, Prabagar B, Vanjari R, Gandon V and Sahoo A K 2020 An Unconventional Sulfur-to-Selenium-to-Carbon Radical Transfer: Chemo- and Regioselective Cyclization of Yne-Ynamides *Green Chem.* **22** 1113; (h) Yang M, Wang X and Zhao J 2020 Ynamide-Mediated Macrolactonization *ACS Catal.* **10** 5230
- Some relevant examples: (a) Smith D L, Goundry W R F and Lam H W 2012 Palladium-catalyzed hydroacyloxylation of ynamides *Chem. Commun.* **48** 1505; (b) Yang Y, Wang L, Zhang J, Jin Y and Zhu G 2014 An unprecedented Pd-catalyzed trans-addition of boronic acids to ynamides *Chem. Commun.* **50** 2347; (c) Yang Y, Wang L, Zhang F and Zhu G 2014 Preparation of (Z)- α,β -Disubstituted Enamides via Palladium-Catalyzed Addition of Boronic Acids to Ynamides *J. Org. Chem.* **79** 9319; (d) Hu L, Xu S, Zhao Z, Yang Y, Peng Z, Yang M, Wang C and Zhao J 2016 Ynamides as Racemization-Free Coupling Reagents for Amide and Peptide Synthesis *J. Am. Chem. Soc.* **138** 13135; (e) Tona V, Ruider S A, Berger M, Shaaban S, Padmanaban M, Xie L-G, González L and Maulide N Divergent ynamide reactivity in the presence of azides – an experimental and computational study 2016 *Chem. Sci.* **7** 6032. (f) Li X Q, Jiang M Y, Zhan T Y, Cao W D and Feng X M 2020 Catalytic Asymmetric Three-component Hydroacyloxylation/1,4-Conjugate Addition of Ynamides *Chem. Asian J.* **15** 1953
- Recent examples: (a) Che J, Li Y, Zhang F, Zheng R, Bai Y and Zhu G 2014 Silver-promoted trans-hydrofluorination of ynamides: a regio- and stereoselective approach to (Z)- α -fluoroenamides *Tetrahedron Lett.* **55** 6240; (b) Xu S, Liu J, Hu D, Bi X Metal-free hydroacyloxylation and hydration reactions of ynamides: synthesis of α -acyloxyenamides and N-acylsulfonamides 2015 *Green Chem.* **17** 184; (c) Pirovano V, Negrato M, Abbiati G, Dell'Acqua M and Rossi E 2016 Gold-Catalyzed cis-Hydroarylation of Ynamides with Indoles: Regio and Stereoselective Synthesis of a Class of 2-Vinylindoles *Org. Lett.* **18** 4798; (d) Shu C, Shen C-H, Wang Y-H, Li L, Li T, Lu X and Ye L-W 2016 Synthesis of 2-Aza-1,3-butadienes through Gold-Catalyzed Intermolecular Ynamide Amination/C–H Functionalization *Org. Lett.* **18** 4630; (e) Sallio R, Corpet M, Habert L, Durandetti M, Gosmini C and Gillaizeau I 2017 Cobalt-Catalyzed Carbozincation of Ynamides *J. Org. Chem.* **82** 1254
- (a) He G, Qiu S, Huang H, Zhu G, Zhang D, Zhang R and Zhu H 2016 Cu(I)- or Ag(I)-Catalyzed Regio- and Stereocontrolled trans-Hydrofluorination of Ynamides *Org. Lett.* **18** 1856; (b) Métayer B, Compain G, Jouvin K, Martin-Mingot A, Bachmann C, Marrot J, Evano G and Thibaudeau S 2015 Chemo- and Stereoselective

- Synthesis of Fluorinated Enamides from Ynamides in HF/Pyridine: Second-Generation Approach to Potent Ureas Bioisosteres *J. Org. Chem.* **80** 3397; (c) Sato A H, Ohashi K and Iwasawa T 2013 Regio- and stereospecific synthesis of (*E*)- α -iodoenamide moieties from ynamides through iodotrimethylsilane-mediated hydroiodation *Tetrahedron Lett.* **54** 1309; (d) Zhu G, Qiu S, Xi Y, Ding Y, Zhang D, Zhang R, He G and Zhu H 2016 (IPr)CuF-catalyzed α -site regiocontrolled trans-hydrofluorination of ynamides *Org. Biomol. Chem.* **14** 7746; (e) Zeng X, Lu Z, Liu S, Hammond G B and Xu B 2017 Metal-free, Regio-, and Stereo-Controlled Hydrochlorination and Hydrobromination of Ynones and Ynamides *J. Org. Chem.* **82** 13179; (f) Ohashi K, Mihara S, Sato A H, Ide M and Iwasawa T 2014 Synthesis of 1-haloethenamides from ynamide through halotrimethylsilane-mediated hydrohalogenation *Tetrahedron Lett.* **55** 632; (g) Ide M, Ohashi K, Mihara S and Iwasawa T 2014 Regio- and stereoselective hydrohalogenation of ynamide components in 1,3-butadiynes with in situ generated HX *Tetrahedron Lett.* **55** 2130; (h) Zeng X J, Li J L, Ng C K, Hammond G B and Xu B 2018 (Radio)Fluoro-Click Reaction Enabled by a Hydrogen Bonding Cluster *Angew. Chem., Int. Ed.* **57** 2924 (i) Li J-L, Lin E, Han X-L, Li Q and Wang H 2019 Synthesis of α -Fluorinated Imides via Direct Fluorohydroxylation of Ynamides *Org. Lett.* **21** 4255 (j) Kim S W, Uma T-W and Shin S 2017 Brønsted Acid-Catalyzed α -Halogenation of Ynamides from Halogenated Solvents and Pyridine-*N*-Oxides *Chem. Commun.* **53** 1
8. (a) Hirano S, Tanaka R, Urabe H and Sato F 2004 Practical Preparation of *N*-(1-Alkynyl)sulfonamides and Their Remote Diastereoselective Addition to Aldehydes via Titanation *Org. Lett.* **6** 727; (b) Saito N, Katayama T and Sato Y 2008 Nickel-Catalyzed Highly Regioselective Multicomponent Coupling of Ynamides, Aldehydes, and Silane: A New Access to Functionalized Enamides *Org. Lett.* **10** 3829; (c) Graux L V, Clavier H and Buono G 2014 Palladium-Catalyzed Addition of 1,3-Diones to Ynamides: An Entry to Alkoxy-Substituted Enamides *ChemCatChem.* **6** 2544
 9. (a) Fadel A, Legrand F, Evano G and Rabasso N 2011 Highly Regio- and Stereoselective Nickel-Catalyzed Addition of Dialkyl Phosphites to Ynamides: an Efficient Synthesis of β -Aminovinylphosphonates *Adv. Synth. Catal.* **353** 263; (b) Dwivedi V, Hari Babu M, Kant R and Sridhar Reddy M 2015 *N*-Substitution dependent stereoselectivity switch in palladium catalyzed hydroalkynylation of ynamides: a regio and stereoselective synthesis of ynamides *Chem. Commun.* **51** 14996; (c) Maity P, Klos M R and Kazmaier U 2013 Syntheses of α -Stannylated and α -Iodinated Enamides via Molybdenum-Catalyzed Hydrostannation *Org. Lett.* **15** 6246
 10. (a) Mulder J A, Kurtz K C M, Hsung R P, Coverdale H, Frederick M O, Shen L and Zifcick C A 2003 Highly Stereoselective Synthesis of Novel α -Haloenamides via a Mild and Efficient Hydrohalogenation of Ynamides *Org. Lett.* **5** 1547; (b) Yabuuchi Y, Kuzuguchi T, Yoshimura T and Matsuo J 2016 Synthesis of α -Halo- γ -hydroxyenamides by Titanium Tetrahalide Mediated Addition of Aldehydes or Ketones to Ynamides *Org. Lett.* **18** 4951; (c) Prabagar B, Nayak S, Mallick R K, Prasad R and Sahoo A K 2016 Triphenylphosphine promoted regio and stereoselective α -halogenation of ynamides *Org. Chem. Front.* **3** 110; (d) Lu Z, Kong W, Yuan Z, Zhao X and Zhu G 2011 Synthesis of Multisubstituted Enamides via Pd-Catalyzed Chloroallylation of Ynamides *J. Org. Chem.* **76** 8524; (e) Cao W, Chen P, Wang L, Wen H, Liu Y, Wang W and Tang Y 2018 A Highly Regio- and Stereoselective Syntheses of α -Halo Enamides, Vinyl Thioethers, and Vinyl Ethers with Aqueous Hydrogen Halide in Two-Phase Systems *Org. Lett.* **20** 4507
 11. (a) Nakanishi M, Minard C, Retailleau P, Cariou K and Dodd R H 2011 Copper(I) Catalyzed Regioselective Asymmetric Alkoxyamination of Aryl Enamide Derivatives *Org. Lett.* **13** 5792; (b) Carbery D R 2008 Enamides: valuable organic substrates *Org. Biomol. Chem.* **3455**; (c) Matsubara R and Kobayashi S 2008 Enamides and Encarbamates as Nucleophiles in Stereoselective C–C and C–N Bond-Forming Reactions *Acc. Chem. Res.* **41** 292; (d) Ma Z-X, Feltenberger J B and Hsung R P 2012 Total Syntheses of Chelidonine and Norchelidonine via an Enamide_Benzyne_[2 + 2] Cycloaddition Cascade *Org. Lett.* **14** 2742; (e) Kuranaga T, Sesoko Y and Inoue M 2014 Cu-mediated enamide formation in the total synthesis of complex peptide natural products *Nat. Prod. Rep.* **31** 514
 12. (a) Gourdet B and Lam H W 2010 Catalytic Asymmetric Dihydroxylation of Enamides and Application to the Total Synthesis of (+)-Tanikolide *Angew. Chem., Int. Ed.* **49** 8733; (b) St. Denis J D, Zajdlik A, Tan J, Trinchera P, Lee C F, He Z, Adachi S and Yudin A K 2014 Boron-Containing Enamine and Enamide Linchpins in the Synthesis of Nitrogen Heterocycles *J. Am. Chem. Soc.* **136** 17669; (c) Song Z, Lu T, Hsung R P, Al-Rashid Z F, Ko C and Tang Y 2007 Stereoselective Simmons–Smith Cyclopropanation of Chiral Enamides *Angew. Chem., Int. Ed.* **46** 4069
 13. (a) Henderson W E and Gangloff W 1916 The action of anhydrous aluminium chloride upon unsaturated compounds *J. Am. Chem. Soc.* **18** 1382; (b) Wang C and Xi Z 2007 Co-operative effect of Lewis acids with transition metals for organic synthesis *Chem. Soc. Rev.* **36** 1395
 14. (a) Yang Y, Liu H, Peng C, Wu J, Zhang J, Qiao Y, Wang X-N and Chang J 2016 AlCl₃-Catalyzed Annulations of Ynamides Involving a Torquoselective Process for the Simultaneous Control of Central and Axial Chirality *Org. Lett.* **18** 5022; (b) Oppenheimer J, Johnson W L, Tracey M R, Hsung R P, Yao P-Y, Liu R and Zhao K 2007 A Rhodium(I)-Catalyzed Demethylation-Cyclization of *o*-Anisole-Substituted Ynamides in the Synthesis of Chiral 2-Amido Benzofurans *Org. Lett.* **9** 2361; (c) Chen L, Yu L, Deng Y, Cui Y, Bian G and Cao J 2016 Synthesis of α -haloenamides via zinc halide mediated direct addition of benzhydryl halides to ynamides *Org. Biomol. Chem.* **14** 564
 15. Our earlier work on ynamides: (a) Siva Reddy A, Nagarjuna Reddy M and Kumara Swamy K C 2014 A simple copper-catalysed tandem cyclisation of ynamides leading to triazolo-1,2,4-benzothiadiazine-1,1-dioxides

- in PEG-400 medium *RSC Adv.* **4** 28359; (b) Siva Reddy A and Kumara Swamy K C 2015 Use of Elemental Sulfur or Selenium in a Novel One-Pot Copper-Catalyzed Tandem Cyclization of Functionalized Ynamides Leading to Benzosultams *Org. Lett.* **17** 2996; (c) Siva Reddy A, Kumari, A L S, Saha S and Kumara Swamy K C 2016 Palladium-Catalyzed Tandem-Cyclization of Functionalized Ynamides: An Approach to Benzosultams *Adv. Synth. Catal.* **358** 1625; (d) Kumari A L S, Siva Reddy A and Kumara Swamy K C 2016 Transition Metal-Free Cascade Cyclization of Epoxy-Ynamides: To Go for 1,3-Oxazines or 1,4-Oxazines? *Org. Lett.* **18** 5752; (e) Anitha M, Shankar M and Kumara Swamy K C 2019 Reactivity of epoxy-ynamides with metal halides: nucleophile (Br/Cl/OH)-assisted tandem intramolecular 5-exo-dig or 6-endo-dig cyclisation and AgF₂-promoted oxidation *Org. Chem. Front.* **6** 1133; (f) Siva Reddy A and Kumara Swamy K C 2017 Ethanol as a Hydrogenating Agent: Palladium-Catalyzed Stereoselective Hydrogenation of Ynamides To Give Enamides *Angew. Chem. Int. Ed.* **56** 6984
16. (a) Ho M L, Flynn A B and Ogilvie W W 2007 Single-Isomer Iodochlorination of Alkynes and Chlorination of Alkenes Using Tetrabutylammonium Iodide and Dichloroethane *J. Org. Chem.* **72** 977; (b) Ide M, Yauchi Y and Iwasawa T 2014 Regio-, and Stereoselective Iodobromination of Ynamides for Synthesis of (*E*)-1-Bromo-2-iodoenamides *Eur. J. Org. Chem.* 3262
17. (a) Zhang Y, Hsung R P, Tracey M R, Kurtz K C M and Vera E L 2004 Copper Sulfate-Pentahydrate-1,10-Phenanthroline Catalyzed Amidations of Alkynyl Bromides. Synthesis of Heteroaromatic Amine Substituted Ynamides *Org. Lett.* **6** 1151; (b) Dateer R B, Shaibu B S and Liu R S 2012 Gold-Catalyzed Intermolecular [4+2] and [2+2+2] Cycloadditions of Ynamides with Alkenes *Angew. Chem., Int. Ed.* **51** 113
18. For another recent example, see: Anitha M and Kumara Swamy K C 2019 Highly functionalised (γ -azido/ γ -fluoro- β -iodo/) vinyl derivatives from phosphorus based allenes or allenates: I...O halogen bonding interactions *Org. Biomol. Chem.* **17** 5736