

Air-sea exchange of CO₂ in the Gulf of Kutch, northern Arabian Sea based on bomb-carbon in corals and tree rings

S CHAKRABORTY, R RAMESH and S KRISHNASWAMI

Physical Research Laboratory, Ahmedabad 380009

Abstract. Radiocarbon analyses were carried out in the annual bands of a 40 year old coral collected from the Gulf of Kutch (22°6'N, 70°E) in the northern Arabian Sea and in the annual rings of a teak tree from Thane (19°14'N, 73°24'E) near Bombay. These measurements were made in order to obtain the rates of air-sea exchange of CO₂ and the advective mixing of water in the Gulf of Kutch. The $\Delta^{14}\text{C}$ peak in the Thane tree occurs in the year 1964, with a value of $\sim 630\%$, significantly lower than that of the mean atmospheric $\Delta^{14}\text{C}$ of the northern hemisphere ($\sim 1000\%$). The radiocarbon time series of the coral was modelled considering the supply of carbon and radiocarbon to the gulf through air-sea exchange and advective water transport from the open Arabian Sea. A reasonable fit for the coral data was obtained with an air-sea CO₂ exchange rate of $11\text{--}12 \text{ mol m}^{-2} \text{ yr}^{-1}$, and an advective velocity of 28 m yr^{-1} between the Arabian Sea and the Gulf of Kutch; this was based on a model generated time series for radiocarbon in the Arabian Sea. The deduced velocity ($\sim 28 \text{ m yr}^{-1}$) of the advective transport of water between the gulf and the Arabian Sea is much lower than the surface tidal current velocity in this region, but can be understood in terms of net fluxes of carbon and radiocarbon to the gulf to match the observed coral $\Delta^{14}\text{C}$ time series.

Keywords. Air-sea interaction; CO₂ exchange; Gulf of Kutch; coral; tree rings; bomb radiocarbon.

1. Introduction

The oceans of the world contain about sixty times as much CO₂ as the atmosphere and regularly exchange CO₂ with it. Thus, they play a significant role in controlling the CO₂ content of the atmosphere and hence the greenhouse warming of our planet (Dickson 1992). The oceanic capacity for uptake of CO₂ is influenced by physical, chemical and biological processes occurring within the sea. One of the aims of the JGOFS (Joint Global Ocean Flux Study), an IGBP core project, is to determine the rate of exchange of carbon dioxide between the atmosphere and the oceans, for a better understanding of the global carbon cycle. The air-sea CO₂ exchange rate (ASCER), has been determined by various investigators in several oceanic regions (Sundquist 1985 and references therein). Such measurements are however meagre in the Arabian Sea, a unique biogeochemical province of the oceans. Realizing the importance of Arabian Sea in the global carbon cycle, detailed JGOFS related studies are planned in this region during the next 2–3 years.

ASCER can be determined by comparing the radiocarbon time histories of the atmosphere and the ocean. The surface water radiocarbon history can be derived from ¹⁴C data in dated coral bands, and the atmospheric radiocarbon variation can be obtained by measuring ¹⁴C concentrations in tree rings. Druffel and Suess (1983) estimated the ASCER in the northwest Atlantic and eastern tropical Pacific region

Table 1. $\Delta^{14}\text{C}$ minimum, maximum and peak shift for different oceanic regions. The minima of $\Delta^{14}\text{C}_{\text{coral}}$ represent the pre-bomb values. The time difference between the peaks in $\Delta^{14}\text{C}$ of the atmosphere and the coral (surface ocean) is termed 'peak shift'. The atmospheric peak was taken to be in the year 1963.

Oceanic region	$\Delta^{14}\text{C}_{\text{coral}}$		Year of maximum	Peak shift (yr)	Reference
	min	max			
Bermuda 32°N, 65°W	-50	168	1975	12	Nozaki <i>et al</i> (1978).
Florida 24°57'N 88°33'W	-55	160	1972	9	Druffel and Linick (1978)
Oahu 21°N, 158°W	—	175	1973	10	Toggweiler <i>et al</i> (1991).
Belize 16°50'N 83°48'W	-55	160	1972	9	Druffel (1980).
North Red Sea 12°N, 43°E	-75	105	1967	4	Cember (1989).
South Red Sea 11.5°N, 43°E	-73	~45	1972	>7	Cember (1989).
Tarawa 1°N, 172°E	—	103	1978	15	Toggweiler <i>et al</i> (1991).
Galapagos 1°S, 90°W	-72	25	1973	10	Druffel (1981).
Fiji 18°S, 179°E	-70	138	1974	11	Toggweiler <i>et al</i> (1991).
Tonga 20°S, 175°W	—	157	1975	12	Toggweiler <i>et al</i> (1991).
Gulf of Kutch 22.6°N, 70°E	-65	172	1968	5	This work.

using coralline $\Delta^{14}\text{C}$. Cember (1989) estimated the ASCER in the Red Sea following a similar approach. The earlier results are summarized in table 1. We have followed this approach to determine ASCER in the northeast Arabian Sea.

2. Methods

We collected a coral *Favia speciosa* from the Pirotan island of the Gulf of Kutch (GKh, 22.6°N, 70°E), northeastern Arabian Sea, (water depth during low tide at coral location ≤ 1 m) in June 1990 (figure 1). This gulf is a part of the western continental shelf of the Indian coast and supports fringing reefs. Tree rings from a teak tree (*Tectona grandis*) that grew in Thane (19°14'N, 73°24'E) near Bombay was also analyzed for radiocarbon. A disc from the stump and its chronology was provided by Pant and Borgaonkar (1983).

A ~ 1 cm thick slice of the coral was sectioned along its growth axis using a diamond tipped circular saw. X-radiography revealed clear bands with a mean band thickness of ~ 4 mm. Oxygen isotope studies showed that these bands were annual in nature,




Figure 1. The Gulf of Kutch showing water depth contours (m) and the coral location (Pirotan island). Inset shows the locations for coral, tree and the GEOSECS station 416.

with an average growth rate of 4 ± 0.8 mm/yr (Chakraborty and Ramesh 1994) for the past 40 years. Because of the low growth rate, 2 or 3 bands were combined for radiocarbon measurements. Radiocarbon measurements were made by the liquid scintillation technique (Chakraborty 1993; Bhushan *et al* 1994) using a TASK benzene synthesizer and a Packard 2250CA Liquid Scintillation Analyzer. The radiocarbon activity is reported as $\Delta^{14}\text{C}$ following standard terminology (Stuiver and Polach 1977). Fractionation correction was done using measured $\delta^{13}\text{C}$ values, (average of a number of samples per band). The reported $\Delta^{14}\text{C}$ values are decay corrected to the year of formation.

For the measurement of radiocarbon in tree rings, wood was powdered and chemically pretreated according to the method of Cain and Suess (1976). Samples were combusted in oxygen to produce CO₂ which was converted to benzene (Noakes *et al* 1965). $\delta^{13}\text{C}$ measurements were not made in these tree rings but was done in the cellulose from an adjacent tree (Ramesh *et al* 1989) which shows values in the range of -24 to -26‰ . We used a value of -25‰ for $\Delta^{14}\text{C}$ calculations. The overall analytical uncertainty in $\Delta^{14}\text{C}$ measurement is $\pm 8\text{‰}$ (1σ).

3. Features of the data

3.1 Gulf of Kutch coral

Figure 2 shows the $\Delta^{14}\text{C}$ time-series in the GKc coral for the years 1950–1990. From a value of -60‰ in the year 1950 it steadily increases to a peak value of 170‰ in 1968, after which it decreases monotonically to a value of 55‰ in 1990. The increase in ^{14}C activity since 1950 results from the injection of bomb produced radiocarbon from the atmosphere to the surface ocean via air-sea CO₂ exchange. Moore and Krishnaswami (1974) had reported $\delta^{14}\text{C}$ time series for a *Favia* coral in the GKc




Figure 2. $\Delta^{14}\text{C}$ time series in the coral *Favia speciosa* from the Gulf of Kutch, northern Arabian Sea. The continuous curve is an eye fit line to the data.

for the years 1951 to 1973. They obtained a peak value of $+230\text{‰}$ for $\delta^{14}\text{C}$ in 1968. This $\delta^{14}\text{C}$ when corrected for isotopic fractionation using a $\delta^{13}\text{C}$ value of -0.27‰ (our measured average value in the GKc coral) yields a $\Delta^{14}\text{C}$ of 169‰ , very close to the peak value of 170‰ obtained in this study. Measurements of radiocarbon activity in the water of Pirotan island in 1992 yielded a value of $46 \pm 8\text{‰}$ (Bhushan *et al* 1994) which is consistent with the $\Delta^{14}\text{C}$ value ($55 \pm 7\text{‰}$) of the coral bands for the years 1988–90.

3.2 Tree rings

The $\Delta^{14}\text{C}$ profile in the annual rings of the teak tree from Thane is shown in figure 3 along with the northern hemispheric (NH) curve (Nydal and Lövseth 1983; Gupta and Polach 1985). The Thane tree ring samples analyzed cover a time span of 1960 to 1980. The prebomb (1934–43 A.D.) tree ring $\Delta^{14}\text{C}$ value was $-25 \pm 7.9\text{‰}$ for the same species (*Tectona grandis*) collected from Kerala (9°N , $76^\circ 5'\text{E}$; Kusumgar 1965). This is in agreement with Stuiver and Quay's (1981) prebomb $\Delta^{14}\text{C}$ values of -22‰ from tree rings. In the early sixties, the $\Delta^{14}\text{C}$ values of the Thane tree rings are indistinguishable from that of the NH values, but the two sets of values differ significantly after mid-sixties through the year 1980. The Thane tree ring $\Delta^{14}\text{C}$ reaches a peak value (630‰) during 1964–65, similar to that for air samples from Bombay during 1963–64 (Kusumgar 1965) which yield values of $\sim 700\text{‰}$. The trend of $\Delta^{14}\text{C}$ time series in the Thane teak tree is similar to that of NH curve, but the $\Delta^{14}\text{C}$ values at Thane are significantly lower than those of NH. For example the peak $\Delta^{14}\text{C}$ value for NH is $\sim 1000\text{‰}$ compared to 630‰ for Thane. Broecker *et al* (1985) have reported atmospheric $\Delta^{14}\text{C}$ time series for three regions: $>20^\circ\text{N}$, $20^\circ\text{N}-20^\circ\text{S}$, and $>20^\circ\text{S}$. The tree analyzed in this study (located at 19°N) belongs to the middle group and its peak $\Delta^{14}\text{C}$ value is $\sim 100\text{‰}$ less compared to that expected from the $20^\circ\text{N}-20^\circ\text{S}$ curve. The reason for the consistently lower $\Delta^{14}\text{C}$ in the tree could not be due to ^{14}C depleted CO_2 from the Arabian Sea, as pre-bomb values for the same species are in good agreement with global values, as mentioned above. More teak trees should be analyzed to verify the above results.




Figure 3. Atmospheric $\Delta^{14}\text{C}$ time variations of the Northern Hemisphere (long dashed line, data from Nydal and Lövseth 1983) and Thane (near Bombay, solid line). The Thane data are based on tree ring measurements. The Thane values are significantly lower than the NH curve after 1963. Typical 1σ error in the $\Delta^{14}\text{C}$ measurement of tree ring is $\pm 7\%$.

4. Air-Sea CO₂ exchange rate (ASCER) in the Gulf of Kutch

4.1 Model for calculation of ASCER

Determination of the CO₂ exchange rate between atmosphere and the surface ocean requires a knowledge of sources and sinks of carbon and radiocarbon in the particular oceanic region. The carbon budget in surface water is controlled mainly by air-sea CO₂ exchange, advective processes (lateral and/or vertical mixing of water masses) and biological productivity. The GK_h area is a shallow continental shelf region having a mean water depth of ~ 30 m. It exchanges water with the adjoining Arabian Sea through tidal currents. The speed of these currents vary from 0.5 to 0.76 m/sec and are mainly towards the west and southeast directions (Srivastava and John 1977). The mass balance equations for ¹²C and ¹⁴C in the GK_h between supply and removal is given by:

$$\left\{ \begin{array}{l} \text{Invasion of CO}_2 + \\ \text{advective transport} \end{array} \right\} = \left\{ \begin{array}{l} \text{Evasion of CO}_2 + \text{advective transport} + \\ \text{biological removal.} \end{array} \right\}$$

For ¹²C this is given by (assuming invasion and evasion of CO₂ are equal)

$$F_{12} + wC_S = F_{12} + wC_G + B \quad (1)$$

$$wC_S = B + wC_G \quad (2)$$

where F_{12} ($\text{mol m}^{-2} \text{yr}^{-1}$) is the CO₂ exchange flux, w (m yr^{-1}) lateral advection velocity, C (mol m^{-3}) carbon concentration and B ($\text{mol m}^{-2} \text{yr}^{-1}$) is the biological removal rate of carbon. The subscripts S and G refer to the surface Arabian Sea and the GK_h respectively. The above equations assume steady state for ¹²C balance. This assumption is an oversimplification as the CO₂ concentration of atmosphere has been increasing steadily during the past century (Bacastow and Keeling 1981; Siegenthaler and Sarmiento 1993) which would affect the invasion-evasion balance.

The mass balance equation for ^{14}C is of the following form (neglecting the radiocarbon decay term).

$$D_G \frac{dC_G^*}{dt} = F_{12}(R_A - R_G) + w(C_S R_S - C_G R_G) \quad (3)$$

where R is the $^{14}\text{C}/^{12}\text{C}$ mol ratio in the respective reservoirs. Subscripts A and G stand for the atmosphere and GKh respectively. C_G^* is the radiocarbon concentration in GKh and D_G is the mean depth of GKh (table 2). The formulation of our model is similar to that used by Cember (1989) for deriving ASCER in the Red Sea region. Equation 3 can be written in the standard $\Delta^{14}\text{C}$ notation. The first term in the right hand side, the net ^{14}C flux, between the atmosphere and the GKh can be expressed as: $k(\Delta^{14}\text{C}_A - \Delta^{14}\text{C}_G)F_{12}$, where k is a factor which takes into consideration the $^{14}\text{C}/^{12}\text{C}$ mol ratio of the NBS oxalic acid standard (1.176×10^{-12}) and fractionation factor for the inter reservoir carbon transfer (Stuiver 1980). The numerical value of k is 1.24×10^{-15} (*op cit.*). $\Delta^{14}\text{C}_A$ and $\Delta^{14}\text{C}_G$ are the atmospheric and Gkh radiocarbon activities. The second term in the right hand side of equation (3) represents the exchange flux of radiocarbon between the Arabian Sea and GKh by advection. This term can be written as:

$$F_{SG} = w \left[m(\Sigma\text{CO}_2)_S \left(1 + \frac{\Delta^{14}\text{C}_S}{1000} \right) - m(\Sigma\text{CO}_2)_G \left(1 + \frac{\Delta^{14}\text{C}_G}{1000} \right) \right] \quad (4)$$

Table 2. Symbols, units and values of parameters used in our model.

Symbol	Parameter	Unit	Value	Reference
F_{12}	Air-sea exchange rate of CO_2	$\text{mol m}^{-2} \text{yr}^{-1}$	model derived	
$C_S, (\Sigma\text{CO}_2)_S$	Total DIC conc. in surface Arabian Sea	mol m^{-3}	2.11	Stuiver and Östlund (1983).
$C_D, (\Sigma\text{CO}_2)_D$	Total DIC conc. of deep Arabian Sea	mol m^{-3}	2.36	Stuiver and Östlund (1983).
$C_G, (\Sigma\text{CO}_2)_G$	Total DIC conc. in GKh	mol m^{-3}	2.11	Stuiver and Östlund (1983).
R	$^{14}\text{C}/^{12}\text{C}$ mol ratio	—	variable	
C^*	Radiocarbon conc.	mol m^{-3}	variable	
D_S	Mixed layer depth of Arabian Sea	m	60	
D_G	Mean depth of GKh	m	30	
$(\Delta^{14}\text{C})_A^0$	Pre-bomb atmospheric $\Delta^{14}\text{C}$	‰	-25	Stuiver and Quay (1981).
u	Upwelling velocity	m yr^{-1}	model derived	
w	Advective velocity Arabian Sea to GKh	m yr^{-1}	model derived	

where m is a constant which is obtained by multiplying the $^{14}\text{C}/^{12}\text{C}$ mol ratio of the NBS Oxalic acid standard and the fractionation term between the nineteenth century wood standard and the marine bicarbonate (Stuiver 1980). That is, $m = (1.176 \times 10^{-12} \times 1.052) = 1.24 \times 10^{-12}$. The solution of equation (3) is given by:

$$\Delta^{14}\text{C}_G(t) = \Delta^{14}\text{C}_G^0 e^{-Jt} + E e^{-Jt} \int_0^t e^{J\tau} \Delta^{14}\text{C}_A(\tau) d\tau + F e^{-Jt} \int_0^t e^{J\tau} \Delta^{14}\text{C}_S(\tau) d\tau \quad (5)$$

where $\Delta^{14}\text{C}_G^0$ is the initial value (pre-bomb) of $\Delta^{14}\text{C}$ in GKh, J , E and F are constants ($E = F_{12}/D_G \Sigma\text{CO}_2$, $F = W/D_G$, $J = E + F$). The values of various parameters used in this model are given in table 2.

In equation (3), the term containing F_{12} is always positive as $\Delta^{14}\text{C}_A > \Delta^{14}\text{C}_{GK}$ during the period under consideration. In contrast, the term containing w is always negative because $R_G > R_S$ ($C_S = C_G$, we neglect the biological removal B , for simplicity), if the trend seen in 1977 GEOSECS (Östlund *et al* 1987) observations is assumed to hold good for the entire period under consideration. Therefore it is possible to get a number of solutions for equation (3) such that they are consistent with the observed $\Delta^{14}\text{C}$ in the GKh coral, as long as the ratio F_{12}/w is kept constant. The timing of the maximum in $\Delta^{14}\text{C}_G$ can be used as a constraint to determine the value of F_{12}/w . At maximum, $dC/dt = 0$ and hence F_{12}/w is given by $[(C_G R_G - C_S R_S)/(R_A - R_S)]_{1968}$. This is numerically equal to ~ 0.4 . Therefore under the assumptions that F_{12} and w are constant during the period of investigation, a number of 'best fit' solutions are possible for equation (3), for the ratio of 0.4 for F_{12}/w . It is difficult to constrain the values of F_{12} and w based only on the model as the $\Delta^{14}\text{C}$ time series trend is governed by the ratio of F_{12}/w such that $F_{12}/w \sim 0.4$. Independent estimates of w are needed to arrive at unique values of F_{12} . As a first step we calculate $\Delta^{14}\text{C}_G(t)$ using (5) with two simplifications, (i) $[\Sigma\text{CO}_2]_S = [\Sigma\text{CO}_2]_G$ and (ii) the advective transport process is negligible, i.e. the ^{14}C time variations in the GKh is controlled only by carbon exchange with the atmosphere. Druffel and Suess (1983) also modelled their ^{14}C data in corals from north-west Atlantic and eastern tropical Pacific to derive exchange time scales and CO₂ exchange rate based on a similar simplified model. Equation 3 with $w = 0$ reduces to the model used by Druffel and Suess (1983) i.e. the time derivative of ^{14}C concentration in sea surface water is proportional to the air-sea gradient of $\Delta^{14}\text{C}$.

Solving (5) with $w = 0$ and for different F_{12} values we obtain a family of curves (figure 4). The curve (a) with $F_{12} = 2 \text{ mol m}^{-2} \text{ yr}^{-1}$ shows that the GKh radiocarbon activity increases monotonically through 1980 without a distinct peak. This is not consistent with our coral data (filled circles in figure 4) which show a distinct peak in 1968. The second, third and fourth curves (b), (c) and (d) with $F_{12} = 6, 8$ and $10 \text{ mol m}^{-2} \text{ yr}^{-1}$ respectively, though show peaks in the $\Delta^{14}\text{C}$ time series, they predict much higher radiocarbon concentration in the GKh compared to that observed in the coral. It is clear from these results that the air-sea exchange alone is not able to simulate the observed $\Delta^{14}\text{C}$ variations in the GKh coral. Therefore carbon and radiocarbon supply to the GKh via advection from the Arabian Sea also needs to be considered, which would 'dilute' the effects of air-sea exchange and thereby provide a better fit to the coral data.

The advective supply of carbon to the GKh could be from the surface layer or




Figure 4. Calculated time variations of radiocarbon activity in the Gulf of Kutch considering only air-sea exchange of CO_2 . (a) $F_{12} = 2$, (b) $F_{12} = 6$, (c) $F_{12} = 8$, (d) $F_{12} = 10 \text{ mol m}^{-2} \text{ yr}^{-1}$. Solid line: simulated curve; filled circle: coral data.

shallow depths of the Arabian Sea. Solution of (5) requires knowledge of time variations in R_S , $^{14}\text{C}/^{12}\text{C}$ ratio of the Arabian sea water contributing radiocarbon to GKk. Since there is no data on $\Delta^{14}\text{C}$ time series of the surface Arabian Sea, we generated the same for the time period 1953 to 1980 (figure 5), based on input from atmosphere and upwelling from the deep Arabian Sea (the detailed calculations are presented in Chakraborty 1993). The dotted curves in this figure represent the simulated radiocarbon activities, $\pm 20\%$ wrt the solid line which was obtained with $F_{12} = 11 \text{ mol m}^{-2} \text{ yr}^{-1}$ and u (upwelling velocity in the northern Arabian Sea) = 10.5 myr^{-1} . Though we constrained this time series with the 1977 mixed layer $\Delta^{14}\text{C}$ value of 59‰ (dot in figure 5) based on the GEOSECS (station 416; Stuiver and Ostlund 1983) measurements, the trend of the time series is not necessarily unique, as values for various parameters used were based not on actual measurements but on reasonable assumptions. However the derived time series provides a likely $\Delta^{14}\text{C}$ data set for the Arabian Sea.

Now we use the Arabian Sea and the atmospheric $\Delta^{14}\text{C}$ time series as the two end members contributing carbon and radiocarbon to GKk to generate the $\Delta^{14}\text{C}$ time series in the GKk; we generate a family of curves for different sets of values of F_{12} and w (figure 6). The solid line is the calculated $\Delta^{14}\text{C}$ time series for the GKk, appear to fit well. Dashed lines represent the model fit with different F_{12} and w values. The filled circles are our coral data.

It is seen from figure 6 that the rise in $\Delta^{14}\text{C}$ of coral during 1953–1960 is faster than that predicted by the model. A factor which would have contributed to this is the




Figure 5. Calculated time variations of radiocarbon activity in the mixed layer of the northern Arabian Sea (GEOSECS Sta 416). See text for explanation. Filled circle is the only data available.




Figure 6. Results of box model calculations. The plots show the simulated radiocarbon time series in the Gulf of Kutch (GKh) for different sets of parameters. (a) constant F_{12} ($= 12 \text{ mol m}^{-2} \text{ yr}^{-1}$) and varying w ($= 20, 28, 36 \text{ m yr}^{-1}$), (b) constant w ($= 28$) and varying F_{12} ($= 18, 12, 6$), (c) constant F_{12} ($= 50$) and varying w ($= 70, 100, 130$) and (d) constant w ($= 216$) and varying F_{12} ($= 70, 100, 130$). Filled circles represent GKh coral data with $\pm 1\sigma$ errors. Solid lines in each case appears to fit well. Dashed lines represent the model fit with different F_{12} and w values.

averaging effect i.e., the $\Delta^{14}\text{C}$ plotted for the year 1957.5 is an average for five coral bands 1955–1960. Another possible cause for this discrepancy can be an error in the year assignment for the bands. As we have taken utmost care in sampling and verified the year assignments with the $\delta^{18}\text{O}$ data this error is unlikely; however, if at all there is an error it is likely to be ± 1 yr.

Our calculations show that the timing of $\Delta^{14}\text{C}$ peak in the GKh is controlled more by F_{12} , whereas both F_{12} and w seem to influence the magnitude of the $\Delta^{14}\text{C}$ peak. A reasonable fit to the coral data can be made using the values of $F_{12} = 12 \text{ mol m}^{-2} \text{ yr}^{-1}$ and $w = 28 \text{ m yr}^{-1}$ based on the $\Delta^{14}\text{C}$ time series of the two end members considered. In this case when $F_{12} = 0$ the $\Delta^{14}\text{C}_G$ approaches $\Delta^{14}\text{C}_s$ with a time constant of ~ 1 yr and when $w = 0$ the $\Delta^{14}\text{C}_G$ approaches to that of atmospheric activity with a time constant of about 5 yrs.

Figure 7 shows the simulated $\Delta^{14}\text{C}$ variations for the gulf. All the curves were plotted with $F_{12} = 12$ and $w = 28$. The input from the Arabian Sea for the solid line was considered with $F_{12} = 11$ and $u = 10.5$ (the solid line in figure 5) and those for the dotted lines were plotted considering the inputs from the Arabian Sea which are 20% off the best fit line (dotted lines in figure 5). Filled circles are our coral data.

If the $\Delta^{14}\text{C}$ values of the Arabian Sea end member contributing radiocarbon to GKh is different from that used in the above model, then the values F_{12} and w for GKh would also differ correspondingly. However, for $< 20\%$ change, there is no significant effect on the coral $\Delta^{14}\text{C}$ (figure 5). There are no reported results on the circulation and hydrographic properties of the GKh and hence it is difficult at present to constrain our model further to have a unique solution. However it is possible to estimate the exchange rate by some other independent means. One of them is to calculate the exchange rate using the ^{14}C inventory. Broecker *et al* (1985) have calculated the inventory of bomb ^{14}C in station 416 to be $5.1 \times 10^9 \text{ atom cm}^{-2}$. According to Stuiver (1980) the inventory is related to the CO_2 air-sea exchange flux as:

$$Q = 1.24 \times 10^{-15} F_{12} \int (\Delta^{14}\text{C}_A - \Delta^{14}\text{C}_s - 45) dt$$




Figure 7. Calculated time variations of the GKh radiocarbon activity considering different inputs from the Arabian Sea. Solid line which appears to fit well was plotted with the Arabian Sea input (solid line in figure 5) and the dotted lines in figure 5 are the corresponding inputs for these two lines.

The value of the integrand between 1953–1977 is $\sim 5500\%$ yr based on our model derived $\Delta^{14}\text{C}$ values of the surface Arabian Sea, atmospheric $\Delta^{14}\text{C}$ values used in this model, and a steady state difference between atmosphere and ocean of about 45%. This yields a value of $F_{12} = 12 \text{ mol m}^{-2} \text{ yr}^{-1}$ for the northern Arabian Sea. This value is similar to that derived for GKh using our model.

Broecker *et al* (1985) have established a relation between the wind speed and the CO₂ invasion flux. Considering a mean annual wind speed in the northern Arabian Sea region as $\sim 6\text{m/s}$ (Esbensen and Kushnir 1981) the CO₂ invasion flux obtained is $13 \text{ mol m}^{-2} \text{ yr}^{-1}$. Our calculation is in good agreement with this estimate. The uncertainties in F_{12} and w are $\sim 10\%$.

5. Conclusions

The air-sea CO₂ exchange rate and the advective transport of water to the Gulf of Kutch were determined by modelling the coral $\Delta^{14}\text{C}$ time series. The estimated CO₂ exchange rate was found to be $11\text{--}12 \text{ mol m}^{-2} \text{ yr}^{-1}$ which was in good agreement with that obtained from other evidences. The model results suggest that it is not possible to reproduce the radiocarbon time series in the Gulf of Kutch coral without considering the advective supply of carbon and radiocarbon to the gulf from the Arabian Sea. Further the model also predicts that with the typical CO₂ exchange rate of $8\text{--}25 \text{ mol m}^{-2} \text{ yr}^{-1}$ the $\Delta^{14}\text{C}$ in corals would be much higher than commonly observed if not diluted by carbon from other sources. The model derived advective velocity ($\sim 28 \text{ m yr}^{-1}$) between the gulf and the Arabian Sea is too low compared to the surface current velocity in this region. This could be explained in terms of fluxes of carbon and radiocarbon to the gulf to match the observed coral $\Delta^{14}\text{C}$ time series.

Acknowledgements

We thank Mr M Bhaskaran and Mr Y B Raval for their help in collecting the coral sample and Dr G B Pant for providing the tree ring sample. Help provided by Mr Ravi Bhushan and Mr J P Bhavsar is gratefully acknowledged. We are thankful to Dr S R Shetye, E M Druffel and Prof D Lal for their comments. Partial financial support was made available by a grant given by Department of Ocean Development to the Physical Research Laboratory, for setting up a radio carbon laboratory; and INSA and ID-GBP (modelling) to one of us (RR).

References

- Bacastow R B and Keeling C D 1981 Atmospheric carbon dioxide concentration and the observed airborne fraction; In: *Carbon Cycle Modelling* (ed.) B Bolin (John Wiley) pp. 103–112
- Bhushan R, Chakraborty S and Krishnaswami S 1994 Physical Research Laboratory (Chemistry) Radiocarbon date list CH-1; *Radiocarbon* (submitted)
- Broecker W S, Peng T H, Ostlund G and Stuiver M 1985 The distribution of bomb radiocarbon in the ocean; *J. Geophys. Res.* **90** 6953–6970
- Cain W F and Suess H E 1976 Carbon 14, in tree rings; *J. Geophys. Res.* **81** 3688–3694
- Cember R 1989 Bomb radiocarbon in the Red sea: A medium scale gas exchange experiment; *J. Geophys. Res.* **94** 2111–2123

- Chakraborty S 1993 Environmental significance of isotopic and trace elemental variations in banded corals; PhD thesis, M.S. University of Baroda, Baroda, 119 + 22 p
- Chakraborty S and Ramesh R 1994 Stable isotope variations in a coral from the Gulf of Kutch: environmental implications; In: *Global Change Studies, scientific results from ISRO Geosphere Biosphere Programme*, ISRO-GBP-SR-42-94, 245–255
- Dickson A G 1992 JGOFS: measuring CO₂ in the ocean; *Eos* **73** 546
- Druffel E M and Linick T W 1978 Radiocarbon in annual coral rings of Florida; *Geophys. Res. Lett.* **5** 913–916
- Druffel E M 1980 Radiocarbon in annual coral rings of Florida and Belize; *Radiocarbon* **22** 363–371
- Druffel E M 1981 Radiocarbon in annual coral rings from the eastern tropical Pacific Ocean; *Geophys. Res. Lett.* **8** 59–62
- Druffel E M and Suess H E 1983 On the radiocarbon record in banded corals: Exchange parameters and net transport of ¹⁴CO₂ between atmosphere and ocean; *J. Geophys. Res.* **88** 1271–1280
- Esbensen S K and Kushnir Y 1981 The heat budget of the global ocean: An Atlas based on estimates from surface marine observations; *Rep 29*, Clim Res Inst., Oreg State Univ, Corvallis
- Gupta S K and Polach H A 1985 Radiocarbon dating practices at ANU, Radiocarbon Laboratory handbook, Research School of Pacific Studies, Australian National University, Canberra, 173 p.
- Kusumgar S 1965 Study of the geophysical processes based on C¹⁴/C¹² ratios in the atmosphere; M.Sc thesis, University of Bombay, Bombay, 55 p.
- Moore W S and Krishnaswami S 1974 Correlation of X-radiography revealed banding in corals with radiometric growth rates; *Proc. 2nd Int. Symp. on Coral Reefs* **2** 269–276
- Noakes J E, Kim S and Stipp J J 1965 Chemical and counting advances in liquid scintillation age dating; In: *Proc. Int. Conf. on ¹⁴C and Tritium Dating* (eds.) R M Chatters and E A Olson, NBS, Washington, DC, USAEC 650–652, 68–92
- Nozaki Y, Rye D M, Turekian K K and Dodge R E 1978 A 200 year record of carbon-13 and carbon-14 variations in a Bermuda coral; *Geophys. Res. Lett.* **5** 825–828
- Nydal R and Lövseth K 1983 Tracing bomb ¹⁴C in the atmosphere 1962–1980; *J. Geophys. Res.* **88** 3621–3642
- Östlund H G, Craig H, Broecker W S and Spencer D 1987 *GEOSECS Atlantic Pacific and Indian Ocean Expedition 7* NSF Washington DC 200 pp
- Pant G B and Borgaonkar H P 1983 Growth rings of teak trees and regional climatology (an ecological study of Thane region); In: *Environmental Management* (eds.) L R Singh *et al* (Allahabad India: Allahabad Geological Society Univ.)
- Ramesh R, Bhattacharya S K and Pant G B 1989 Climatic significance of δD variations in a tropical tree species from India; *Nature (London)* **337** 149–150
- Siegenthaler U and Sarmiento J L 1993 Atmospheric carbon dioxide and the ocean; *Nature (London)* **365** 119–125
- Srivastava P S and John V C 1977 Current regime in the Gulf of Kutch; *Indian J. Mar. Sci.* **6** 39–48
- Stuiver M 1980 ¹⁴C distribution in the Atlantic ocean; *J. Geophys. Res.* **85** 2771–2778
- Stuiver M and Polach H A 1977 Discussion, reporting of ¹⁴C data; *Radiocarbon* **19** 355–363
- Stuiver M and Quay P D 1981 Atmospheric ¹⁴C changes resulting from fossil fuel CO₂ release and cosmic ray flux variability; *Earth Planet. Sci. Lett.* **53** 349–362
- Stuiver M and Östlund H G 1983 GEOSECS Indian Ocean and Mediterranean radiocarbon; *Radiocarbon* **25** 1–29
- Sundquist E T 1985 Geological perspective on carbon dioxide and the carbon cycle; In: *The carbon and atmospheric CO₂: natural variations Archean to Present* (ed.) E T Sundquist and W S Broecker (Washington DC: Geophysical Monograph 32, AGU)
- Toggweiler J R, Dixon K and Broecker W S 1991 The Peru upwelling and the ventilation of the South Pacific thermocline; *J. Geophys. Res.* **96** 20467–20497