Mineralogical and chemical characteristics of newer dolerite dyke around Keonjhar, Orissa: Implication for hydrothermal activity in subduction zone setting

PIYALI SENGUPTA*, ARIJIT RAY and SAYANTANI PRAMANIK

Department of Geology, Presidency University, 86/1 College Street, Kolkata 700 073, India.

**Corresponding author. e-mail: piysen@yahoo.co.in*

The newer dolerite dykes around Keonjhar within the Singbhum Granite occur in NE–SW, NW–SE and NNE–SSW trends. The mafic dykes of the present study exhibit several mineralogical changes like clouding of plagioclase feldspars, bastitisation of orthopyroxene, and development of fibrous amphibole (tremolite–actinolite) from clinopyroxene, which are all considered products of hydrothermal alterations. This alteration involves addition and subtraction of certain elements. Graphical analyses with alteration index and elemental abundances show that elements like Rb, Ba, Th, La and K have been added during the alteration process, whereas elements like Sc, Cr, Co, Ni, Si, Al, Fe, Mg and Ca have been removed. It is observed that in spite of such chemical alteration, correlation between major and trace elements, characteristic of petrogenetic process, is still preserved. This might reflect systematic alteration (addition or subtraction) of elements without disturbing the original element to element correlation. It has also been established by earlier workers that the evolution of newer dolerite had occurred in an arc-back arc setting which may also be true for newer dolerites of the present study. This is evident from plots of pyroxene composition and whole rock composition of newer dolerite samples in different tectonic discrimination diagrams using immobile elements. The newer dolerite dykes of the Keonjhar area may thus be considered to represent an example of hydrothermal activity on mafic rocks in an arc setting.

1. Introduction

Proterozoic mafic dyke swarms are fairly common in the shield areas all over the world. In India, Proterozoic mafic dykes are widespread in the cratonic areas of Bundelkhand, Dharwar, Singbhum (Saha 1994; Halls *et al.* 2007; Pati *et al.* 2008). These dykes are disposed in NE–SW, NW–SE direction and they have tholeiitic composition. Many of them bear the signature of low grade metamorphic alteration (Mallikharjuna Rao *et al.* 2005; Bose 2008).

The mafic dyke swarm, popularly known as the newer dolerite suite, traverses the Precambrian

Singbhum granitic complex in the districts of Singbhum, Keonjhar and Mayurbhanj in eastern India (Dunn 1929; Jones 1934; Krishnan 1936; Dunn and Dey 1942; Saha 1948, 1952; Saha *et al.* 1972, 1973). The reticulate dyke system defines two broad populations trending NW–SE and NE–SW in the Singbhum craton (Dunn 1929; Dunn and Dey 1942; Saha *et al.* 1973; Bose 2008). These orientations of the dykes apparently imply a principal axis of stress from the north resulting in brittle deformation of the rigid Singbhum pluton (Dunn and Dey 1942). The dykes in the northern domain of the Singbhum pluton appear to be emplaced along planes of conjugate shear fractures

Keywords. Newer dolerite; hydrothermal activity; alteration; subduction.

(Dunn and Dey 1942; Saha 1994; Mukhopadhyay 2001). It has been proposed that hybrid fracturing of the rigid granitic mass evolved through both extension and shear embracing nucleation, propagation and coalescence at varying relative rates (Mandal *et al.* 2006). Few radiometric age data are available for these mafic dykes using K–Ar, Rb–Sr, Sm–Nd system of dating (Saha 1994; Mallick and Sarkar 1994) which broadly assign a Proterozoic status to these mafic dykes. It appears from previous works that the newer dolerites were evolved in arc-back arc setting (Bose 2008; Mir *et al.* 2010, 2013).

The present research comprises field, petrographical and geochemical study of the newer dolerite dykes around Keonjhar of Orissa situated in the western marginal parts of the Singhbhum

craton respectively (figure 1). The newer dolerite dykes occur as distinct long ridges, hillocks and also as stony wastes dominantly showing a NE–SW, NW–SE and NNE–SSW trend within the Singhbhum Granite around Keonjhar (figure 1). In all the exposures, rocks are very hard, compact and massive. The widths of the dykes vary from a few meters to over 1 km. The mafic dykes of the present study show the effect of hydrothermal alteration which imparts a green colour to these rocks causing significant mineralogical changes.

The aim of the present research is to characterize and document the mineralogical changes of the newer dolerite dykes due to hydrothermal activity and to understand the mechanism of such alteration processes which involve addition and subtraction of certain elements,

Figure 1. Geological map of Singhbhum region (modified after Saha 1994) indicating study areas denoted by solid blocks A and B around Keonjhar. The two blocks have been blown up to show occurrences of newer dolerite dykes within Singhbhum granite around Keonjhar. Important towns and villages in and around the study areas have also been marked.

especially trace elements during such hydrothermal activity.

2. General geology

The Sinbhum craton is one of the oldest cratonic nuclei of the Indian landmass (Basu *et al.* 1981; Goswami *et al.* 1995; Misra *et al.* 1999; Mandal *et al.* 2006; Mukherjee *et al.* 2008). It preserves a complex geological association involved in several episodes of magmatism, sedimentation and orogenies over a protracted period of time in the Archaean (Goswami *et al.* 1995; Misra *et al.* 1999; Mukhopadhyay 2001; Roy and Bhattacharya 2012). The Singbhum Granite batholithic complex within Singbhum craton comprises at least 12 granitic units, which are considered to have been emplaced in three successive but closely related phases: Phase I, Phase II and Phase III (Saha 1994). The Singbhum granite pluton is invaded by a group of dykes, the 'Newer Dolerites' as reported by Dunn and Dey (1942) marking the stabilisation of the craton as a whole. The chronostratigraphic succession of rocks of Singbhum craton (Misra 2006) indicate that the newer dolerite dykes are the youngest mafic magmatic unit.

3. Petrography and mineralogy

Petrographic studies were carried out with Nikon Polarizing microscope (Model no. 064333) at the Department of Geology, Presidency University, Kolkata and composition of constituent minerals were determined from diamond polished thin section with carbon coating by Electron Probe Micro Analyses (EPMA) in the laboratory of Geological Survey of India, Kolkata using CAMECA SX 100 with accelerating voltage 15 kV, 12 nA current and a beam size of 1 micron. All natural standards are used for analyses of Ca, Fe, Si, Al, Na, Mg, Cr, K, Ni, P elements. For analyses of Mn and Ti, synthetic mineral standard has been used.

Mafic dykes of the study area are represented by both fine-grained porphyritic variety and medium to coarse-grained nonporphyritic variety. Porphyritic varieties are characterized by occurrence of phenocrysts of clinopyroxene and plagioclase. Clinopyroxenes show a faint pink to green colour and occur as single prismatic grains and as clusters. Compositionally they show a wide range of variations from $Wo_{43.9} En_{44.3} Fs_{11.8}$ to $Wo_{13.9} En_{27.7} Fs_{58.4}$ (table 1) forming augite, sub-calcic augite, ferroaugite to pigeonite (figure 2). Zoning is observed in some large clinopyroxene grains commonly showing iron enrichment trend from core to rim. In few cases, the augite core of a grain is surrounded by a pigeonite rim. In

some dykes, clinopyroxenes are replaced by chlorite, tremolite and actinolitic amphibole (figure 3). In the terminology of Leake *et al.* (1997), all the studied amphiboles are of calcic type, namely ferro actinolite, actinolite, actinolitic hornblende (figure 4). Orthopyroxene occurs as a major mineral in some dykes but is mostly replaced by bastite leaving a few relict grains (figure 5) (table 1). Dykes with orthopyroxenes are sometimes associated with anorthitic plagioclase ($An_{86.7}$). These orthopyroxene bearing dykes may be classified as norite or noritic gabbro. Fresh plagioclase grains are very rare in the dolerites of the present study area. They are mostly clouded showing dark to light grey coloured dusty appearance (figure 6) and are nearly isotropic between crossed polars. Wide variation of composition (from $An_{95.3}$ to $Ab_{99.5}$; table 2) is observed in different parts of unzoned clouded plagioclase grains. Alterations like saussuritization and sericitization are also observed in some plagioclase grains. Uncoloured small grains of plagioclase show extreme albitic ($Ab_{97.4}$) compositions with some intermediate ones ($Ab_{48} An_{51.8}$) (table 2). Figure 7 shows the variable compositions of both clouded and unclouded plagioclase grains covering a wide range of compositions from albite to anorthite. The common accessory minerals are magnetite, occasionally associated with ilmenite, biotite and sphene. Interstitial quartz and high modal abundance of granophyric intergrowth are commonly developed and often constitute a significant proportion of the rock.

4. Geochemistry

In order to get an idea about petrogenetic process and subsequent alteration, chemical composition of mafic rocks appear to be important. For this purpose 37 samples of newer dolerite of the present study have been analyzed to determine major element abundances by XRF method from Wadia Institute of Himalayan Geology, Dehradun, India. Trace elements including rare earth element abundances of 13 samples were determined by inductively coupled plasma mass spectrometry techniques using Perkin Elmer, Sciex ELAN DRC II system at National Geophysical Research Institute, Hyderabad, India. International rock standards like BHVO-1, BR (major elements) and JB2 (trace and REE) were run along with the samples to check the precision and accuracy of measurement. The analyzed and certified values of BHVO-1 and JB2 are given in tables 3 and 4. The major oxides of these 37 samples show a wide range of compositional variation; SiO_2 content (46.77–56.99%), TiO_2 (0.39–2.65%), Al_2O_3 (7.48–14.04%), FeO (7.51–13.04%), Fe_2O_3

Table 1. Representative EPMA data of pyroxene grains in newer dolerite dykes of Keonjhar.

Sl. no.	1	2	3	4	5	6	7	8	9	10	11	13	14	16
Des	Cpx	Cpx	Cpx	Cpx	Cpx	Cpx core	rim	Relict opx	Pig	Cpx	Cpx	Cpx	Cpx	Relict opx
SiO ₂	49.96	51.9	50.89	51.31	49.41	50.43	51.34	54.62	51.17	49.76	52.16	50.29	49.33	54.68
TiO ₂	1.1	0.22	0.22	0.04	0.73	0.56	0.24	0.06	2.88	0.67	0.39	0.38	0.56	0.11
Al ₂ O ₃	2.79	1.62	2.94	1.89	2.04	1.8	1.29	1.14	0.82	1.7	2.08	2.18	0.93	1.49
FeO	12.35	9.65	7.01	20.02	16.5	12.37	18.45	13.21	26.12	19.63	8.6	14.57	28.62	12.1
Cr ₂ O ₃	0	0.04	0.85	0.1	0.08	0	0.13	0.4	0	0	0	0.19	0	0.5
MnO	0.23	0.24	0.27	0.18	0.4	0.21	0.43	0.21	0.29	0.42	0.21	0.27	0.51	0.2
MgO	13.9	16.49	15.36	11.41	12.83	15.93	21.19	28.15	7.04	11.25	16.68	15.91	12.05	28.08
CaO	19.17	18.55	21.2	12.68	17.55	17.77	5.84	2.34	4.89	16.29	19.25	15.07	7.63	2.37
Na ₂ O	0.33	0.19	0.35	0.29	0.28	0.31	0.12	0.02	3.62	0.23	0.22	0.24	0.11	0.01
K ₂ O	0.02	0	0	0.07	0	0	0.02	0.06	0.95	0	0.02	0	0	0.01
Number of ions on the basis of 6 oxygen														
Si	1.876	1.938	1.892	2.016	1.882	1.887	1.919	1.948	2.04	1.919	1.928	1.897	1.943	1.959
Al	0.053	0.062	0.128	0.087	0.819	0.079	0.057	0.048	0.038	0.077	0.098	0.097	0.043	0.0625
Fe ³⁺	0.087	0.052	0.074	0	0.122	0.136	0.099	0.045	0.038	0.063	0.048	0.099	0.046	0
Ti	0.031	0.006	0.006	0.001	0.021	0.016	0.007	0.002	0.086	0.019	0.011	0.011	0.017	0.003
Fe ²⁺	0.3	0.249	0.144	0.658	0.404	0.25	0.478	0.349	0.833	0.571	0.217	0.361	0.897	0.363
Cr	0	0.001	0.025	0.003	0.002	0	0.004	0.011	0	0	0	0.006	0	0.014
Mg	0.778	0.918	0.851	0.668	0.728	0.889	1.181	1.497	0.418	0.647	0.919	0.895	0.707	1.5
Mn	0.007	0.008	0.009	0.006	0.013	0.007	0.014	0.006	0.01	0.014	0.007	0.009	0.017	0.006
Ca	0.771	0.742	0.845	0.534	0.716	0.713	0.234	0.089	0.209	0.673	0.762	0.609	0.322	0.091
Na	0.024	0.014	0.025	0.022	0.021	0.022	0.009	0.001	0.28	0.017	0.016	0.018	0.008	0.001
K	0.001	0	0	0.004	0	0	0.001	0.003	0.048	0	0.001	0	0	0
Wo	39.66	37.69	43.93	28.61	36.11	35.71	11.67	4.50	13.85	34.23	39.02	30.89	16.19	4.64
En	40.02	46.62	44.29	35.82	36.73	44.55	58.89	75.35	27.75	32.89	47.04	45.37	35.57	76.54
Fs	20.32	15.69	11.78	35.58	27.15	19.74	29.44	20.15	58.40	32.89	13.94	23.75	48.25	18.81

Des: description, Cpx: Clinopyroxene, Opx: orthopyroxene, Pig: pigeonite.

Figure 2. Representative compositions of pyroxenes in newer dolerite dykes of Keonjhar area.

(1.46–2.53%), MgO (3.19–14.07%), MnO (0.14–0.23%), Na₂O (1.11–4.34%), K₂O (0.43–1.72%), and P₂O₅ (0.06–0.58%). The dolerites are basalt to basaltic andesite in composition (figure 8). A prominent iron-enrichment trend is observed in the AFM diagram (figure 9).

Figure 3. Photomicrograph showing replacement of clinopyroxene grain by tremolite–actinolite needles between crossed polars.

In a majority of bivariate diagrams with mg number against TiO₂, FeO, Na₂O + K₂O, P₂O₅ (figure 10), a distinct linear differentiation trend is observed supporting the co-genetic nature of the dolerite dykes. Inter element correlation and variation is an important aspect to portray magmatic differentiation mechanism. It is observed

Figure 4. Representative compositions of amphiboles in newer dolerite dykes of Keonjhar area.

Figure 5. Photomicrograph showing bastitised orthopyroxene grain with some relict parts in plane polarized light.

that TiO_2 , FeO , $(\text{Na}_2\text{O} + \text{K}_2\text{O})$, P_2O_5 show a good negative correlation with mg number whereas oxides such as TiO_2 , FeO , P_2O_5 show a positive correlation. $\text{K}_2\text{O}/\text{TiO}_2$ ratio shows a positive correlation with mg number. Absence of good correlation is observed in CaO vs. MgO and SiO_2 vs. MgO .

Binary variation diagrams constructed with some important trace elements like Ba, Y, Ni, Th, Zr, Nb, La, etc., clearly demonstrate the co-genetic nature of the dolerite dykes (figure 11). In all the plots, the different dykes show a distinct trend of evolution except Rb–Y. Rb is highly sensitive to alteration and as a result does not show any correlation with Y. A good positive correlation is observed between Ba–Y, Th–Y, Zr–Nb, La–Zr while a negative correlation is found in case of Ni–Zr.

REE abundances of mafic dykes of the present study are plotted in a chondrite-normalized diagram (figure 12). The plots of the newer dolerite

suite show a fractionated REE pattern with LREE enrichment. All the dykes show a similar pattern but their overall abundances vary from 50 to 100 times higher compared to chondrite. Similar REE pattern of the dolerite dykes also supports their co-genetic nature.

5. Signatures of alteration from petrographic and geochemical studies

5.1 Petrographical features

We will discuss some of the typical alteration features of some minerals which may throw light on the possible alteration processes which operated during the formation of these rocks.

5.1.1 Clouding in feldspar

Detailed observation of thin sections revealed that plagioclase feldspar laths under plane polarized light, commonly show grey to black turbid patchy discolouration or clouding (figure 6). This clouding occurs due to presence of ultramicroscopic dust like inclusions (MacGregor 1931; Poldervanrt and Gilkey 1954; Pichamuthu 1959; Zhang 1988; Halls *et al.* 1994). Previous works on clouded plagioclase (Knopf and Jonas 1929) show that these dusty inclusions are mostly magnetite, i.e., much of the iron in clouded feldspar is in the divalent state.

Spatial distribution of clouding is variable among different grains in a specimen and among different specimens in the study area. But an overall account shows that many of the rocks show intense clouding in plagioclase while others show medium to weak clouding effects. The optical density varies across a grain. The outer margin is sometimes clearer than the interior because it contains no inclusions. Feldspar clouding is easily distinguishable from feldspar saussuritization. Feldspar clouding forms a brown to dark grey stain in plane polarized light, formed by the presence of sub-microscopic magnetite. Between crossed polars these clouded grains appear opaque. Saussurite forms a dusty grey amalgamation of visible minerals in plane polarized light, with a speckled appearance between cross nicols.

Previous observation revealed that clouded feldspar is very common in mafic dykes that transect Proterozoic shield areas such as satellite dykes of the Great Dyke in Zimbabwe (Robertson and Van Breeman 1970), Matachewan dykes of the Canadian shield (Halls and Palmer 1990), dykes on Wyoming shield, USA (Armbrustmacher and Banks 1974), Dharwar craton in India (Pichamuthu 1959) and many others. It has been

Figure 6. Photomicrograph showing clouded plagioclase grain in plane polarized light.

shown that cloudiness is a product of thermal metamorphism caused by regional heating or by the heat of neighbouring igneous magma (MacGregor 1931). We prefer the regional heating model of thermal metamorphism as the possible cause for feldspar clouding in our study area as there is no evidence of intrusive rocks in the vicinity. It is also evident from petrographic studies that the dolerite dykes of the present study area have suffered hydrothermal alteration.

Among many hypotheses, we consider the model of Poldervanrt and Gilkey (1954) suitable to explain the clouding of feldspar of the study area. According to the hypothesis, weak clouding may result from exsolution of iron incorporated into the feldspar during growth, but intense clouding is produced by diffusion of material into the feldspar after growth by means of channels produced by unmixing of plagioclase into regions of albite and anorthite. The clouded feldspar of the newer dolerites around Keonjhar shows extreme compositional variation ($Ab_{4.7}$ to $Ab_{99.5}$; table 2) within a single grain. This variation is considered the result of unmixing of plagioclase of intermediate composition. Such unmixing might have occurred when the plagioclase was held at high temperatures for a long time in the presence of water. Such conditions occur during metamorphism of basic rocks of high iron content and from entrance of Fe-rich solutions along the zones of unmixing precipitating submicroscopic grains of iron mineral (Poldervanrt and Gilkey 1954).

5.1.2 Large bastitised orthopyroxene grain

Some of the dykes contain numerous large bastitised orthopyroxene (figure 5) grains leaving relict

Table 2. Representative EPMA data of plagioclase grains in newer dolerite dykes of Keonjhar.

Sl. no.	1	2	3	4	5	6	7	8
Des.	Plag	Plag	Plag	Plag	Relict plag	Plag core	Cloudy plag	Cloudy plag
SiO ₂	60.9	64.56	58.87	56.45	58.37	65.85	68.36	48.87
TiO ₂	0.11	0.07	0.04	0.16	0	0.12	0	0
Al ₂ O ₃	23.11	20.89	23.75	25.37	25.07	17.73	19.29	25.57
FeO	0.53	0.26	0.56	0.59	1.06	1.7	0.14	4.13
MnO	0.05	0.04	0	0.03	0	0.04	0	0.05
MgO	0.01	0	0.05	0.03	0.39	1.08	0.04	0.36
CaO	5.14	2.59	7.35	8.92	5.16	1.26	0.08	13.04
Na ₂ O	8.1	9.64	7	6.12	6.99	10.57	11.61	4.07
Cr ₂ O ₃	0	0.03	0.06	0	0.02	0	0.05	0
K ₂ O	0.78	0.62	0.73	0.55	2.11	0.23	0.08	1.72
Cations on the basis of 16 oxygen								
Si	5.529	5.792	5.423	5.231	5.313	6.074	6.005	4.95
Al	2.471	2.207	2.577	2.769	2.687	1.926	1.995	3.05
Ti	0	0.01	0	0	0	0.008	0	0
Fe ²	0.01	0.056	0.065	0.022	0.106	0.04	1.008	0.07
Mg	0.004	0.008	0.006	0.001	0.18	0.001	0.004	0.016
Ca	0.047	1.02	1.43	0.479	1.24	0.5	2.917	0
Na	1.923	1.029	0.668	1.438	1.15	1.426	0.145	1.88
K	0.005	0.051	0.018	0.041	0.004	0.09	0	0.009
Ab	97.4	49.0	31.6	73.4	48.0	70.7	4.7	99.5
An	2.4	48.6	67.5	24.5	51.8	24.8	95.3	0.0
Or	0.2	2.4	0.9	2.1	0.2	4.5	0.0	0.5

Plag: Plagioclase feldspar, Des: description.

Figure 7. Composition of both clouded and unclouded plagioclase grains in Ab–An–Or diagram.

parts of unaltered orthopyroxenes. The composition of these orthopyroxene grains varies from $Wo_{4.3} En_{72.5} Fs_{23.2}$ to $Wo_{4.2} En_{80.4} Fs_{15.4}$. The grains are associated with rare augite and plagioclase grains. Bose (2008) has reported presence of partly bastitised orthopyroxene in a norite sample within newer dolerite dykes from Parsola but due to the absence of an interlocking boundary of orthopyroxene grains, Bose (2008) suggested that the rock could be a cumulate. In our study, as the orthopyroxene grains in some of the dykes show interlocking texture with associated plagioclase and clinopyroxenes, it is more likely that they might have been formed by the contamination of the upcoming magma with the granitic country rock (Deer *et al.* 1997). Similar orthopyroxene bearing dolerite dyke has been reported from Chaibasa area of Singhbhum region (Sengupta and Ray 2012) where co-existence of bastitised orthopyroxenes, anorthitic plagioclase and clinopyroxene has been explained as product of contamination of granitic country rock and mafic melt. In the present study, coexistence of orthopyroxene ($Wo_{3.5} En_{82} Fs_{14.5}$) and anorthitic plagioclase ($An_{93.1}$) possibly suggest similar assimilation and transformation of clinopyroxene to orthopyroxene coupled with migration of Ca ion into plagioclase during petrogenesis. Formation of bastite in orthopyroxene indicates hydrothermal activity during greenschist facies metamorphism (Johnson *et al.* 2004; Plumper *et al.* 2012).

5.1.3 Actinolite-tremolite assemblage

In many dykes, fine needles of tremolite and actinolite are observed along marginal parts of

clinopyroxene grains (figure 3). In some cases, the clinopyroxene has been altered to fine tremolite and actinolite needles. The presence of a few discrete primary amphibole grains and frequent hydrothermal alteration of clinopyroxene to chlorite and actinolitic amphibole suggest influence of water during late magmatic stage and metamorphic alteration (Devaraju *et al.* 2008). According to Devaraju *et al.* (2008), the mafic dyke of Western Dharwar craton has widespread evidence of deuteritic alteration. Minerals such as uraltite, chlorite, epidote are considered products of the interaction of deuteritic liquid with ferromagnesian minerals, pyroxene. The composition of deuteritic liquids has been assessed by them as one with high silica, alumina, alkali, Ti, P and REE. The actinolite-tremolite assemblage of newer dolerite dykes around Keonjhar may be considered products of interaction of late magmatic hydrous fluid and early magmatic clinopyroxene where the reaction involves migration of volatile phases along with dissolved silica, alkali, REE from hydrothermal fluid to the clinopyroxene grains. Similar alterations of clinopyroxene grains to actinolite, tremolite needles have been recorded from low grade metamorphosed mafic volcanic rocks of the Mira terrane of Avalonia, southeastern Cape Breton Island, Nova Scotia (Mc Mullin *et al.* 2010).

6. Changes in chemical composition of the mafic dykes due to alteration

Ocean floor rocks show a range of consistent mineralogical alterations during low temperature hydrothermal processes (Frey *et al.* 1974; Mottl 1983). Newer dolerite rocks of the present area also exhibit several mineralogical changes like alteration of clinopyroxenes to tremolite–actinolite needles, bastitization of orthopyroxenes, clouding in feldspars which have been discussed in detail in previous sections. It is obvious that changes of these mineralogical compositions should also be reflected in the chemical composition of the rock; i.e., the altered rocks would show some changes in major and trace element abundances from their fresh counterparts.

Several studies on altered mafic rocks have shown that most, if not all, elements can be added or subtracted from a rock during low temperature ocean floor metamorphism (Mottl and Holland 1978; Seyfried *et al.* 1988) and the extent of the effect of alteration varies from one member to the other within a group although they show overall similar mineralogical changes. These rocks are particularly effective for such a geochemical study as they show a range of mineralogical

Table 3. Major (in wt%) element data of the newer dolerite dykes within Singbhum Granitoid complex near Keonjhar, Orissa.

Sl. no. Sample	1 P29	2 P44	3 P49	4 P53	5 P57	6 P61	7 P62	8 P43A	9 P30	10 P31	11 P32A
SiO ₂	53.75	53.99	47.55	46.77	51.96	53.9	50.71	54.7	50.59	51.18	52.2
TiO ₂	0.55	0.56	0.43	0.39	0.71	0.51	1.06	0.47	1.69	1.18	2.49
Al ₂ O ₃	10.94	11.01	11.54	12.15	7.62	12.63	10.76	9.41	12.65	13.38	12.56
FeO	8.24	8.41	9.04	9.33	10.4	7.51	9.99	9.01	11.97	11.24	12.39
Fe ₂ O ₃	1.6	1.63	1.76	1.81	2.02	1.46	1.94	1.75	2.32	2.18	2.4
MgO	12.51	11.28	13.89	14.07	11.43	8.23	10.79	9.65	4.93	4.82	3.43
CaO	6.89	7.6	9.87	9.09	8.93	9.75	7.33	8.35	8.39	9.23	7.35
MnO	0.16	0.16	0.16	0.17	0.18	0.14	0.16	0.17	0.2	0.21	0.21
Na ₂ O	1.11	1.56	1.33	1.47	2.01	1.75	2.52	2	2.51	2.7	2.84
K ₂ O	1.72	1.04	0.46	0.67	0.62	1.16	1.32	1.14	1.13	0.89	1.24
P ₂ O ₅	0.08	0.09	0.08	0.08	0.13	0.07	0.15	0.06	0.34	0.17	0.54
LOI	3.01	2.03	3	3.26	1.84	2.3	2	2.14	1.14	1.77	1.39
Total	100.56	99.36	99.11	99.26	97.85	99.41	98.73	98.85	97.86	98.95	99.04
mg no.	64	62	65	64	57	57	56	56	33	34	25
Sl. no. Sample	12 P33	13 P34	14 P36	15 P37	16 P38	17 P39	18 P41	19 P46	20 P48	21 P54	22 P55
SiO ₂	51.91	52.09	56.55	48.88	48.88	50.33	51.46	56.36	50.81	52.29	51.05
TiO ₂	2.63	2.65	1.08	1.64	1.91	1.65	1.97	1.05	1.36	1.08	1.13
Al ₂ O ₃	12.56	12.64	11.56	12.93	12.76	12.8	13.26	11.48	13.08	13.78	13.16
FeO	12.6	12.6	10.92	11.74	12.54	11.67	10.82	10.95	10.69	10.76	11.22
Fe ₂ O ₃	2.45	2.45	2.12	2.28	2.43	2.27	2.1	2.13	2.08	2.09	2.18
MgO	3.19	3.3	3.56	5.28	4.53	4.46	5.33	3.67	5.26	5.22	5.13
CaO	7.51	7.48	7.57	8.87	8.79	8.3	7.46	7.58	8.28	9.31	9.46
MnO	0.22	0.22	0.21	0.2	0.22	0.2	0.17	0.21	0.18	0.2	0.21
Na ₂ O	2.75	2.77	2.85	2.82	2.53	2.5	3.66	2.88	2.69	2.71	2.56
K ₂ O	1.21	1.21	1.34	0.73	0.81	1.32	1.17	1.3	1.3	0.83	0.78
P ₂ O ₅	0.57	0.58	0.26	0.32	0.39	0.39	0.17	0.27	0.33	0.17	0.15
LOI	1.54	1.48	1.4	1.6	1.55	1.93	2.09	1.55	1.93	1.31	1.52
Total	99.14	99.47	99.42	97.29	97.34	97.82	99.66	99.43	97.99	99.75	98.55
mg no.	23	24	28	35	30	31	37	29	37	37	35

Sl. no. Sample	23 P58	24 P59	25 P60	26 P63	27 P52	28 P50	29 P28A	30 P32B	31 P35	32 P40
SiO ₂	53.33	52.98	47.7	49.95	56.99	53.24	53.2	53.01	49.41	51.82
TiO ₂	2.17	2.16	2.11	1.59	1.02	1.62	1	0.92	1.39	0.93
Al ₂ O ₃	13.12	13.08	12.09	12.89	12.94	10.21	11.88	10.92	14.04	12.86
FeO	10.88	10.82	13.04	11.69	8.54	10.69	9.65	9.75	9.44	9.36
Fe ₂ O ₃	2.11	2.1	2.53	2.27	1.66	2.08	1.87	1.89	1.83	1.82
MgO	4.38	4.45	5.88	4.97	4.58	5.89	6.7	9.58	7.13	7.25
CaO	6.85	6.82	8.49	8.68	6.77	6.02	8.59	6.51	9.44	9.37
MnO	0.17	0.17	0.21	0.19	0.15	0.18	0.17	0.15	0.14	0.16
Na ₂ O	3.69	3.65	2.73	2.42	2.97	4.34	2.5	1.88	2.76	2.46
K ₂ O	1.45	1.45	0.72	0.98	1.54	1.68	0.79	1.46	1.22	0.82
P ₂ O ₅	0.21	0.21	0.43	0.33	0.38	0.16	0.14	0.11	0.08	0.16
LOI	1.7	1.72	1.96	1.72	1.6	1.19	2.1	2.26	2.08	2.19
Total	100.06	99.61	97.89	97.68	99.14	97.3	98.59	98.44	98.96	99.2
mg no.	32	33	35	34	39	40	45	54	47	48

Sl. no. Sample	33 P42	34 P45	35 P56	36 P65D	37 P47	38 P51	BHVO-1 analyzed	BHVO-1 Govindaraju (1994)
SiO ₂	52.47	51.57	50.07	50.75	54.43	55.24	49.08	49.94
TiO ₂	1.43	0.83	0.73	0.86	0.65	0.84	2.72	2.71
Al ₂ O ₃	12.38	11.41	9.11	11.91	7.48	12.01	13.89	13.8
FeO	9.62	9.57	9.95	9.46	11.57	9.46	9.29	9.36
Fe ₂ O ₃	1.87	1.86	1.93	1.84	2.25	1.84	1.8	1.82
MgO	7.1	8.82	9.99	8.98	7	5.73	6.98	7.23
CaO	7.64	9.09	8.37	10.03	8.82	7.9	11.2	11.4
MnO	0.16	0.17	0.17	0.17	0.23	0.18	0.16	0.17
Na ₂ O	3.07	2.19	2.64	2.09	3.11	2.69	2.37	2.26
K ₂ O	1.49	0.75	0.77	0.43	0.84	1.06	0.5	0.52
P ₂ O ₅	0.14	0.11	0.13	0.11	0.06	0.15	0.25	0.27
LOI	2.1	2.18	5.23	1.55	1.75	2.06		
Total	99.47	98.55	99.09	98.18	98.19	99.16		
mg no.	47	52	54	53	42	42		

Table 4. Trace element and REE data of the newer dolerite dykes within Singbhum Granitoid complex near Keonjhar, Orissa.

Sl. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	JB2 analysed	JB2 Govindaraju (1994)
Sp no.	P49	P53	P44	P61	P29	P 43A	P57	P40	P32B	P34	P58	P50	P37		
Sc	40.0	37.0	29.0	32.0	28.0	39.9	27.0	30.0	30.5	30.6	27.6	26.7	35.0	54.336	54.4
V	158.0	198.4	193.0	168.0	202.0	213.0	223.0	246.0	201.6	252.0	321.4	183.9	314.0	581.447	578
Cr	1077.0	1410.5	917.0	581.0	981.0	1194.3	1337.0	472.0	1193.3	276.1	86.8	538.1	270.0	27.456	27.40
Co	83.0	89.7	86.0	83.0	82.0	91.0	95.0	75.0	87.3	76.7	88.9	87.9	86.0	39.868	39.80
Ni	471.0	509.6	234.0	128.0	247.0	132.5	540.0	146.0	390.8	87.5	49.2	272.1	107.0	14.155	14.20
Ga	10.8	11.2	15.0	15.0	14.1	19.6	12.7	15.7	17.0	17.4	22.0	15.2	18.2	17.076	17.00
Rb	19.0	24.7	50.0	53.0	68.0	46.0	28.0	27.0	73.4	38.3	42.9	68.3	26.0	6.192	6.20
Sr	99.0	133.6	111.0	137.0	191.0	116.6	233.0	227.0	205.7	255.2	257.5	526.0	159.0	178.115	178.00
Y	18.0	21.7	25.0	32.0	27.0	36.3	19.0	21.0	28.1	22.0	36.2	27.8	40.0	25.004	24.00
Nb	3.4	4.4	5.8	12.7	6.7	13.2	5.5	5.2	11.4	8.2	10.5	12.1	9.6	0.809	0.80
Zr	40.0	65.0	95.0	211.0	96.0	170.0	92.0	71.0	160.0	140.0	180.4	200.2	171.0	51.25	51.40
Ba	81.0	116.2	196.0	236.0	247.0	258.6	122.0	172.0	202.1	163.5	246.7	141.3	223.0	208.98	208.00
Th	0.9	1.8	5.0	7.1	5.9	5.9	2.1	1.7	6.2	3.1	5.3	5.5	2.0	0.304	0.33
La	5.0	6.2	15.6	29.4	15.8	27.7	9.2	9.9	25.1	11.0	19.5	15.0	16.1	2.397	2.37
Ce	9.9	12.5	31.1	56.6	31.9	61.0	18.1	21.2	55.0	25.6	42.1	31.6	36.8	6.833	6.77
Nd	1.2	5.6	3.6	6.6	3.7	25.7	2.2	2.7	26.3	11.7	22.2	14.1	4.9	6.80	6.70
Pr	4.5	1.6	13.0	23.2	13.2	7.1	8.3	10.6	7.1	3.1	5.6	4.0	20.1	0.945	0.96
Sm	1.0	1.3	2.8	4.8	2.8	6.0	1.8	2.6	5.7	2.8	5.7	3.5	5.1	2.256	2.25
Eu	0.4	0.5	0.8	1.0	0.7	1.5	0.6	0.9	1.7	0.9	1.8	0.8	1.6	0.897	0.86
Gd	1.4	1.7	3.1	5.3	3.1	6.9	2.0	3.0	5.6	3.2	6.0	4.1	5.9	3.476	3.28
Tb	0.3	0.3	0.5	0.9	0.5	1.1	0.3	0.5	0.8	0.5	0.9	0.7	1.0	0.638	0.62
Dy	1.9	2.2	3.0	5.3	2.9	6.5	1.8	3.0	4.0	2.9	5.0	4.1	6.3	3.748	3.66
Ho	0.5	0.5	0.6	1.2	0.6	1.4	0.4	0.7	0.8	0.6	1.0	0.9	1.4	0.831	0.81
Er	1.5	1.6	1.7	3.2	1.7	3.8	1.1	1.7	1.9	1.6	2.4	2.4	3.7	2.561	2.63
Tm	0.3	0.3	0.3	0.5	0.3	0.6	0.2	0.3	0.3	0.3	0.3	0.4	0.6	0.455	0.45
Yb	1.8	2.0	1.8	3.3	1.8	4.0	1.0	1.7	1.7	1.7	2.2	2.4	3.9	2.463	2.51
Lu	0.3	0.3	0.3	0.5	0.3	0.6	0.2	0.3	0.2	0.3	0.3	0.4	0.6	0.392	0.39

Figure 8. Total alkali silica classification diagram (TAS) for mafic dykes (after Le Bas 2000) showing the plots of newer dolerite dykes of Keonjhar in the field of basalt and basaltic andesite.

Figure 9. Plots of the newer dolerite dykes of Keonjhar in the AFM diagram (after Irvine and Baragar 1971). The iron enrichment trend can be clearly noted.

alterations. It has been observed that mobile elements like Ba, Rb, Th, etc., and even some immobile elements as Nb and Zr can be affected under specific conditions (Hellman *et al.* 1979; Hynes 1980). Behaviour of a particular element is dependent on the element concentration in the rock, on its concentration in solutions entering the rock, and on partitioning between the fluid phase and the low-temperature secondary minerals that are stable in the rock (Frey *et al.* 1974; Mottl 1983).

In order to document quantitatively the effects of alteration on newer dolerites of the study area, we

are following the methods adopted by Greenough *et al.* (1990). In this process, to assess chemical changes caused by alteration, we have compared the least altered dolerite sample of the area with altered samples that had initial compositions similar to those of unaltered rocks. It has been observed, that in case of low temperature alteration of mafic rocks, one of the most obvious effects of alteration is highly variable Rb values within a group of rocks (Backman *et al.* 1988). The dolerite rocks of the study area show large variations in concentration of Rb (19–73.42 ppm) which cannot be accounted for only in terms of differentiation. So, alteration must have played an important role in creating large variations in Rb content among different dolerite samples. Therefore, we assume that Rb is an alteration sensitive element and the sample with lowest Rb value represents the least altered dolerite dyke of the area. Among several established immobile elements like Zr, Nb, Y, etc., we observe that Y concentration values show least variation (table 4) among the newer dolerite dykes of our study area. Therefore, to quantify the geochemical effects of alteration of the mafic dykes, we assume that Y was immobile and use it to evaluate the mobility of all other elements as described below. To assess the mobility of different elements we are using a parameter ‘alteration index’ of elements which calculates the percentage change between element/Y ratios in altered rocks as compared to least altered rock or fresh rock (i.e., sample with lowest Rb concentration).

$$\text{Alteration Index} = \frac{\text{Element/Y (altered rock)} - \text{Element/Y (least altered rock)}}{\text{Element/Y (least altered rock)}}$$

Figure 10. Bivariate diagrams using mg number against major element oxide composition (first five diagrams) and TiO_2 vs. FeO (sixth diagram) of newer dolerite dykes of Keonjhar.

Figure 13(a, b) shows plots of alteration index of different elements (Y axis) vs. absolute Rb concentrations of different altered rocks arranged in ascending order (X axis) denoting samples showing a gradual increase in degree of alteration. For a particular Rb concentration, i.e., a particular altered rock we calculate the alteration index of different elements. Positive values on the Y axis indicate that an element was added to the rock during alteration process but negative values suggest that the element was removed. The lines denote successive changes of Alteration Index of an element with gradual increase of degree of alteration from least altered sample (i.e., lowest Rb concentration) to the most altered sample (highest Rb concentration).

From figure 13(a), it is observed that elements like Rb, Ba, Th, La, K show positive values on the Y axis suggesting that these elements have been added during the alteration process whereas elements like Sc, Cr, Co, Ni, Si, Al, Fe, Mg, Ca show negative values on the Y axis (figure 13b) implying removal of these elements. The least altered

sample of newer dolerite dyke (Rb–19 ppm) occurs in the northern part of the study area and is considered to have localized in the distal part. The degree of alteration increases more or less along southward direction and the most altered sample (Rb–73.42 ppm) might represent proximal zone of alteration front (figure 1).

7. Possible paleotectonic setting

To get an idea about the tectonic setting of the dolerites we consider several mineralogical and geochemical aspects. As the dolerites show strong effects of hydrothermal alterations as discussed under petrography, it is quite reasonable to rule out the possibility of their emplacement in a continental setting where such alterations are very rare. Formation under an oceanic setting is more likely for the dolerite rocks showing albitisation, bastitisation and other features of hydrothermal alteration. Several workers have suggested a subduction zone related origin of these newer dolerite dykes

Figure 11. Binary variation diagrams constructed with some important trace element abundances of newer dolerite dykes of Keonjhar.

Figure 12. Chondrite normalized REE plot (Sun and McDonough 1989) for the newer dolerite dykes of the study area.

of Singhbhum region (Bose 2008; Mir *et al.* 2010, 2011, 2013). On the basis of detailed geochemical studies of newer dolerite dykes of Singhbhum craton, Mir *et al.* (2010) suggested that these dykes were derived from a mantle source which has been metasomatized by a subduction component (e.g., fluids derived by dehydration of the subducting slab). It has also been demonstrated that newer

dolerites show geochemical signatures similar to those of back-arc basaltic rocks (Mir *et al.* 2011, 2013). With a backdrop of this knowledge we made an attempt to understand the tectonic settings of the newer dolerite dykes of our study area in terms of a few discrimination diagrams using relatively immobile major and trace elements. In tectonic discrimination diagrams of $\text{TiO}_2\text{--MnO} \times 10\text{--P}_2\text{O}_5 \times 10$ (after Mullen 1983), Cr–Y (Pearce 1982), Ti–V (Shervais 1982) and 3Zr–Nb–Y (Meschede 1986) the plots (figure 14) of the dolerite rocks of study area mostly fall in the fields of ocean floor, oceanic ridge, and volcanic arc basalt suggesting a possible arc–back arc setting for newer dolerites of present study.

Chemical composition of constituent minerals is also important to understand tectonic setting. Partition of elements like Al, Si, Ti into clinopyroxene structure may also provide evidences for tectonic environment (Leterrier *et al.* 1982; Burns 1985; Loucks 1990; Koksai 2003). In some covariation plots with Ti, Na, Al(t), Al(IV) contents of clinopyroxenes (pfu), plots of clinopyroxene composition of newer dolerite dyke of our study area occupy the ‘Arc’ fields and the field for ‘Boninite’ (figure 15a, b, c). This possibly suggests crystallization of clinopyroxene and evolution of newer dolerite of Keonjhar area in an arc setting.

Figure 13. Graph showing the effects of increasing alterations on element concentrations in newer dolerite rocks. Y axis denotes alteration index of different elements while X axis represents absolute Rb concentrations of different altered samples.

8. Discussion

Mafic dykes are important components of the Proterozoic rock record and act as significant time markers within stabilized Archean cratons (Ernst and Buchan 2001; Bleeker 2004). In India, the Proterozoic mafic dykes are found in the major cratonic blocks of Aravalli craton, Bundelkhand craton, Dharwar craton, and Singhbhum cratons under different extensional and compressional tectonic regimes (Murthy 1987, 1995; Saha 1994; Halls *et al.* 2007; Pati *et al.* 2008). In Singhbhum region, newer dolerite dykes exhibit mineralogical

and chemical compositions which point towards emplacement in an arc setting in which hydrothermal activity and alteration are common features during emplacement (Sengupta and Ray 2012). Albitisation of plagioclase, bastitisation of orthopyroxene, and development of fibrous amphibole (tremolite–actinolite) from clinopyroxene are all considered products of hydrothermal activity in oceanic conditions (Mottl and Holland 1978). Geochemical signatures of subduction related processes in a back arc setting in newer dolerite dykes of Singhbhum pluton have been reported by several workers (Bose 2008; Mir *et al.* 2010; Sengupta and

Figure 14. Tectonic discrimination diagrams of newer dolerite dykes of Keonjhar area using major and trace element abundances.

Ray 2012). Similar features of mineralogical alteration have also been reported from other Proterozoic mafic rocks of India. The Proterozoic mafic dykes within Bundelkhand Granite Massif (BGM) show clouded feldspar, chloritized or amphibolitized clinopyroxenes as a result of hydrothermal alteration (Mallikharjuna Rao *et al.* 2005). One of the first detailed studies on the clouded feldspar in dykes on regional scale was carried out by Pichamuthu (1959) from Dharwar craton. Subsequently, it has been realized that such feldspar clouding, specially in Proterozoic mafic dyke swarms, has significant tectonic implications (e.g., Zhang and Halls 1995). Early Proterozoic mafic dykes of Kalyadi area of western Dharwar craton also exhibit tea coloured feldspar, i.e., clouded feldspar and secondary amphiboles around clinopyroxenes (Chandrasekharam *et al.* 2008).

Although reports on mineralogical alteration in Proterozoic mafic dykes of India are fairly common (Mallikharjuna Rao *et al.* 2005; Sengupta and Ray 2012), detailed studies on behaviour of major trace elements during such alterations have hardly been attempted. In the present research we have identified the alteration features of some important minerals like plagioclase, clinopyroxene, orthopyroxene, etc., and have made an attempt to find out chemical changes associated with such

mineralogical alterations of the mafic dykes. Studying the major and trace element abundances of the altered newer dolerite dykes, we have observed that elements like Rb, Ba, Th, La, K have been added whereas elements like Sc, Cr, Co, Ni, Si, Al, Fe, Mg, Ca have suffered removal from the rock during the alteration process. It has also been established that the evolution and emplacement of newer dolerite dykes had occurred in an arc-back arc setting as evidenced from different tectonic diagrams (figures 14 and 15). In general, it has been observed that alteration processes of mafic rocks in an oceanic setting yield somewhat consistent chemical changes (Frey *et al.* 1974; Mottl 1983). Greenough *et al.* (1990) have documented quantitatively, the effects of alteration on hotspot basalts from sunken oceanic island in the Indian Ocean. They have shown that the mineralogical changes involve chemical additions of K, Rb, Cs, Li, Si, Sc, Fe, etc., and removal of Ca, Mg, Mn, Ni and others. Similar geochemical studies have also been performed in ophiolitic basalts as well as ocean-floor basalts (Gills and Robinson 1985, 1988). The newer dolerite dykes of the Keonjhar area, which have suffered several mineralogical changes due to hydrothermal alteration along with addition and removal of several elements, may represent a case-study of hydrothermal activity on arc related mafic rocks.

Figure 15. Tectonic discrimination diagrams of newer dolerite dykes of Keonjhar area using partition of elements like Al, Si, Ti into clinopyroxene structure. IAT: Island Arc Tholeiite, BON: Boninite, MORB: Mid Oceanic Ridge Basalt.

Acknowledgements

The authors thank the Department of Science and Technology, New Delhi for providing research grant for the present work. They also thank the Head of the Department of Geology, Presidency University for providing facilities to carry out this work.

References

- Armbrustmacher T J and Banks N G 1974 Clouded plagioclase in metadolerite dykes, southeastern Bighorn Mountains, Wyoming; *Am. Mineral.* **59** 656–665.
- Backman J, Duncan R A et al. 1988 *Proc. ODP, Init. Repts* **115** College Station, TX (Ocean Drilling Program), doi: [10.2973/odp.proc.ir.115.1988](https://doi.org/10.2973/odp.proc.ir.115.1988).
- Basu A R, Ray S L, Saha A K and Sarkar S N 1981 Eastern Indian 3800 million year old crust and early mantle differentiation; *Science* **212** 1502–1506.
- Bleeker W 2004 Taking the pulse of planet Earth: A proposal for a new multi-disciplinary flagship project in Canadian solid Earth Sciences; *Geosci. Canada* **31** 179–190.
- Bose M K 2008 Proterozoic dykes from Singhbhum granite pluton; In: *Indian dykes* (eds) Srivastava R K, Sivaji Ch and Rao C (New Delhi: Narosa Publ.), pp. 413–445.
- Burns L E 1985 The border ranges ultramafic and mafic complex, southcentral Alaska: Cumulate fractionates of island arc volcanics; *Can. J. Earth. Sci.* **22** 1020–1038.
- Chandrasekharam D, Mellus L, Cucciniello C, Mathew B and Perini G 2008 Petrogenesis of E–W trending dykes from Kalyadi, Dharwar craton, southwestern India; In: *Indian dykes* (eds) Srivastava R K, Sivaji Ch and Rao C (New Delhi: Narosa Publ.), pp. 199–214.
- Deer W A, Howie R A and Jussman J 1997 *An Introduction to the Rock-Forming Minerals*; The English Language Book Society and Longman, London.
- Devaraju T C, Alapieti T T, Sudhakara T L and Kaukonen R J 2008 Mafic dyke swarms of volcanic arc, ocean floor, and N–MOR basalt affinity with destructive plate margin emplacement features in the northern segment of western Dharwar craton; In: *Indian dykes* (eds) Srivastava R K, Sivaji Ch and Rao C (New Delhi: Narosa Publ.), pp. 215–237.
- Dunn J A 1929 The geology of north Singhbhum including parts of Ranchi and Manbhum districts; *Geol. Surv. India Memoir* **54(2)** 166p.
- Dunn J A and Dey A K 1942 Geology and petrology of eastern Singhbhum and surrounding areas; *Geol. Surv. India Memoir* **69(2)** 261–456.
- Ernst R E and Buchan K L 2001 Large mafic magmatic events through time and links to mantle-plume heads; In: *Mantle plumes: Their identification through time* (eds) Ernst R E and Buchan K L, *Geol. Soc. Am. Spec. Paper* **352** 483–575.
- Frey F A, Bryan W B and Thompson G 1974 Atlantic Ocean floor: Geochemistry and petrology of basalts from legs 2 and 3 of the Deep Sea Drilling Project; *J. Geophys. Res.* **79** 5507–5527.
- Gills K M and Robinson P T 1985 Low-temperature alteration of the extrusive sequence, Troodos ophiolite, Cyprus; *Can. Mineral* **23** 431–441.
- Gills K M and Robinson P T 1988 Distribution of alteration zones in the upper oceanic crust; *Geology* **16** 262–266.
- Goswami J N, Mishra S, Wiedenbeck M, Ray S L and Saha A K 1995 3.55 Ga old zircon from Singhbhum–Orissa iron ore craton, eastern India; *Curr. Sci.* **69** 1008–1011.
- Greenough John D, Fryer Brian J and Robinson Paul T 1990 Geochemical effects of alteration on mafic rocks from Indian ocean site 706; *Proceedings of the Ocean Drilling Program, Scientific Results* **115** 85–92.
- Irvine T A and Baragar W R A 1971 A guide to chemical classification of common volcanic rocks; *Can. J. Earth Sci.* **8** 523–548.
- Halls H C and Palmer H C 1990 The tectonic relationship of two early Proterozoic dyke swarms to the Kapuskasing

- structural zone: A paleomagnetic study; *Can. J. Earth Sci.* **27** 78–103.
- Halls H C, Palmer H C, Bates M P and Phenney W C 1994 Constraints on the nature of the Kapuskasing structural zone from the study of Proterozoic dyke swarms; *Can. J. Earth Sci.* **31** 1182–1196.
- Halls H C, Kumar A, Srinivasan R and Hamilton M A 2007 Paleomagnetism and U–Pb geochronology of easterly trending dykes in the Dharwar craton, India: Feldspar clouding, radiating dyke swarms and the position of India at 2.37 Ga; *Precamb. Res.* **151** 47–68.
- Hellman P I, Smith R E and Henderson P 1979 The mobility of rare earth elements: Evidence and implications from selected terrains affected by burial metamorphism; *Contrib. Mineral. Petrol.* **71** 23–44.
- Hynes A 1980 Carbonatization and mobility of Ti, Y and Zr in Ascot Formation metabasalts, southeast Quebec; *Contrib. Mineral. Petrol.* **75** 79–87.
- Johnson P R, Kattan F H and Al-Saleh A M 2004 Neoproterozoic ophiolites in the arabian shield: Field relations and structure; In: *Developments in Precambrian Geology* (eds) Timothy M Kusky and Condie K C (series editor), Elsevier, **13** 129–162.
- Jones H C 1934 The iron ore deposits of Bihar and Orissa; *Geol. Surv. India Memoir* **63(2)** 167–302.
- Knopf E B and Jonas A I 1929 The geology of the crystalline rocks of Baltimore County; In: *Baltimore County, Maryland Geol. Survey*, pp. 97–199.
- Koksall F 2003 Petrology of the phlogopite-bearing ultramafic–mafic plutonic rocks within Central Anatolian Crystalline complex, Turkey; Ph.D. Thesis.
- Krishnan M S 1936 The dyke rocks of Keonjhar state, Bihar and Orissa; *Rec. Geol. Surv. India* **71** 105–120.
- Leake B E, Woolley A R, Arps C E S, Birch W D, Gilbert M C, Grice J D, Hawthorne F C, Kato A, Kisch H J, Krivovichev V G, Linthout K, Larid J, Mandarino J A, Maresch W V, Nickel E H, Rock N M S, Schmacher J C, Smith D C, Stephenson N C N, Ungaretti L, Whittaker E J W and Youzhi G 1997 Nomenclature of amphiboles: Report of the subcommittee on amphiboles of the International Mineralogical Association, Commission on New Minerals and Mineral Names; *Am. Mineral.* **82** 1019–1037.
- Le Bas M J 2000 IUGS reclassification of the high-Mg and picritic volcanic rocks; *J. Petrol.* **41** 1467–1470.
- Leterrier J, Maurry R C, Thonon P, Girard D and Marchal M 1982 Clinopyroxene composition as a method of identification of the magmatic affinities of paleo-volcanic series; *Earth Planet. Sci. Lett.* **59** 139–154.
- Loucks R R 1990 Discrimination of ophiolitic from nonophiolitic ultramafic–mafic allochthons in orogenic belts by the Al/Ti ratio in clinopyroxene; *Geology* **18** 346–349.
- MacGregor A G 1931 Clouded feldspars and thermal metamorphism; *Min. Mag.* **22** 524–538.
- Mallick A K and Sarkar A 1994 Geochronology and geochemistry of mafic dykes from precambrians of Keonjhar Orissa; *India Min.* **48** 3–24.
- Mallikharjuna Rao J, Poornachandra Rao M, Widdowson M and Kelley S P 2005 Evolution of proterozoic mafic dyke swarms of the Bundelkhand Granite massif, Central India; *Curr. Sci.* **88(3)** 502–506.
- Mandal N, Mitra A K, Misra S and Chakraborty C 2006 Is the outcrop topology of dolerite dykes of Precambrian Singhbhum craton fractal?; *J. Earth. Syst. Sci.* **115** 643–660.
- Mc Mullin David W A, Barr Sandra M and Raeside Robert P 2010 Very low and low grade metamorphism of mafic volcanic rocks of the Mira terraine (Avalonia), southeastern cape Breton Island, Nova Scotia; *Atlantic Geology* **46**, doi: 10.4138/atlgol.2010.006.
- Meschede M 1986 A method of discriminating between different types of mid ocean ridge basalts and continental tholeiites with the Nb–Zr–Y diagram; *Chem. Geol.* **56** 207–218.
- Mir A R, Alvi S H and Balaram V 2010 Geochemistry of mafic dikes in the Singhbhum Orissa craton: Implications for subduction-related metasomatism of the mantle beneath the eastern Indian craton; *Inter. Geol. Rev.* **52** 79–94.
- Mir A R, Alvi S H and Balaram V 2011 Geochemistry of the mafic dykes in parts of the Singhbhum Granitoid Complex: Petrogenesis and tectonic setting; *Arab. J. Geosci.* **4** 933–943.
- Mir A R, Alvi S H, Balaram V, Bhat F A, Sumira Z and Dar S A 2013 A subduction zone geochemical characteristic of the newer dolerite dykes in the Singhbhum craton, Eastern India; *Int. Res. J. Geol. Mining* **3** 213–223.
- Misra S, Deomurari M P, Wiedenbeck M, Goswami J N, Ray S and Saha A K 1999 $^{207}\text{Pb}/^{206}\text{Pb}$ zircon age and the evolution of the Singhbhum craton, eastern India: An ion microprobe study; *Precamb. Res.* **93** 139–151.
- Misra S 2006 Precambrian chronostratigraphic growth of Singhbhum–Orissa Craton, eastern Indian Shield: An alternative model; *J. Geol. Soc. India* **67** 356–378.
- Mottl M J 1983 Metabasalts, axial hot springs, and structure of hydrothermal systems at mid-ocean ridges; *Geol. Soc. Am. Bull.* **94** 161–180.
- Mottl M J and Holland H D 1978 Chemical exchange during hydrothermal alteration of basalt by sea water–Experimental results for major and minor components of seawater; *Geochim. Cosmochim. Acta* **42** 1103–1115.
- Mukherjee J, Beukes N J, Armstrong R A, Zimmermann V, Ghosh G and Medda R A 2008 Dating the oldest greenstone in India: A 3.51 Ga precise U–Pb SHRIMP zircon age for dacitic lava of the southern iron ore group, Singhbhum Craton; *J. Geol.* **116** 449–461.
- Mukhopadhyay D 2001 The Archaeana nucleus of Singhbhum: The present state of knowledge; *Gondwana Res.* **4(3)** 307–318.
- Mullen E D 1983 MnO/TiO₂/P₂O₅: A major element discriminant for basaltic rocks of oceanic environments and its implication for petrogenesis; *Earth Planet. Sci. Lett.* **62** 53–62.
- Murthy N G K 1987 Mafic dyke swarms of the Indian shield; In: Mafic dyke swarms (eds) Halls H C and Fahrig W F, *Geol. Assoc. Canada Spec. Paper* **34** 393–400.
- Murthy N G K 1995 Proterozoic mafic dykes in southern peninsular India; *Geol. Soc. India Memoir* **33** 81–98.
- Pati J K, Raju S, Malviya V P, Bhusan R, Prakash K and Patel S C 2008 Mafic dykes of Bundelkhand Craton, central India: Field, petrological and geochemical characteristics; In: *Indian dykes* (eds) Srivastava R K, Sivaji Ch and Rao C (New Delhi: Narosa Publ.), pp. 548–569.
- Pearce J A 1982 Trace element characteristics of lavas from destructive plate boundaries; In: *Andesites* (ed.) Thorpe R S, John Wiley and Sons, pp. 525–548.
- Pichamuthu C S 1959 The significance of clouded plagioclase in the basic dykes of Mysore state, India; *J. Geol. Soc. India* **1** 68–79.
- Plumper O, Piazzolo S and Austrheim H 2012 Olivine pseudomorphs after serpentinized orthopyroxene record transient oceanic lithospheric mantle dehydration (Leka Ophiolite Complex, Norway); *J. Petrol.* **53(9)** 1943–1968.
- Poldervaart A and Gilkey A K 1954 On clouded feldspars; *Am. Mineral.* **39** 75–91.
- Robertson L D M and Van Breeman O 1970 The southern satellite dykes of the Great Dyke, Rhodesia; *Geol. Soc. South Af. Spec. Publ.* **1** 621–644.

- Roy A B and Bhattacharya H N 2012 Tectonostratigraphic and geochronologic reappraisal constraining the growth and evolution of Singhbhum Archean craton, eastern India; *J. Geol. Soc. India* **80** 455–469.
- Saha A K 1948 Kolhan series: Iron ore series boundary to the east and southwest of Chaibasa; *Sci. Cult.* **14** 77–79.
- Saha A K 1952 On porphyry dykes near Dalsara, Mayurbhanj; *Sci. Cult.* **18** 250–253.
- Saha A K 1994 Crustal evolution of north Singhbhum-Orissa, eastern India; *Geol. Soc. India Memoir* **27** 341.
- Saha A K, Bose M K, Sankaran A V and Bhattacharyya T K 1972 Petrology and geochemistry of the ultramafic intrusion of Keshargaria, Singhbhum, Bihar; *J. Geol. Soc. India* **13** 113–121.
- Saha A K, Sankaran A V and Bhattacharyya T K 1973 Geochemistry of the newer dolerite suite of intrusions within the Singhbhum granite; *J. Geol. Soc. India* **14** 324–346.
- Sengupta P and Ray A 2012 Newer dolerite dykes, Jharkhand, India: A case study of magma generation, differentiation and metasomatism in a subduction zone setting; *Geochem. J.* **46** 477–491.
- Seyfried W E Jr, Berndt M E and Seewald J S 1988 Hydrothermal alteration processes at mid ocean ridges: Constraints from thermal alteration experiments, hot-spring fluids and composition of the oceanic crust; *Can. Mineral.* **26** 787–804.
- Shervais J W 1982 Ti-V plots and the petrogenesis of modern and ophiolitic lavas; *Earth Planet. Sci. Lett.* **59** 101–118.
- Sun S S and McDonough W F 1989 Chemical and isotopic systematics of oceanic basalts: Implication for mantle composition and processes; *Geol. Soc. Spec. Publ.* **42** 313–345.
- Zhang B 1988 Preliminary study on Proterozoic dyke swarms in north Sanxin; M.Sc. thesis, Peking University (in Chinese with English Abstract).
- Zhang B and Halls H C 1995 The origin and age of the feldspar clouding in the Matachewan dyke swarm; In: *Phys. Chem. Dykes* (eds) Baer G and Heimann A, Balkema, Rotterdam, pp. 171–176.

MS received 4 October 2013; revised 15 January 2014; accepted 23 January 2014