

## RESEARCH NOTE

# Cytosine deletion at AP2-box region of *HSP70* promoter and its influence on semen quality traits in crossbred bulls

RAJIB DEB<sup>1\*</sup>, BASAVRAJ SAJJANAR<sup>2</sup>, UMESH SINGH<sup>1</sup>, SUSHIL KUMAR<sup>1</sup>, GYANENDRA SINGH SENGAR<sup>1</sup>, RANI ALEX<sup>1</sup>, A. K. DAS<sup>1</sup>, S. TYAGI<sup>1</sup>, T. V. RAJA<sup>1</sup>, R. R. ALYETHODI<sup>1</sup>, RANI SINGH<sup>1</sup>, V. BHANUPRAKASH<sup>1</sup> and B. PRAKASH<sup>1</sup>

<sup>1</sup>Molecular Genetics Laboratory, ICAR-Central Institute for Research on Cattle, Meerut 250 001, India

<sup>2</sup>School of Atmospheric Stress Management, ICAR-National Institute of Abiotic Stress Management, Baramati 413 115, India

[Deb R., Sajjanar B., Singh U., Kumar S., Sengar G. S., Alex R., Das A. K., Tyagi S., Raja T. V., Alyethodi R. R., Singh R., Bhanuprakash V. and Prakash B. 2016 Cytosine deletion at AP2-box region of *HSP70* promoter and its influence on semen quality traits in crossbred bulls. *J. Genet.* **95**, 1035–1038]

## Introduction

Present investigation identified the effect of genetic polymorphism within the AP2 box region 12 located at bovine *HSP70* promoter on certain semen quality traits among crossbred bulls revealed the fact that superior capability of wild type promoter genotype over the deletion mutation at *HSP70* promoter may partly elucidate better semen quality traits for their contribution in selection of better quality bulls.

Although artificial insemination (AI) is now extensively used in cattle breeding, there is about 40–50% loss in bull semen quality during the freezing–thawing process, including sperm motility, effective survival time and antioxidant enzymes activity (Hu *et al.* 2011). Recently, studies have been carried out in our laboratory to unveil the genetic basis of semen quality traits in cattle (Kumar *et al.* 2014, 2015a, b). It is well known fact that mammalian spermatogenic cells are sensitive to heat stress (Yin *et al.* 1997). Studies have established that heat shock proteins (HSPs), particularly, the abundantly expressed 70 kDa HSP (*HSP70*), plays an important role in acquired thermotolerance (Nover 1991) and *HSP70* has been suggested to function as an indicator of thermotolerance in cells (Leung *et al.* 1996). *HSP70* is known to play an important role in sperm function after ejaculation and was identified in proteins extracted from sperm of bull (Kamaruddin *et al.* 1996) and boar (Huang *et al.* 1999). Huang *et al.* (1999) reported that both constitutive and inducible forms of *HSP70* are present in ejaculated spermatozoa of boars and both the levels of *HSP70* in spermatozoa and semen quality traits of boars significantly declined during the hot season.

Owing to the ability of *HSP70* to combat heat stress, the study was designed to analyse the genetic variation at AP2 box region of *HSP70* (*HSPA1A*) promoter and their association with semen quality parameters among Frieswal crossbred bulls (HF × Sahiwal) maintained in India. Our previous studies indicated that mutant variants of AP2 box region of *HSP70* significantly affected cellular thermotolerance, physiological parameters and milk production traits among Frieswal crossbred cows (Deb *et al.* 2013). The present study is the further extension of our earlier reports to investigate the effect of AP2 box variation on the semen quality traits among crossbred bulls.

## Materials and methods

A total of 96 mature crossbred bulls maintained at bull rearing unit were included in the present investigation. Semen samples were collected using artificial vagina from each bull during the period, from December 2010 to November 2011. Immediately after collection, the ejaculates were stored at 34°C in a water bath to evaluate the fresh semen quality traits including semen volume per ejaculate (mL), sperm motility (%), viability (%) and sperm concentration (M/mL). The details on the evaluation of good/poor sperm quality can be obtained from our earlier work by Kumar *et al.* (2015c). The fresh semen was then diluted with glycerol–egg yolk–citrate–Tris extender, processed and cryopreserved. After storing in liquid nitrogen for 1–2 days, two straws were randomly obtained from each ejaculate and thawed at 37°C for 60 s and immediately evaluated for postthaw motility (PTM) (%) using light microscopy. Genomic DNA was extracted from the sperm using GenElute™ Blood Genomic DNA kit (Sigma, Aldrich, USA). The DNA samples were

\*For correspondence. E-mail: drrajibdeb@gmail.com.

**Keywords.** AP2 box; promoter; *HSP70*; semen quality; crossbred bulls.


collected in elution buffer (supplied with kit) and stored at  $-20^{\circ}\text{C}$  for future use. Genomic DNA obtained from the ejaculated spermatozoa was subjected to genotyping using a double polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) (*ScrFI*) methodology developed earlier by our groups (Deb *et al.* 2013). At base position, in 895 of AP2 box region of *HSP70* (GenBank accession number M98823), deletion of cytosine residue was observed which created a *ScrFI* restriction site. The PCR was performed in a total volume of 100  $\mu\text{L}$  containing  $\sim 0.5$   $\mu\text{g}$  genomic DNA, 10 mM Tris-HCl, 50 mM KCl, 1.5 mM  $\text{MgCl}_2$ , 0.01% gelatine, 250 mM of each dNTP, and 5 U *Taq* polymerase. Initial denaturation was for 5 min (at  $94^{\circ}\text{C}$ ) followed by 35 cycles of  $94^{\circ}\text{C}$  denaturation (1 min),  $63^{\circ}\text{C}$  annealing (1 min) and  $72^{\circ}\text{C}$  extension (2 min). The PCR product was detected by electrophoresis in 1% agarose. Two sets of diagnostic primers used for first round amplification to generate *HSP70* gene-specific DNA fragment followed by a mutated set of nested primers (Deb *et al.* 2013) to reamplify the generated PCR products from first round PCR to abolish three native restriction sites for *ScrFI* enzyme and thus, only the diagnostic *ScrFI* site remain intact. Frequencies of the existing polymorphisms in the crossbred bulls were resolute through restriction digestion of the second round PCR products with *ScrFI* restriction enzyme. Different genotypes obtained were subjected for TA cloning using pTZ57R plasmid (InsTAclone PCR Cloning kit, Fermentas, USA). The positive clones were confirmed by restriction enzyme analysis. Positive clone from each genotype was subjected to DNA sequencing. The nested PCR products (encoding the transcription factor binding site) obtained from different genotypic groups were subjected to electromobility assay using Gel Shift Assay kit (Promega, Madison, USA). Briefly, PCR product (1  $\mu\text{L}$ ) from each genotype was incubated with HeLA cell extract (9  $\mu\text{L}$ ) and subjected to 4% polyacrylamide gel electrophoresis. Silver staining technique was used for visualizing the banding pattern of each group. Appropriate positive (AP2 oligo and HeLA cell extract) and negative (only AP2 oligo) controls were run along with the samples. The effect of the cytosine deletion at AP2 box region of *HSP70* on semen quality parameters, including semen volume, sperm concentration, motility, postthaw motility and viability were investigated using independent *t*-test using SAS 9.3 (SAS Institute, Cary, USA). Specific transformation was applied on semen quality traits, namely, motility, postthaw motility and viability, which was obtained by considering Box and Cox (1964) transformation class and the results were expressed as mean  $\pm$  standard error in the original scale.

## Results and discussion

The double PCR-RFLP using *ScrFI* confirmed the occurrence of homozygous wild-type genotype (CC: 49 and 47 bp) and heterozygous cytosine deletion mutant genotype (C-: 96, 49 and 47 bp) (figure 1, a-c). The genotype frequency

of CC was lower (0.47) than C- (0.52) (table 1). The electromobility gel-shift assays of AP2 binding pattern in two different genotypes with nuclear extracts from HeLA cells are depicted in figure 1d. The 96 base pair nested PCR product from each genotype was subjected to gel shift assay and it was concluded that the concentration of free DNA was higher in heterozygous deletion mutant (C-) than homozygous wild (CC) types. CC genotypes significantly ( $P < 0.01$ ) contribute to better fresh spermatozoal motility ( $58.78 \pm 0.89$ ) and viability ( $71.83 \pm 1.06$ ) as well as postthaw motility ( $56.87 \pm 1.24$ ) compared to C-genotypes (table 1).

Binding of AP2 transcription factor at the AP2 box region of the *HSP70* promoter sequences were reported by Tacchini *et al.* (1995). Schwerin *et al.* (2003) reported that polymorphisms within the promoter may alter the potential *cis*-acting elements and are also linked with relative thermotolerance in terms of 'productive life time' in Holstein breed of cattle. Further, our earlier studies showed that polymorphism within the promoter region of *HSP70* gene is strongly associated with the thermotolerance and other production traits in crossbred cows (Deb *et al.* 2013). In the present investigation, we have identified polymorphism within the *cis*-acting element of *HSP70* promoter by double PCR-RFLP method among the cross bred bull semen. Our present findings revealed that presence of the heterozygous cytosine deletion mutant genotype (C-) in majority with the complete absence of homozygous cytosine deletion mutant genotypes (--) in the targeted population. However, Schwerin *et al.* (2003), reported the presence of polymorphism at the same position having all three genotypes in pure Holstein dairy cows in contrast to the presence of only two genotypes in Indian HF crossbred cows (Deb *et al.* 2013). Gel shifting analysis revealed that AP2 transcription factor can interact almost equally at the promoter region of both the genotypes; however, the level of free DNA residues was more in C-genotypes than CC. It may be due to the basis that as the C-DNA are derived from heterozygous, the interaction of transcription factor with promoter is partial. Percentage of normal sperm, motility and sperm concentration are common criteria for evaluating semen quality (Den Daas 1992). Many studies have demonstrated that heat stress during the summer decreases semen quality (Kuo *et al.* 1997). Ejaculated spermatozoa are highly differentiated cells which are short of the biosynthetic machinery to cope with adverse environmental impacts (Hammersdtedt *et al.* 1990). Association analysis with the fresh semen quality traits, the deletion showed significant association ( $P < 0.01$ ) with individual spermatozoal motility and viability traits. Correspondingly, Shrum *et al.* (2010) and Gafer *et al.* (2015) reported that associations between a SNP in *HSP70* promoter and sperm motility as well as velocity traits in bulls, which may be due to persuade of *HSP70* on defense of proteins related to respiration activity and energy level in spermatozoa (Nascimento *et al.* 2008). The study also highlighted that deletion at AP2 box region also significantly ( $P < 0.01$ ) effect on postthaw motility of bull semen after cryopreservation. As the motility of



**Figure 1.** Double PCR-RFLP for genotyping AP2 box region of *HSP70* promoter and its influence on AP2 factor binding. (a) First round (550 bp) PCR with F(W) and R(W) primer and (b) Second round (96 bp) PCR with F(M) and R(M) primer set. M: 100-bp ladder. (c) PCR-RFLP banding pattern of AP2 genotypes. Lane 1, 5–8: CC genotypes (49 and 47 base pair); Lane 2–4: C-genotypes (96, 49 and 47 base pair), M: 100-bp DNA ladder. (d) Electromobility gel-shift assays of AP2 binding pattern in two different genotypes with nuclear extracts from HeLA cells. Lane 1: positive control (HeLA extrac. + AP2 oligo), lanes 2–3: CC genotype (HeLA + PCR product of CC), lanes 4–5: C-genotypes (HeLA + PCR product of C–), lane 6: negative control (only AP2 oligo).

**Table 1.** Genotype frequency and their association with semen quality traits among crossbred bulls.

Genotypes	Genotype frequency	Fresh samples				Cryopreserved samples
		Volume (mL)	Concentration (M/mL)	Motility (%)**	Viability (%) **	Postthaw motility (%)**
C– (n = 46)	0.47	4.36 ± 0.27	970.86 ± 111.11	45.00 ± 2.31	48.64 ± 2.69	33.37 ± 1.87
CC (n = 50)	0.52	4.86 ± 0.66	970.07 ± 97.87	58.78 ± 0.89	71.83 ± 1.06	56.87 ± 1.24

\*\**P* < 0.01.

the postthawed semen indicate its fertility status, we have not studied the viability of postthawed semen samples. Moreover, under Indian climatic conditions, the semen samples are generally evaluated for their viability before freezing and not after freezing. It is a routine practice that the frozen semen samples are evaluated only on the basis of their postthaw motility and not by viability after cryopreservation. However, the study on the viability of the postthawed semen using computer assisted semen analysis (CASA) would add much more knowledge on the fertilizing capacity of the frozen semen samples. Polymorphisms at AP2 box region of

*HSP70* gene can influence on semen quality traits and thus, suggest to be used in marker-assisted selection for antiheat stress as well as superior reproductive bulls. Nevertheless, further study on the evaluation of viability of the postthaw semen samples using CASA needs to be conducted to support the results obtained in the present study. Further, the present results could be validated even in larger bull populations using CASA to find the effect of deletion at AP2 box region on postthaw motility and viability parameters of cryopreserved semen before utilizing it as an early marker for selection of bulls in artificial insemination programmes.

### Acknowledgements

We are thankful to Director, ICAR-CIRC for providing necessary facilities to conduct the work. We also acknowledge Science and Engineering Research Board, Government of India, for providing financial support under the project YSS/2014/000279 to RD. We are also thankful to the Military Farm, Meerut, India, for providing the animals used in the present experiment.

### References

- Box G. E. and Cox D. R. 1964 An analysis of transformations. *J. Roy Stat. Soc.* **26**, 211–252.
- Deb R., Sajjanar B., Singh U., Kumar S., Brahmane M. P., Singh R. et al. 2013 Promoter variants at AP2 box region of *HSP70.1* affect thermal stress response and milk production traits in Frieswal cross bred cattle. *Gene* **15**, 230–235.
- Den Daas N. 1992 Laboratory assessment of semen characteristics. *Anim. Reprod. Sci.* **28**, 87–94.
- Gafer J. A., Abd El-Rahman G. H. and Rawash Z. M. 2015 Association of *HSP70* gene polymorphism and bull semen quality in winter and summer seasons. *Alexandria J. Vet. Sci.* **46**, 1–14.
- Hammersdtedt R. H., Graham J. K. and Nolan J. P. 1990 Cryopreservation of mammalian sperm: what we ask them to survive. *J. Androl.* **11**, 73–88.
- Hu J. H., Zhao X. L., Tian W. Q., Zan L. S. and Li Q. W. 2011 Effects of vitamin E supplementation in the extender on frozen-thawed bovine semen preservation. *Animal* **5**, 107–112.
- Huang S. Y., Kuo Y. H., Lee W. C., Tsou H. L., Lee Y. P., Chang H. L. et al. 1999 Substantial decrease of heat-shock protein 90 precedes the decline of sperm motility during cooling of boar spermatozoa. *Theriogenology* **51**, 1007–1016.
- Kamaruddin M., Kroetsch T., Basrur P. K. and King W. A. 1996 Heat shock protein 70 in bovine semen. *Biol. Reprod.* **54**, 222.
- Kumar S., Deb R., Singh U., Ganguly I., Mandal D. K., Singh R. et al. 2014 SNPs at exonic region of Aquaporin-7 (AQP-7) gene may effects semen quality parameters among crossbred bulls. *J. Genet.* **93**, Online Resources. e108–e112.
- Kumar S., Deb R., Singh U., Ganguly I., Mandal D. K., Singh R. et al. 2015a Bovine circadian locomotor output cyclic kaput (CLOCK) and cluster in (CLU) mRNA quantitation in ejaculated crossbred bull spermatozoa. *Reprod. Domest. Anim.*, doi: 10.1111/rda.12522.
- Kumar S., Singh U., Deb R., Tyagi S., Mandal D. K., Kumar M. et al. 2015b A SNP (g.358A>T) at intronic region of CD9 molecule of crossbred bulls may associate with spermatozoal motility. *Meta Gene*, (doi: 10.1016/j.mgene.2015.07.004).
- Kumar S., Singh U., Ganguly I., Deb R., Singh R., Mann S. et al. 2015c Protamine 3 expressions in crossbred bull spermatozoa may not be a prognostic marker for differentiating good and poor quality semen. *African J. Biotechnol.* **13**, 20.
- Kuo Y. H., Huang S. Y. and Lee Y. P. 1997 Effects of breed and season on semen characteristics of boars in subtropical area. *J. Chin. Soc. Vet. Sci.* **23**, 114–122.
- Leung S. M., Senisterra G., Ritchie K. P., Sadis S. E., Lepock J. R. and Hightower L. E. 1996 Thermal activation of the bovine *HSC70* molecular chaperone at physiological temperatures: physical evidence of a molecular thermometer. *Cell Stress. Chaperon.* **1**, 78–89.
- Nascimento J. M., Shi L. Z., Tam J., Chandsawangbhuwana C., Durrant B., Botvinick E. L. et al. 2008 Comparison of glycolysis and oxidative phosphorylation as energy sources for mammalian sperm motility, using the combination of fluorescence imaging, laser tweezers, and real-time automated tracking and trapping. *J. Cell Physiol.* **217**, 745–751.
- Nover L. 1991 *Heat shock response*. CRC Press, Boca Raton, USA.
- Schwerin M., Sanftleben H. and Grupe S. 2003 Genetic predisposition for productive life is associated with functional inactivation of a AP2-binding site in the promoter of the stress protein 70.1-encoding gene in cattle. *Arch. Tierz., Dummerstorf.* **46**, 177–185.
- Shrum K., Lester T., Rorie R., Reiter S., Looper M. and Rosenkrans J. C. F. 2010 Effects of heat shock protein 70 haplotype and tall fescue variety on bull sperm characteristics. *Arkansas Agric. Res. Stat. Res. Ser.* **584**, 39–44.
- Tacchini L., Pogliaghi G., Radice L., Anzon E. and Bernelli-Zazzera 1995 A differential activation of heat-shock and oxidation-specific stress genes in chemically induced oxidative stress. *Biochem. J.* **309**, 453–459.
- Yin Y., Hawkins K. L. and Dewolf W. C. 1997 Heat stress causes testicular germcell apoptosis in adult mice. *J. Androl.* **18**, 159–165.

Received 13 February 2016, in revised form 13 April 2016; accepted 5 May 2016

Unedited version published online: 6 May 2016

Final version published online: 29 November 2016

Corresponding editor: SILVA GARAGNA