

## **An Operator Perturbation Method of Polarized Line Transfer V. *Diagnosis of Solar Weak Magnetic Fields***

K. N. Nagendra<sup>1,2\*</sup>, H. Frisch<sup>2</sup>, M. FaurobertScholl<sup>2</sup> & F. Paletou<sup>2</sup>

<sup>1</sup>*Indian Institute of Astrophysics, Bangalore 560034, India.*

<sup>2</sup>*Observatoire de la Côte d'Azur, (CNRS/UMR 6529), BP. 4229, F06304, Nice Cedex 4, France.*

*\*email:knn@iiap.ernet.in*

**Abstract.** We present an application of the PALI (Polarized Approximate Lambda Iteration) method to the resonance scattering in spectral lines formed in the presence of weak magnetic fields. The method is based on an operator perturbation approach, and can efficiently give solutions for oriented vector magnetic fields in the solar atmosphere.

*Key words.* Polarization—magnetic fields, radiative transfer—stars: atmospheres—methods: numerical.

### **1. Introduction**


We refer to polarized spectral lines formed outside the active regions having ‘weak’ oriented fields ( $0 \leq B \leq 300$  G) which cause de-polarization in Stokes- $Q$ , and rotation of the plane of polarization in Stokes- $U$  parameter of the ‘resonantly scattered’ line radiation. This phenomenon called ‘Hanle Effect’, is invoked to explain the linear polarization changes observed in lower chromospheric resonance lines such as Ca I 4227 Å, Sr 4607 Å and Sr II 4078 Å (see Bianda *et al.* 1999 for observational Hanle diagnostics). The theoretical interpretation of such data demands the solution of NLTE polarized line transfer equation, for several combinations of independent parameters.

### **2. Results and discussions**

The details of the Hanle scattering problem are described in Nagendra *et al.* (1998, 1999). The coherent superposition of radiatively broadened Zeeman sub-states in weak fields ( $\Delta v_L \leq \Gamma_R$ ), is responsible for the Hanle effect. The Stokes- $V$  parameter is negligible in weak fields.

#### *2.1 The two-parameter polarization diagrams*

Determination of field parameters from observed data can be attempted with the help of ‘two-parameter polarization diagrams’ showing a network of *iso-strength* and



**Figure 1.** (a): Hanle effect in weak magnetic fields ( $B = 10 - 300$  G), when ( $\Delta v_L \approx \Gamma_R$ ). Pure Zeeman effect ( $B = 1000 - 3000$  G) corresponds to ( $\Delta v_L \gg \Gamma_R$ ); (b): Geometry specifying the direction of the magnetic field  $\mathbf{B}$  and of the line-of-sight  $\mathbf{n}$ . Angles  $\theta$  and  $\theta_B$  are the colatitudes of  $\mathbf{n}$  and  $\mathbf{B}$ , respectively. The azimuthal angles  $\varphi$  and  $\varphi_B$  are measured from the  $x$ -axis in an anti-clockwise direction in the  $xy$ -plane – that also represents the 1-D plane parallel slab atmosphere. The  $z$ -axis is the vertical direction. The light gray bands represent the radiative width  $\Gamma_R$ .

*iso-azimuth* curves. For a given line of sight, determined by the values of  $\theta$  and  $\varphi$  (Fig. 1b), we choose a value of  $\theta_B$  and vary the two other parameters of the vector magnetic field,  $\gamma_B$  and  $\varphi_B$ . The field strength parameter is  $\gamma_B = 2\pi\Delta v_L g_U / \Gamma_R$ , where  $g_U$  is the Lande  $g$ -factor of the upper level with a radiative width  $\Gamma_R$ , and  $\Delta v_L$  is the Larmour frequency (Fig. 1a). If an observational data point ( $U/I$ ,  $Q/I$ ) falls within an interval defined by  $[\Delta\gamma_B, \Delta\varphi_B]$ , we get upper and lower limits on the possible values of  $\gamma_B$  and  $\varphi_B$ . This approach is useful when an independent estimate of  $\theta_B$  is available. We now point out the generalization of the relevant equations in Nagendra *et al.* (1998, 1999), where the meaning of mathematical symbols in equations (1) and (2), can be found. For PRD mechanism,  $\bar{\mathbf{J}}(\tau, x)$  is given by (when  $\varphi_B$  is constant with depth):


$$\bar{\mathbf{J}}(\tau, x) = \hat{\mathbf{R}}(\varphi_B) \bar{\mathbf{J}}^0(\tau, x), \quad (1)$$

where  $\bar{\mathbf{J}}^0(\tau, x)$  is the *reduced mean intensity* computed for the special case of field azimuth  $\varphi_B = 0$ . Notice that one can get the  $\bar{\mathbf{J}}(\tau, x)$  for an arbitrary value of  $\varphi_B$ , using the above equation. The corresponding line source vector is:

$$\mathbf{S}_l(\tau, x) = (1 - \varepsilon(\tau)) \hat{\mathbf{H}}_B(\theta_B, \varphi_B, \mathbf{B}) \bar{\mathbf{J}}(\tau, x) + \mathbf{S}^{th}(\tau), \quad (2)$$

which gives the *reduced specific intensities* by application of a formal solution of the transfer equation. The iso-azimuth curves are computed by solving the Hanle transfer problem fully, for several values of  $\gamma_B$ . Iso-strength curves however are computed rapidly by taking advantage of the Hanle symmetry mentioned above. The polarization diagrams are easy to construct since one can express  $I$ ,  $Q$  and  $U$  in terms of the six components of the ‘azimuth independent reduced specific intensity vector’ written as  $\mathbf{I} = (I_1, I_Q, I_{+1}, I_{-1}, I_{+2}, I_{-2})^T$  (see equations (81)(83) in Nagendra *et al.* 1998).

Details of constructing the well known “two-parameter polarization diagrams” are presented in Nagendra *et al.* (1998). Exactly identical model parameters are employed



**Figure 2.** The twoparameter polarization diagrams for Hanle effect in an optically thin line. A comparison of CRD and PRD mechanisms is presented. The magnetic field is assumed to be horizontal ( $\theta_B = 90^\circ$ ). The field strength parameter  $\gamma_B$  and field azimuth  $\varphi_B$  are independent parameters.  $\gamma_B$  varies from 0–100 (dashed horizontal lines moving down to up); and  $\varphi_B$  varies from 0–180° (solid vertical lines moving from left to right). The *iso-azimuth curves* (solid lines) are drawn by fixing  $\varphi_B$  and varying  $\gamma_B$ . The *iso-strength curves* (dashed lines) are drawn by fixing  $\gamma_B$  and varying  $\varphi_B$ . The line of sight (LOS) is fixed at  $(\theta, \varphi) = (90^\circ, 0^\circ)$ . The results for four frequencies  $x = 0, 3, 5.3$  and 10 are presented. The iso-strength curves show the *Hanle de-polarization* and *saturation effects* clearly. The iso-azimuth curves show the effect of *rotation of the plane of polarization* when  $\gamma_B$  varies. Iso-azimuth curves meet at the point  $\gamma_B = 0$  where  $(U/I = 0)$ , and for  $\varphi_B = (90^\circ, 270^\circ)$ , they are parallel to the  $(U/I) = 0$  axis. The Hanle de-polarizing ability is maximum for  $\theta_B = 90^\circ$ .

in computing Fig. 15 of that paper, and the Fig. 2 in this paper. In Fig. 2, the two basic mechanisms of line scattering are considered: the *Partial Frequency Redistribution* (PRD) which is physically more realistic for resonance lines, compared to the *Complete Frequency Redistribution* (CRD). The CRD diagrams uniformly decrease in size (degree of linear polarization ( $Q/I, U/I$ )), and reach a constant level for frequencies  $x > 10$ . The PRD diagrams show a strong sign reversal at  $x = 3$ , where polarization is  $\approx 0$  (panel 2); become fully positive (panel 3); and reach zero polarization for large frequencies (panel 4). They follow the well known physical characteristics of CRD and PRD (see Fig. (1b) of Faurobert 1987).

### 3. Conclusions

The PALI is about 100 times faster than the conventional methods of solving the Hanle scattering problem. It is suitable for realistic modeling of the Hanle effect observations in spectral lines formed on the Sun. The Hanle effect is useful in exploring the spatially unresolved vector magnetic fields in a parameter space where the ordinary Zeeman effect is not practically useful.

### References

- Bianda, M., Stenflo, J. O., Solanki, S. K. 1999, in *Solar Polarization, 2nd SPW*, (ed.) K. N. Nagendra & J. O. Stenflo, (Dordrecht: Kluwer), ASS **243**, 31.
- Faurobert-Scholl M. 1987, *Astron. Astrophys.* **178**, 269.
- Nagendra, K. N., Frisch, H., Faurobert-Scholl, M. 1998, *Astron. Astrophys.* **332**, 610.
- Nagendra, K. N., Paletou, F., Frisch, H., Faurobert-Scholl. 1999, in *Solar Polarization, 2nd SPW*, (ed.) K. N. Nagendra & J. O. Stenflo, (Dordrecht: Kluwer), ASSL **243**, 127.