

Kolmogorov's existence theorem for Markov processes in C^* algebras

B V RAJARAMA BHAT and K R PARTHASARATHY

Indian Statistical Institute, 7, S.J.S. Sansanwal Marg, New Delhi 110016, India

Dedicated to the memory of Professor K G Ramanathan

Abstract. Given a family of transition probability functions between measure spaces and an initial distribution Kolmogorov's existence theorem associates a unique Markov process on the product space. Here a canonical non-commutative analogue of this result is established for families of completely positive maps between C^* algebras satisfying the Chapman-Kolmogorov equations. This could be the starting point for a theory of quantum Markov processes.

Keywords. Completely positive map; Markov process; GNS principle.

1. Introduction

Let (X_i, \mathcal{F}_i) , $i = 0, 1, 2, \dots$ be Polish measurable spaces and let $P_i(x_i, dx_{i+1})$ be a transition probability from (X_i, \mathcal{F}_i) to $(X_{i+1}, \mathcal{F}_{i+1})$ for each i . Given a probability measure μ on (X_0, \mathcal{F}_0) it follows from Kolmogorov's extension theorem that there exists a unique probability measure P_μ on the infinite product space $(\Omega, \mathcal{F}) = \bigotimes_{i=0}^{\infty} (X_i, \mathcal{F}_i)$ such that, for every finite n , its projection or marginal distribution P_μ^n in $\bigotimes_{i=0}^n (X_i, \mathcal{F}_i)$ is given by

$$P_\mu^n(E_0 \times E_1 \times \dots \times E_n) = \int_{E_0 \times E_1 \times \dots \times E_n} \mu(dx_0) P_0(x_0, dx_1) P_1(x_1, dx_2) \dots P_n(x_{n-1}, dx_n) \quad (1.1)$$

for all $E_i \in \mathcal{F}_i$, $i = 0, 1, 2, \dots, n$. The probability space $(\Omega, \mathcal{F}, P_\mu)$ describes the Markov process with initial distribution μ and transition probability $P_i(\cdot, \cdot)$ for transition from a state at time i to a new state at time $i + 1$. This can be described in a $*$ algebraic language as follows. Denote by \mathcal{A}_i the commutative $*$ algebra of all complex valued bounded measurable functions on (X_i, \mathcal{F}_i) . Introduce the positive unital operator $T(i, i + 1): \mathcal{A}_{i+1} \rightarrow \mathcal{A}_i$ by

$$(T(i, i + 1)g)(x_i) = \int g(x_{i+1}) P_i(x_i, dx_{i+1}).$$

For any $i \leq k$ define $T(i, k): \mathcal{A}_k \rightarrow \mathcal{A}_i$ by

$$T(i, k) = \begin{cases} \text{identity} & \text{if } i = k, \\ T(i, i + 1) T(i + 1, i + 2) \dots T(k - 1, k) & \text{if } i < k. \end{cases}$$

The family $\{T(i, k), i \leq k\}$ of transition operators obeys the Chapman-Kolmogorov equations:

$$T(i, k) T(k, \ell) = T(i, \ell) \quad \text{for } i \leq k \leq \ell.$$

Let \mathcal{H} be the Hilbert space $L^2(P_\mu)$ and $F(i)$ denote the Hilbert space projection on the subspace of functions depending only on the first $i + 1$ coordinates (x_0, x_1, \dots, x_i) of $\omega = (x_0, x_1, x_2, \dots)$ in Ω . Then $\{F(i)\}$ is an increasing sequence of projections in \mathcal{H} . For any $g \in \mathcal{A}_i$ define the operator $j_i(g)$ in \mathcal{H} by

$$(j_i(g)\phi)(\omega) = g(x_i)(F(i)\phi)(\omega), \quad \omega = (x_0, x_1, \dots).$$

Then j_i is a $*$ homomorphism from \mathcal{A}_i into the $*$ algebra $\mathcal{B}(\mathcal{H})$ of all bounded operators in \mathcal{H} . The Markov property of the stochastic process $(\Omega, \mathcal{F}, P_\mu)$ is encapsulated in the operator relations

$$j_k(1) = F(k), \tag{1.2}$$

$$F(i)j_k(g)F(i) = j_i(T(i, k)g), \quad g \in \mathcal{A}_k, \quad i \leq k. \tag{1.3}$$

The relations (1.1) can be expressed as

$$\begin{aligned} &\langle u, j_0(g_0)j_1(g_1)\cdots j_n(g_n)v \rangle \\ &= \int (\bar{u}v g_0)(x_0)g_1(x_1)\cdots g_n(x_n)dP_\mu(\omega) \end{aligned} \tag{1.4}$$

for all u, v in the range of $F(0)$ and $g_i \in \mathcal{A}_i, i = 0, 1, 2, \dots, n$. Here ω denotes the sequence (x_0, x_1, \dots) . We may call the triple $(\mathcal{H}, F, j_k, k = 0, 1, 2, \dots)$ consisting of the Hilbert space \mathcal{H} , the filtration of projections $F(k)$ increasing in k and the family $\{j_k, k = 0, 1, 2, \dots\}$ of $*$ (but nonunital) homomorphisms, a Markov process with transition operators $\{T(i, j), i \leq j\}$. A similar description of a Markov process in continuous time is also possible.

In the context of quantum or non-commutative probability theory there have been several partial attempts (for example, by Accardi, Frigerio and Lewis [AFL], Emch [E], Sauvageot [S] and Vincent-Smith [Vi-S]) to construct Markov processes when transition probabilities between measurable spaces, or equivalently, the transition operators between the corresponding commutative $*$ algebras of bounded measurable functions are replaced by unital and completely positive linear maps between unital $*$ algebras of operators in Hilbert spaces. In the present paper we shall start with a family of completely positive maps between C^* algebras which obey the Chapman-Kolmogorov equations and build a unique canonical minimal Markov process, using the GNS principle. Rather remarkably, this minimal process, when restricted to the centres of the different C^* algebras that are involved, can be obtained as a conditional expectation of a completely commutative process. The definition of a Markov process that we shall adopt is inspired by the equations (1.2)–(1.4).

2. The basic construction

Let \mathcal{A}_t be a unital C^* algebra of bounded operators in a complex Hilbert space \mathcal{H}_t , for every $t \geq 0$. The time index t here may be discrete or continuous. It is useful to

imagine any hermitian element $x \in \mathcal{A}_t$ as a real valued observable concerning a system at time t . For every $0 \leq s \leq t < \infty$ let $T(s, t): \mathcal{A}_t \rightarrow \mathcal{A}_s$ be a linear, unital and completely positive map (hereafter called simply a c.p. map) satisfying the following: (i) $T(s, s)$ is the identity map on \mathcal{A}_s ; (ii) $T(r, t) = T(r, s)T(s, t)$ for all $0 \leq r \leq s \leq t < \infty$. When (i) and (ii) hold we say that the family $\{T(s, t)\}$ of c.p. maps obeys the Chapman–Kolmogorov equations and call it a family of *transition operators*. Complete positivity is equivalent to the condition

$$\sum_{i,j} X_i^* \{ T(s, t)(Y_i^* Y_j) \} X_j \geq 0$$

for all bounded operators X_i in \mathcal{X}_s , and elements $Y_i \in \mathcal{A}_t$, the summation being over any finite index set. Another equivalent description of complete positivity is that, for every finite n , the matrix $((T(s, t)(Y_{ij})))_{1 \leq i, j \leq n}$, viewed as an operator in the n -fold direct sum $\mathcal{X}_s \oplus \dots \oplus \mathcal{X}_s$, is positive whenever $((Y_{ij}))_{1 \leq i, j \leq n}$ is positive in the n -fold direct sum $\mathcal{X}_t \oplus \dots \oplus \mathcal{X}_t$ with $Y_{ij} \in \mathcal{A}_t$ for each i, j .

Denote by $\Gamma_0(\mathbb{R}_+) = \Gamma_0$ the set $\{\sigma \subset \mathbb{R}_+, 0 \in \sigma, \#\sigma < \infty\}$, where $\#\sigma$ denotes the cardinality of σ . When $\#\sigma = n$ and $t_i \in \sigma, i = 1, 2, \dots, n$ are distinct we always express it as $\sigma = \{t_1, t_2, \dots, t_n\}$ with $t_1 > t_2 > \dots > t_n = 0$. When $X_{t_i} \in \mathcal{A}_{t_i}$ for each $i = 1, 2, \dots, n$ we denote the n -length sequence $\{X_{t_1}, X_{t_2}, \dots, X_{t_n}\}$ by $X(\sigma)$. Suppose that $\sigma = \{s_1, s_2, \dots, s_m\}$, $\delta = \{t_1, t_2, \dots, t_n\}$ and $\sigma \cup \delta = \{r_1, r_2, \dots, r_k\}$ are in Γ_0 . For any $X(\sigma)$ with $X_{s_i} \in \mathcal{A}_{s_i}$, we write $X^\sigma(\sigma \cup \delta)$ for the sequence $Y(\sigma \cup \delta)$ defined by

$$Y_{r_i} = \begin{cases} X_{s_j} & \text{if } r_i = s_j \text{ for some } j = 1, 2, \dots, m, \\ I_{r_i} & \text{otherwise,} \end{cases}$$

where I_r is the identity element in \mathcal{A}_r . Denote by \tilde{A} the set of all sequences of the form $X(\sigma)$ with σ varying in Γ_0 and write

$$\mathcal{M} = \tilde{A} \times \mathcal{X}_0, \tag{2.1}$$

$$\mathcal{M}_t = \begin{cases} \{(X(\sigma), u) \in \mathcal{M}, \sigma = (t, t_2, \dots, t_n), n = 2, 3, \dots\} & \text{if } t > 0 \\ \mathcal{A}_0 \times \mathcal{X}_0 & \text{if } t = 0 \end{cases} \tag{2.2}$$

To the family $\{T(s, t)\}$ of transition operators we now associate a function L_T on the set $\mathcal{M} \times \mathcal{M}$ as follows:

$$L_T((X(\sigma), u), (Y(\delta), v)) = \langle u, X_0^* \{ T(0, t_{n-1})(X_{t_{n-1}}^* \{ T(t_{n-1}, t_{n-2}) (\dots X_{t_2}^* \{ T(t_2, t_1)(X_{t_1}^* Y_{t_1}) \} Y_{t_2} \dots) \} Y_{t_{n-1}}) Y_0 \} v \rangle$$

if $\sigma = \{t_1, t_2, \dots, t_n\}$, (2.3)

and

$$L_T((X(\sigma), u), (Y(\delta), v)) = L_T((X^\sigma(\sigma \cup \delta), u), (Y^\delta(\sigma \cup \delta), v)). \tag{2.4}$$

PROPOSITION 2.1.

L_T is a positive definite kernel on $\mathcal{M} \times \mathcal{M}$, i.e., for any $n = 1, 2, \dots$, complex scalars c_i and elements $(X_i(\sigma_i), u_i) \in \mathcal{M}, i = 1, 2, \dots, n$ the following inequality holds:

$$\sum_{1 \leq i, j \leq n} \bar{c}_i c_j L_T((X_i(\sigma_i), u_i), (X_j(\sigma_j), u_j)) \geq 0 \tag{2.5}$$

Proof. We claim that for a pair of elements of the form $(X(\sigma), u), (Y(\delta), v)$ in \mathcal{M} and $\delta \in \Gamma_0$

$$L_T((X(\sigma), u), (Y(\delta), v)) = L_T((X^\sigma(\sigma \cup \delta), u), (Y^\delta(\sigma \cup \delta), v)). \tag{2.6}$$

It suffices to prove this relation when $\delta = \{t, 0\}$, $\sigma = \{t_1, t_2, \dots, t_{n-1}, 0\}$, $t \neq t_i$ for every i , since the more general case would follow by induction. In this special case (2.6) follows easily from (2.3) with σ replaced by $\sigma \cup \delta$ and the Chapman–Kolmogorov equations. In view of (2.4) it is enough to prove (2.5) when $\sigma_i = \sigma$ for each i , for otherwise, we may replace all the σ_i 's by $\sigma = \bigcup_i \sigma_i$. Let $\sigma = \{t_1, t_2, \dots, t_{m-1}, t_m = 0\}$ and

$$X_i(\sigma) = (X_{i_1}, X_{i_2}, \dots, X_{i_m}), \quad i = 1, 2, \dots, n.$$

Define inductively the following operators:

$$\begin{aligned} Z_{ij}(t_1) &= X_{i_1}^* X_{j_1} \\ Z_{ij}(t_r) &= X_{i_r}^* T(t_r, t_{r-1})(Z_{ij}(t_{r-1})) X_{j_r}, \\ r &= 2, 3, \dots, m. \end{aligned}$$

Clearly, the matrix $((Z_{ij}(t_1)))$ is a positive operator in the n -fold direct sum $\mathcal{X}_{t_1} \oplus \dots \oplus \mathcal{X}_{t_1}$. If $((Z_{ij}(t_{r-1})))$ is a positive operator in $\mathcal{X}_{t_{r-1}} \oplus \dots \oplus \mathcal{X}_{t_{r-1}}$ the complete positivity of $T(t_r, t_{r-1})$ implies that $((Z_{ij}(t_r)))$ is positive in $\mathcal{X}_{t_r} \oplus \dots \oplus \mathcal{X}_{t_r}$. Thus, by induction, $((Z_{ij}(t_m)))$ is a positive operator in $\mathcal{X}_0 \oplus \dots \oplus \mathcal{X}_0$. If we write

$\xi = \bigoplus_{i=1}^n c_i u_i$ in $\mathcal{X}_0 \oplus \dots \oplus \mathcal{X}_0$ we have

$$\sum_{1 \leq i, j \leq n} \bar{c}_i c_j L_T((X_i(\sigma), u_i), (X_j(\sigma), u_j)) = \langle \xi, ((Z_{ij}(t_m))) \xi \rangle \geq 0. \quad \blacksquare$$

PROPOSITION 2.2.

There exists a Hilbert space \mathcal{H} and a map $\lambda: \mathcal{M} \rightarrow \mathcal{H}$ satisfying the following:

- (i) $\langle \lambda(X(\sigma), u), \lambda(Y(\delta), v) \rangle \equiv L_T((X(\sigma), u), (Y(\delta), v))$;
- (ii) The set $\{\lambda(X(\sigma), u) | (X(\sigma), u) \in \mathcal{M}\}$ is total in \mathcal{H} ;
- (iii) If \mathcal{H}' is another Hilbert space and $\lambda': \mathcal{M} \rightarrow \mathcal{H}'$ satisfying (i) and (ii) with (\mathcal{H}, λ) replaced by (\mathcal{H}', λ') then there exists a unitary operator $W: \mathcal{H} \rightarrow \mathcal{H}'$ such that $W\lambda = \lambda'$;
- (iv) $\lambda((X(\sigma), u)) = \lambda(X^\sigma(\sigma \cup \delta), u)$ for all $(X(\sigma), u) \in \mathcal{M}$ and $\delta \in \Gamma_0$.

Proof. (i), (ii) and (iii) are immediate from Proposition 2.1 and the G.N.S. principle. (See, for example, Proposition 15.4, [P]). By (2.3) and (2.4) we have

$$\begin{aligned} L_T((X(\sigma), u), (X(\sigma), u)) &= L_T((X(\sigma), u), (X^\sigma(\sigma \cup \delta), u)) \\ &= L_T((X^\sigma(\sigma \cup \delta), u), (X^\sigma(\sigma \cup \delta), u)) \end{aligned}$$

and hence by (i) in the proposition

$$\begin{aligned} \|\lambda(X(\sigma), u) - \lambda(X^\sigma(\sigma \cup \delta), u)\|^2 &= \|\lambda(X(\sigma), u)\|^2 + \|\lambda(X^\sigma(\sigma \cup \delta), u)\|^2 \\ &\quad - 2 \operatorname{Re} \langle \lambda(X(\sigma), u), \lambda(X^\sigma(\sigma \cup \delta), u) \rangle = 0. \quad \blacksquare \end{aligned}$$

Remark. When $\sigma = \{t_1, t_2, \dots, t_n\}$ is fixed it is a consequence of (i) in Proposition 2.2 that $\lambda((X_{t_1}, X_{t_2}, \dots, X_{t_n}), u)$ is multilinear on $\mathcal{A}_{t_1} \times \dots \times \mathcal{A}_{t_n} \times \mathcal{H}_0$.

PROPOSITION 2.3.

In Proposition 2.2 let \mathcal{H}_t be the closed linear span of the set $\{\lambda(X(\sigma), u) | (X(\sigma), u) \in \mathcal{M}_t\}$ where \mathcal{M}_t is defined by (2.1) and (2.2). Then $\{\mathcal{H}_t, t \geq 0\}$ is an increasing family of subspaces of \mathcal{H} and the map $V: u \rightarrow \lambda(I_0, u)$ is a unitary operator from \mathcal{H}_0 to \mathcal{H}_0 .

Proof. Let $0 \leq s < t < \infty$. Suppose $\sigma = \{s, s_2, \dots, s_m\}$. Then by property (iv) in Proposition 2.2 we have

$$\lambda((X_s, X_{s_2}, \dots, X_{s_m}), u) = \lambda((I_t, X_s, X_{s_2}, \dots, X_{s_m}), u)$$

and the right hand side belongs to \mathcal{H}_t by definition. This proves the first part. To prove the second part we first observe that

$$\langle \lambda(I_0, u), \lambda(I_0, v) \rangle_{\mathcal{H}} = \langle u, v \rangle_{\mathcal{H}_0}.$$

Thus V is an isometry from \mathcal{H}_0 into \mathcal{H}_0 . Furthermore (2.3) implies

$$\begin{aligned} & \|\lambda(X_0, u) - \lambda(I_0, X_0 u)\|^2 \\ &= L_T((X_0, u), (X_0, u)) + L_T((I_0, X_0 u), (I_0, X_0 u)) \\ &\quad - 2\operatorname{Re} L_T((X_0, u), (I_0, X_0 u)) \\ &= \langle u, X_0^* X_0 u \rangle + \langle X_0 u, X_0 u \rangle \\ &\quad - 2\operatorname{Re} \langle u, X_0^* (X_0 u) \rangle = 0. \end{aligned}$$

■

For any Hilbert space \mathcal{H} we denote by $\mathcal{B}(\mathcal{H})$ the C^* algebra of all bounded operators on \mathcal{H} .

PROPOSITION 2.4.

Let \mathcal{H} , \mathcal{H}_t , λ , V be as in Proposition 2.3. Then there exists a unique $*$ unital homomorphism $j_t^0: \mathcal{A}_t \rightarrow \mathcal{B}(\mathcal{H}_t)$ for every $t \geq 0$ satisfying the relations:

$$j_t^0(Y) \lambda((X_t, X_{t_2}, \dots, X_{t_n}), u) = \lambda((YX_t, X_{t_2}, \dots, X_{t_n}), u) \quad (2.7)$$

for all $Y \in \mathcal{A}_t$, $t > t_2 > \dots > t_n = 0$, $u \in \mathcal{H}_0$. Furthermore

$$V^* j_0^0(X) V = X \quad \text{for all } X \in \mathcal{A}_0.$$

Proof. Let $Y \in \mathcal{A}_t$ be unitary. By (2.3) and the fact that $\{T(s, t)\}$ is a family of transition operators it follows immediately that

$$\begin{aligned} & \langle \lambda((YX_t, X_{t_2}, \dots, X_{t_n}), u), \lambda((YZ_t, Z_{t_2}, \dots, Z_{t_n}), v) \rangle \\ &= L_T(((YX_t, X_{t_2}, \dots, X_{t_n}), u), ((YZ_t, Z_{t_2}, \dots, Z_{t_n}), v)) \\ &= L_T(((X_t, X_{t_2}, \dots, X_{t_n}), u), ((Z_t, Z_{t_2}, \dots, Z_{t_n}), v)) \\ &= \langle \lambda((X_t, X_{t_2}, \dots, X_{t_n}), u), \lambda((Z_t, Z_{t_2}, \dots, Z_{t_n}), v) \rangle \end{aligned}$$

for all $X_t, Y_t \in \mathcal{A}_t, X_{t_1}, Y_{t_1} \in \mathcal{A}_{t_1}, u, v \in \mathcal{X}_0$. This together with property (iv) of Proposition 2.2 implies that

$$\begin{aligned} & \langle \lambda(YX_t, X_{t_2}, \dots, X_{t_n}), u \rangle, \lambda(YZ_t, Z_{t_1}, Z_{t_2}, \dots, Z_{t_n}), v \rangle \\ &= \langle \lambda(X_t, X_{t_2}, \dots, X_{t_n}), u \rangle, \lambda(Z_t, Z_{t_1}, Z_{t_2}, \dots, Z_{t_n}), v \rangle \end{aligned}$$

Thus for any unitary Y in \mathcal{A}_t there exists a unitary operator $j_t^0(Y)$ in \mathcal{H}_t satisfying (2.7). If Y_1, Y_2 are unitary elements in \mathcal{A}_t it follows from the definitions that $j_t^0(Y_1)j_t^0(Y_2) = j_t^0(Y_1 Y_2)$. Since $\lambda((X_t, X_{t_1}, \dots, X_{t_n}), u)$ is linear in the variable X_t and any element in \mathcal{A}_t is a linear combination of at most four unitary elements in \mathcal{A}_t it follows that $j_t^0(\cdot)$ defined for unitary elements extends linearly to \mathcal{A}_t as a $*$ unital homomorphism from \mathcal{A}_t into $\mathcal{B}(\mathcal{H}_t)$. The uniqueness part is obvious. To prove the last part we have to only note that by the definition of V in Proposition 2.3 and the last part of its proof

$$\begin{aligned} j_0^0(X)Vu &= j_0^0(X)\lambda(I_0, u) = \lambda(X, u) \\ &= \lambda(I_0, Xu) = VXu \end{aligned}$$

for all $u \in \mathcal{X}_0$. ■

Theorem 2.5. Let \mathcal{A}_t be a unital C^* algebra of operators in a Hilbert space \mathcal{H}_t for every $t \geq 0$ and let $T(s, t): \mathcal{A}_t \rightarrow \mathcal{A}_s, s \leq t$ be a family of transition operators. Then there exists a Hilbert space \mathcal{H} , an increasing family $\{F(t), t \geq 0\}$ of projection operators on \mathcal{H} , a family of contractive $*$ homomorphisms $j_t: \mathcal{A}_t \rightarrow \mathcal{B}(\mathcal{H}), t \geq 0$ and a unitary isomorphism V from \mathcal{X}_0 onto the range of $F(0)$ satisfying the following:

- (i) $j_t(I_t) = F(t), I_t$ being the identity operator in \mathcal{H}_t ;
- (ii) for any $0 \leq s \leq t < \infty, X \in \mathcal{A}_t$

$$F(s)j_t(X)F(s) = j_s(T(s, t)(X));$$

- (iii) the set $\{j_{t_1}(X_1) \cdots j_{t_n}(X_n) Vu, t_1 > t_2 > \cdots > t_n = 0, X_i \in \mathcal{A}_{t_i}$ for each $i, n = 1, 2, \dots, u \in \mathcal{X}_0\}$ is total in \mathcal{H} ;
- (iv) $j_0(X)V = VX$ for all $X \in \mathcal{A}_0$ and for any $u, v \in \mathcal{X}_0, \sigma = \{s_1 > s_2 > \cdots > s_m = 0\}, \delta = \{t_1 > t_2 > \cdots > t_n = 0\}$,

$$\begin{aligned} & X_i \in \mathcal{A}_{s_i}, Y_j \in \mathcal{A}_{t_j}, i = 1, 2, \dots, m, j = 1, 2, \dots, n \\ & \langle j_{s_1}(X_1)j_{s_2}(X_2) \cdots j_{s_m}(X_m) Vu, j_{t_1}(Y_1)j_{t_2}(Y_2) \cdots j_{t_n}(Y_n) Vv \rangle \\ &= L_T((X(\sigma), u), (Y(\delta), v)), \end{aligned}$$

where L_T is given by (2.3) and (2.4).

Proof. Let $\mathcal{H}, \mathcal{H}_t, \lambda, V$ and j_t^0 be as in Proposition 2.4. Define $F(t)$ to be the projection on the subspace \mathcal{H}_t . By Proposition 2.3, $F(t)$ is increasing in t . Define, for any $X \in \mathcal{A}_t$, the operator $j_t(X)$ in \mathcal{H} by

$$j_t(X) = j_t^0(X)F(t) \text{ for any } t \geq 0.$$

Since j_t^0 is a $*$ unital homomorphism from \mathcal{A}_t into $\mathcal{B}(\mathcal{H}_t)$ and $F(t)$ is a projection it follows that $\|j_t(X)\| \leq \|X\|$ and $j_t(I_t) = F(t)$. To check that $j_t(X)j_t(Y) = j_t(XY)$ it is

enough to verify this on vectors of the form $\lambda((X_1, X_{t_1}, \dots, X_{t_n}), u)$. This is immediate from (2.7). Since $j_t^0(X)F(t) = F(t)j_t^0(X)F(t)$ it follows that $j_t(X)^* = j_t(X^*)$.

To prove (ii) it is enough to check that, for $s < t$,

$$\begin{aligned} &\langle \lambda((X_s, X_{s_2}, \dots, X_{s_m}), u), j_t^0(X)\lambda((Y_s, Y_{s_2}, \dots, Y_{s_m}), v) \rangle = \\ &\quad \langle \lambda((X_s, X_{s_2}, \dots, X_{s_m}), u), \lambda((T(s, t)(X) Y_s, Y_{s_2}, \dots, Y_{s_m}), v) \rangle \end{aligned}$$

for all $X \in \mathcal{A}$. By definitions the left hand side is equal to

$$\langle \lambda((I_t, X_s, X_{s_2}, \dots, X_{s_m}), u), \lambda((X, Y_s, Y_{s_2}, \dots, Y_{s_m}), v) \rangle$$

which, by property (i) in Proposition 2.2 and 2.3, is equal to the right hand side.

(iii) is just a restatement of property (ii) in Proposition 2.2 because

$$j_{t_1}(X_1) \cdots j_{t_n}(X_n) Vu = \lambda(X(\sigma), u)$$

with $\sigma = \{t_1, t_2, \dots, t_n\}$.

The first part of (iv) is contained in the last part of Proposition 2.4. The remaining part of (iv) follows from property (i) in Proposition 2.2. ■

Remark. It is interesting to compare the properties of $\{F(t)\}$ and $\{j_t\}$ in Theorem 2.5 with (1.2)–(1.4) in the case of classical Markov processes. This motivates the following definition: suppose $\mathcal{A}_t, \mathcal{X}_t$ and $T(s, t), s \leq t$ are as in Theorem 2.5. Then any quadruple $(\mathcal{H}, F, \{j_t\}, V)$ consisting of a Hilbert space \mathcal{H} , an increasing family $\{F(t)\}$ of projections in \mathcal{H} , contractive $*$ homomorphisms j_t from \mathcal{A}_t into $\mathcal{B}(\mathcal{H})$ and a unitary isomorphism V from \mathcal{X}_0 onto the range of $F(0)$ is called a *conservative Markov flow* with transition operators $T(\cdot, \cdot)$ if

$$j_t(I_t) = F(t), \quad F(s)j_t(X)F(s) = j_s(T(s, t)(X)) \text{ for } 0 \leq s \leq t < \infty$$

and $j_0(X)V = VX$ for all $X \in \mathcal{A}_0$, the flow is said to be *minimal* if, in addition, property (iii) of Theorem 2.5 holds. Two such minimal conservative Markov flows $(\mathcal{H}, F, \{j_t\}, V)$ and $(\mathcal{H}', F', \{j'_t\}, V')$ with the same transition operators $T(\cdot, \cdot)$ are called *equivalent* if there exists a unitary isomorphism $W: \mathcal{H} \rightarrow \mathcal{H}'$ such that

$$WF(t)W^{-1} = F'(t), \quad Wj_t(X)W^{-1} = j'_t(X), \quad WV = V'$$

for all $t \geq 0, X \in \mathcal{A}_t$ [BP], [M].

We shall establish soon that upto equivalence the minimal Markov flow constructed in Theorem 2.5 is unique.

PROPOSITION 2.6.

Let $(\mathcal{H}, F, \{j_t\}, V)$ be a minimal conservative Markov flow with transition operators $T(\cdot, \cdot)$ then the following hold:

(i) Let $0 \leq t_1 < t_2 < t_3 < \infty$. Then for any $X_i \in \mathcal{A}_{t_i}, i = 1, 2, 3$

$$j_{t_1}(X_1)j_{t_2}(X_2)j_{t_3}(X_3) = \begin{cases} j_{t_1}(X_1)T(t_1, t_2)(X_2)j_{t_3}(X_3) & \text{if } t_1 \geq t_3 \\ j_{t_1}(X_1)j_{t_3}(T(t_3, t_2)(X_2)X_3) & \text{if } t_1 < t_3 \end{cases}$$

(ii) Let \mathcal{N} be the set of all pairs of sequences of the form $(t_1, t_2, \dots, t_n; X_1, X_2, \dots, X_n)$ where $0 \leq t_1, t_2, \dots, t_n < \infty$, $X_i \in \mathcal{A}_{t_i}$, $i = 1, 2, \dots, n$, $n = 1, 2, \dots$. Then there exists a map $\alpha: \mathcal{N} \rightarrow \mathcal{A}_0$ independent of the Markov flow such that

$$F(0)j_{t_1}(X_1)j_{t_2}(X_2) \cdots j_{t_n}(X_n)F(0) = j_0(\alpha(\mathbf{t}, \mathbf{X})) \quad (2.8)$$

for all $(\mathbf{t}, \mathbf{X}) = (t_1, t_2, \dots, t_n; X_1, X_2, \dots, X_n) \in \mathcal{N}$.

Proof. Let t_1, t_2, t_3 be as in (i) and $t_1 \geq t_3$. Then

$$\begin{aligned} & j_{t_1}(X_1)j_{t_2}(X_2)j_{t_3}(X_3) \\ &= j_{t_1}(X_1)F(t_1)j_{t_2}(X_2)F(t_1)j_{t_3}(X_3) \\ &= j_{t_1}(X_1)j_{t_1}(T(t_1, t_2)(X_2))j_{t_3}(X_3) \\ &= j_{t_1}(X_1 T(t_1, t_2)(X_2))j_{t_3}(X_3), \end{aligned}$$

which proves the first part of (i). Its second part is proved in the same manner.

To prove (ii) observe that

$$\begin{aligned} & F(0)j_{t_1}(X_1)j_{t_2}(X_2) \cdots j_{t_n}(X_n)F(0) \\ &= j_0(I_0)j_{t_1}(X_1)j_{t_2}(X_2) \cdots j_{t_n}(X_n)j_0(I_0). \end{aligned} \quad (2.9)$$

Without loss of generality assume that $0 < t_1 < t_2 < \dots < t_{k-1} > t_k$. Then by (i) the product $j_{t_{k-2}}(X_{k-2})j_{t_{k-1}}(X_{k-1})j_{t_k}(X_k)$ can be reduced to a product of size 2 of the form $j_{t_{k-2}}(X'_{k-2})j_{t_k}(X_k)$ or $j_{t_{k-2}}(X_{k-2})j_{t_k}(X'_k)$ where the primed operators depend only on (\mathbf{t}, \mathbf{X}) and $T(\cdot, \cdot)$ and not on the particular flow under consideration. Thus the n -fold product between the two $j_0(I_0)$'s on the right hand side of (2.9) can be reduced to an $(n-1)$ -fold product. A successive reduction of the sequence $(0, t_1, t_2, \dots, t_n, 0; I_0, X_1, X_2, \dots, X_n, I_0)$ applying (i) yields in the end an element $\alpha(\mathbf{t}, \mathbf{X})$ satisfying (2.8). ■

Theorem 2.7. Let $\mathcal{A}_t, \mathcal{X}_t, T(s, t)$, $0 \leq s \leq t < \infty$ be as in Theorem 2.5. Then any two minimal conservative Markov flows with transition operators $T(\cdot, \cdot)$ are equivalent.

Proof. Let $(\mathcal{A}, F, \{j_t\}, V)$ and $(\mathcal{A}', F', \{j'_t\}, V')$ be two Markov flows satisfying the conditions of the theorem. Suppose that $s_1 > s_2 > \dots > s_m = 0$, $t_1 > t_2 > \dots > t_n = 0$, $X_i \in \mathcal{A}_{s_i}$, $Y_j \in \mathcal{A}_{t_j}$, $i = 1, 2, \dots, m, j = 1, 2, \dots, n$. Consider $(\mathbf{r}, \mathbf{Z}) \in \mathcal{N}$ (where \mathcal{N} is as in Proposition 2.6) defined by

$$\begin{aligned} \mathbf{r} &= (s_m, s_{m-1}, \dots, s_1, t_1, t_2, \dots, t_n), \\ \mathbf{Z} &= (X_m^*, X_{m-1}^*, \dots, X_1^*, Y_1, Y_2, \dots, Y_n). \end{aligned}$$

Since $s_m = t_n = 0$ it follows from Proposition 2.6 that there exists $\alpha(\mathbf{r}, \mathbf{Z}) \in \mathcal{A}_0$ such that

$$\begin{aligned} & j_{s_m}(X_m^*)j_{s_{m-1}}(X_{m-1}^*) \cdots j_{s_1}(X_1^*)j_{t_1}(Y_1) \cdots j_{t_n}(Y_n) = j_0(\alpha(\mathbf{r}, \mathbf{Z})), \\ & j'_{s_m}(X_m^*)j'_{s_{m-1}}(X_{m-1}^*) \cdots j'_{s_1}(X_1^*)j'_{t_1}(Y_1) \cdots j'_{t_n}(Y_n) = j'_0(\alpha(\mathbf{r}, \mathbf{Z})). \end{aligned}$$

Thus for any $u, v \in \mathcal{X}_0$ we have

$$\begin{aligned} & \langle j_{s_1}(X_1) \cdots j_{s_m}(X_m) V u, j_{t_1}(Y_1) \cdots j_{t_n}(Y_n) V v \rangle \\ & \langle j'_{s_1}(X_1) \cdots j'_{s_m}(X_m) V' u, j'_{t_1}(Y_1) \cdots j'_{t_n}(Y_n) V' v \rangle \\ & = \langle u, \alpha(r, Z) v \rangle. \end{aligned}$$

From the minimality of the two flows it follows that \mathcal{X} and \mathcal{X}' are spanned by vectors of the form $j_{t_1}(Y_1) \cdots j_{t_n}(Y_n) V u$ and $j'_{t_1}(Y_1) \cdots j'_{t_n}(Y_n) V' u$ respectively. Hence there exists a unitary isomorphism $W: \mathcal{X} \rightarrow \mathcal{X}'$ such that

$$W j_{t_1}(Y_1) \cdots j_{t_n}(Y_n) V u = j'_{t_1}(Y_1) \cdots j'_{t_n}(Y_n) V' u$$

for all $u \in \mathcal{X}_0$, $t_1 > t_2 > \cdots > t_n = 0$, $Y_i \in \mathcal{A}_{t_i}$, $i = 1, 2, \dots, n$. That W is the required isomorphism implementing the equivalence of the two flows is immediate. \blacksquare

Remark. Let $(\mathcal{X}, F, \{j_t\}, V)$ be a minimal conservative Markov flow with transition operators $T(\cdot, \cdot)$. Denote by \mathcal{B} and \mathcal{B}_t respectively the C^* algebras generated by $\{j_s(X), X \in \mathcal{A}_s, 0 \leq s < \infty\}$ and $\{j_s(X), X \in \mathcal{A}_s, 0 \leq s \leq t\}$. By the same arguments as in the proof of Proposition 2.6 it is easy to see that for $t_i \geq s$, $i = 1, 2, \dots, n$ an expression of the form $F(s) j_{t_1}(X_1) \cdots j_{t_n}(X_n) F(s)$ can be expressed as $j_s(\alpha_s(t, X))$ where $\alpha_s(t, X) \in \mathcal{A}_s$. In particular the map E_{s_1} defined by

$$E_{s_1}(Z) = F(s) Z F(s), \quad Z \in \mathcal{B}$$

maps \mathcal{B} onto \mathcal{B}_{s_1} . We may call E_{s_1} the *conditional expectation map* from \mathcal{B} onto \mathcal{B}_{s_1} . If ρ_0 is a state on \mathcal{A}_0 then a state ρ on \mathcal{B} is uniquely determined by

$$\rho(Z) = \rho_0(V^* F(0) Z F(0) V), \quad Z \in \mathcal{B}.$$

It is legitimate to call the filtered quantum probability space $(\mathcal{B}, \mathcal{B}_t, \rho)$ the Markov process with initial state ρ_0 and transition operators $T(\cdot, \cdot)$.

Let \mathcal{Z}_t denote the centre of \mathcal{A}_t for each t . It is possible that $T(s, t)$ may not map \mathcal{Z}_t into \mathcal{Z}_s . In the minimal flow with transition operators $T(\cdot, \cdot)$, the operators $\{j_t(Z), Z \in \mathcal{Z}_t, t \geq 0\}$ need not be a commutative family. However, by following an idea in Bhat [B], we shall modify the construction in Proposition 2.4 in order to arrive at a family of $*$ unital homomorphisms $k_t: \mathcal{Z}_t \rightarrow \mathcal{B}(\mathcal{X})$ so that $\{k_t(Z), Z \in \mathcal{Z}_t, t \geq 0\}$ is a commutative family and $j_t(Z)$ is obtained from $k_t(Z)$ by a conditional expectation.

Theorem 2.8. *Let $(\mathcal{X}, F, \{j_t\}, V)$ be as in Theorem 2.5. Then there exists a unique $*$ unital homomorphism $k_t: \mathcal{Z}_t \rightarrow \mathcal{B}(\mathcal{X})$ satisfying the following:*

(i) for any $t_1 > t_2 > \cdots > t_n = 0$, $X_i \in \mathcal{A}_{t_i}$, $i = 1, 2, \dots, n$, $Z \in \mathcal{Z}_{t_1}$ and $u \in \mathcal{X}_0$

$$k_t(Z) \lambda((X_{t_1}, X_{t_2}, \dots, X_{t_n}), u) = \begin{cases} \lambda((X_{t_1}, X_{t_2}, \dots, X_{t_{i-1}}, Z X_{t_i}, X_{t_{i+1}}, \dots, X_{t_n}), u) & \text{if } t = t_i \text{ for some } i \\ \lambda((Z, X_{t_1}, \dots, X_{t_n}), u) & \text{if } t > t_1, \\ \lambda((X_{t_1}, X_{t_2}, \dots, X_{t_{i-1}}, Z, X_{t_i}, \dots, X_{t_n}), u) & \text{if } t_{i-1} > t > t_i \text{ for some } i; \end{cases} \quad (2.10)$$

(ii) the family $\{k_t(Z), Z \in \mathcal{X}_t, t \geq 0\}$ is commutative;

(iii) $j_t(Z) = F(t)k_t(Z)F(t)$ for all $t \geq 0, Z \in \mathcal{X}_t$.

Proof. As in the proof of Proposition 2.4 consider a unitary element $Z \in \mathcal{X}_t$. Suppose $t = t_i$ for some $i = 1, 2, \dots, n$. For any $X_{t_i}, Y_{t_i} \in \mathcal{X}_{t_i}, i = 1, 2, \dots, n$ we have

$$\begin{aligned} &\langle \lambda((X_{t_1}, X_{t_2}, \dots, X_{t_{i-1}}, ZX_{t_i}, X_{t_{i+1}}, \dots, X_{t_n}), u), \\ &\lambda((Y_{t_1}, Y_{t_2}, \dots, Y_{t_{i-1}}, ZY_{t_i}, Y_{t_{i+1}}, \dots, Y_{t_n}), v) \rangle = \\ &\langle u, X_{t_n}^* (\dots X_{t_i}^* Z^* T(t_i, t_{i-1}) (\dots (X_{t_2}^* T(t_2, t_1) (X_{t_1}^* Y_{t_1}) Y_{t_2}) \dots) ZY_{t_i} \dots) Y_{t_n} v \rangle. \end{aligned}$$

Since Z and $Z^* \in \mathcal{X}_{t_i}$ and $Z^*Z = 1$ it follows that the right hand side is independent of Z . The same argument in the remaining cases together with the Chapman-Kolmogorov equations for $T(\cdot, \cdot)$ and (iv) in Proposition 2.2 imply that $k_t(Z)$ defined by (2.10) on elements of the form $\lambda(X(\sigma), u)$ is scalar product preserving. Hence $k_t(Z)$ extends to a unitary operator on \mathcal{H} . Furthermore for any two unitary elements $Z, Z' \in \mathcal{X}_t$, we have $k_t(Z)k_t(Z') = k_t(ZZ')$. Once again by (iv) in Proposition 2.2, $k_t(I_t)$ is the identity operator in \mathcal{H} . Exactly as in the proof of Proposition 2.4 we extend $k_t(\cdot)$ to a $*$ unital homomorphism from \mathcal{X}_t into $\mathcal{B}(\mathcal{H})$. This proves (i).

If $t \neq t', Z \in \mathcal{X}_t, Z' \in \mathcal{X}_{t'}$, it follows from (2.10) by straightforward verification that

$$k_t(Z)k_{t'}(Z')\lambda(X(\sigma), u) = k_{t'}(Z')k_t(Z)\lambda(X(\sigma), u)$$

where $\sigma = \{t_1 > t_2 > \dots > t_n = 0\}$. This proves (ii).

When $t = t_1 > t_2 > \dots > t_n, X_{t_i}, Y_{t_i} \in \mathcal{X}_{t_i}, u, v \in \mathcal{X}_0$ we have

$$\begin{aligned} &\langle \lambda((X_{t_1}, X_{t_2}, \dots, X_{t_n}), u), k_t(Z)\lambda((Y_{t_1}, Y_{t_2}, \dots, Y_{t_n}), v) \rangle \\ &= \langle \lambda((X_{t_1}, X_{t_2}, \dots, X_{t_n}), u), \lambda((ZY_{t_1}, Y_{t_2}, \dots, Y_{t_n}), v) \rangle \\ &= \langle \lambda((X_{t_1}, X_{t_2}, \dots, X_{t_n}), u), j_t(Z)\lambda((Y_{t_1}, Y_{t_2}, \dots, Y_{t_n}), v) \rangle. \end{aligned}$$

Since vectors of the form $\lambda((X_{t_1}, X_{t_2}, \dots, X_{t_n}), u)$ span the range \mathcal{H}_t of $F(t)$, property (iii) is immediate. Uniqueness of $\{k_t\}$ follows from the minimality of $\{j_t\}$ and property (i). ■

References

[AFL] Accardi L, Frigerio A and Lewis J T, Quantum stochastic processes, *Publ. RIMS, Kyoto Univ.* **18** (1982) 97–133
 [B] Bhat B V, Rajarama, *Markov dilations of nonconservative quantum dynamical semigroups and a quantum boundary theory*, Ph.D. Thesis (submitted to Indian Statistical Institute, New Delhi) (1993)
 [BP] Bhat B V, Rajarama and Parthasarathy K R, *Markov dilations of nonconservative quantum dynamical semigroups and a quantum boundary theory*, Indian Statistical Institute, New Delhi preprint (1993)
 [E] Emch G G, Minimal dilations of CP flows, *C* algebras and applications to physics*, Springer LNM **650** (1978) 156–159
 [M] Meyer P A, Processus de Markov non-commutatifs d'après Bhat-Parthasarathy (to appear in *Séminaire de Probabilités* (ed.) Meyer Azéma and Yor) (Strasbourg preprint) (1993)
 [P] Parthasarathy K R, An Introduction to Quantum Stochastic Calculus, *Monographs in Mathematics* (Basel: Birkhauser Verlag) (1992)
 [S] Sauvageot J L, Markov quantum semigroups admit covariant Markov *C** dilations, *Commun. Math. Phys.* **106** (1986) 91–103
 [Vi-S] Vincent-Smith G F, Dilations of a dissipative quantum dynamical system to a quantum Markov process, *Proc. Lond. Math. Soc.* (3) **49** (1984) 58–72