

Structure of a code related to $\text{Sp}(4, q)$, q even

N S NARASIMHA SASTRY¹ and R P SHUKLA²

¹Stat.-Math. Unit, Indian Statistical Institute, 8th Mile Mysore Road, R.V. College Post, Bangalore 560 059, India

²Department of Mathematics, University of Allahabad, Allahabad 211 002, India
 Email: nsastry@isibang.ac.in; rps@mri.ernet.in

MS received 13 February 2006

Abstract. We determine the socle and the radical series of the binary code associated with a finite regular generalized quadrangle of even order, considered as a module for the commutator of each of the orthogonal subgroups in the corresponding symplectic group.

Keywords. Generalized quadrangle; symplectic group; socle layers; Steinberg module.

1. Introduction and statement of the result

Let $V(q) = \mathbb{F}_q^4$, $q = 2^n$, be the vector space of dimension four over \mathbb{F}_q endowed with a nondegenerate symplectic bilinear form \tilde{h} and $W(q)$ denote the incidence system with the set P of all one-dimensional subspaces of $V(q)$ as its *point-set*, the set L of all two-dimensional subspaces of $V(q)$ which are isotropic with respect to \tilde{h} as its *line-set* and symmetrized inclusion as the *incidence*. Then, $W(q)$ is a regular generalized quadrangle of order q (p. 37 of [12]). Further, elements of the symplectic group G defined by \tilde{h} act as incidence-preserving permutations on the sets P and L .

Let k be an algebraically closed extension field of \mathbb{F}_q . We denote by \mathcal{C} the image of the kG -module homomorphism from the permutation G -module k^L on L over k to the permutation G -module k^P on P over k , taking $l \in L$ to $\sum_{p \in l} p \in k^P$. The endomorphism of k^P taking each element of P to the ‘all-one’ vector $\mathbf{1} = \sum_{p \in P} p$ in k^P is a G -module homomorphism onto the trivial G -submodule $k\mathbf{1}$ of k^P . If Y_P is the kernel of the ‘augmentation map’ from k^P to k taking $p \in P$ to $1 \in k$, then

$$k^P = k\mathbf{1} \oplus Y_P.$$

Since every element of P is incident with $q + 1$ elements of L , it follows that $\mathbf{1} \in \mathcal{C}$ and

$$\mathcal{C} = k\mathbf{1} \oplus \overline{\mathcal{C}},$$

where $\overline{\mathcal{C}} = Y_P \cap \mathcal{C}$. The Loewy structure of the kG -module $\overline{\mathcal{C}}$ is determined in Theorem 2, p. 486 of [14]. Here, we determine the Loewy structure of $\overline{\mathcal{C}}$ as a $k\Omega(f)$ -module, where $\Omega(f)$ is the commutator subgroup of the orthogonal group $O(f) \subset G$ defined by a nondegenerate quadratic form f on $V(q)$ polarizing to \tilde{h} . That is, $\tilde{h}(x, y) = f(x + y) - f(x) - f(y)$ holds for all $x, y \in V(q)$. There are two such quadratic forms on $V(q)$, up to G -equivalence (see Theorem 6, p. 214 of [3]). They are distinguished by the presence

(hyperbolic case) or otherwise (elliptic case) of isotropic projective lines in the set $V_q(f)$ of zeroes of f in the projective 3-space $P(3, q)$ over \mathbb{F}_q ; that is, they correspond to the cases when the Witt index of f is 2 or 1 (§4.1, p. 18 of [3]). The subgroups $O(f)$ and $\Omega(f)$ of G are isomorphic to $(SL(2, q) \times SL(2, q)) \cdot 2$ and $SL(2, q) \times SL(2, q)$, respectively, when the Witt index of f is 2; and to $SL(2, q^2) \cdot 2$ and $SL(2, q^2)$, respectively, when the Witt index of f is 1. We also mention that $Sp(4, q)$ contains exactly 2 conjugacy classes of subgroups of each of the types $SL(2, q^2)$ and $SL(2, q) \times SL(2, q)$ (see Corollary, p. 247 of [6]).

To state our theorem, we describe the simple kG -modules. Let $N = \{0, 1, \dots, 2n - 1\}$, with addition taken modulo $2n$. Let $V = V(q) \otimes k \simeq k^4$ and extend the symplectic form h to V . Then G is the subgroup of the algebraic group $Sp(V) \simeq Sp(4, k)$ fixed by the n -th power of the Frobenius map σ (which is the ‘square-the-matrix-entries’ map on $GL(4, k)$). It is well-known that $Sp(V)$ has an endomorphism τ with $\tau^2 = \sigma$ (Theorem 28, p. 146 of [19]). For any non-negative integer i , we denote by V_i the $Sp(V)$ -module whose vector space structure is the same as that of V and an element g of $Sp(V)$ acts on V_i as $\tau^i(g)$ acts on V . For $I \subseteq N$, let V_I denote the kG -module $\otimes_{i \in I} V_i$ (with $V_\emptyset = k$). Then, by Steinberg’s tensor product theorem (§11 of [18]), $\{V_I : I \subseteq N\}$ is a complete set of inequivalent simple kG -modules.

Let $N_1 = \{0, 1, \dots, n - 1\}$, with addition taken modulo n . For $I \subseteq N$, define $I_e = \{i \in N_1 : 2i \in I\}$ and $I_o = \{i \in N_1 : 2i - 1 \in I\}$. For $n > 1$, we denote by \mathcal{N} the set of all subsets I of N which has no consecutive elements, that is, all I such that I_e and I_o are disjoint and if $i \in I_e$, then $i + 1 \notin I_o$. For each subset I of N_1 , the subset $N_1 \setminus (I \cup \{i + 1 | i \in I\})$ of N_1 will be called the *admissible complement* of I and will be denoted by I^{ac} . We observe that for $I \subseteq N_1$, the subset $K = \{2i : i \in I\} \cup \{2i - 1 : i \in I^{ac}\}$ is the unique maximal subset of N such that $K_e = I$ and $K_o = I^{ac}$. Also, $I^{ac} = \emptyset$ if $|I| \geq n - 1$ or if $N_1 \setminus I = \{i, j\}$, $i < j$ and $j \neq i + 1$. For each $m \in \{0, 1, \dots, n - 2, n\}$, let \mathcal{A}_m denote the set of all subsets I of N_1 such that $|I^{ac}| = m$. We observe that $\mathcal{A}_n = \{\emptyset\}$ and $\mathcal{A}_{n-1} = \emptyset$. For a module M admitting a chain of submodules $M = M_0 \supseteq M_1 \supseteq \dots \supseteq M_r$, we indicate the chain of factor modules $E_i = M_i/M_{i+1}$ as $E_{r-1} \setminus \dots \setminus E_0$ or $E_0/\dots/E_{r-1}$. For a kG -module M , we write M as $\text{rad}^0(M)$, denote by $\text{rad}^1(M)$ the *radical of M* (that is, the smallest submodule of M with semisimple quotient) and, for a positive integer i , define the i -th radical $\text{rad}^i(M)$ of M recursively as $\text{rad}^i(M) = \text{rad}^1(\text{rad}^{i-1}(M))$. Dually, we write $\text{soc}^0(M) = \{0\}$, denote by $\text{soc}^1(M)$ the *socle of M* (that is, the largest semisimple submodule of M) and, for a positive integer i , define the i -th *socle* $\text{soc}^i(M)$ of M recursively by $\text{soc}^i(M)/\text{soc}^{i-1}(M) = \text{soc}(M/\text{soc}^{i-1}(M))$. The *radical length* (respectively *socle length*) of M is the positive integer r such that $\text{rad}^r(M) = 0$ but $\text{rad}^{r-1}(M) \neq 0$ (respectively $\text{soc}^r(M) = M$ but $\text{soc}^{r-1}(M) \neq M$). We refer to $M/\text{rad}^1(M)$ as the *head of M* and write it as $\text{hd}(M)$.

Theorem 1.

(a) *Let $n = 1$. As $k\Omega(f)$ -modules,*

- (i) *when f is of Witt index 1, V_0 and V_1 are semisimple; \bar{C} is multiplicity-free and is isomorphic to the direct sum of V_0 and a $k\Omega(f)$ -module X with $\text{hd}(X) \simeq V_1$ and $\text{rad}(X) \simeq k$; and*
- (ii) *when f is of Witt index 2, \bar{C} is semisimple and each composition factor of \bar{C} appears with multiplicity one.*

(b) Let $n \geq 2$. As a $k\Omega(f)$ -module,

- (i) V_I is semisimple for each $I \in \mathcal{N}$ and \bar{C} is multiplicity-free;
- (ii) socle length of \bar{C} is $n+1$ and its j -th socle layer $\text{soc}^j(\bar{C})/\text{soc}^{j-1}(\bar{C})$, $1 \leq j \leq n+1$, is isomorphic to

$$\bigoplus_{I \in \mathcal{N} \text{ and } |I_o|=j-1} V_I;$$

(iii)

$$\bar{C} \simeq \bigoplus_{m \in \{0, 1, \dots, n-2, n\}} X_m \quad \text{and} \quad X_m = \bigoplus_{I \in \mathcal{A}_m} X_{m, I},$$

where $X_{m, I}$ is the unique indecomposable $k\Omega(f)$ -submodule of \bar{C} with head V_K with K denoting the unique element of \mathcal{N} such that $K_e = I$ and $K_o = I^{ac}$. The radical length of $X_{m, I}$ is $m+1$ and its j -th radical layer $\text{rad}^j(X_{m, I})/\text{rad}^{j+1}(X_{m, I})$, $0 \leq j \leq m$, is isomorphic to

$$\bigoplus_{J \in \mathcal{N}, J_e=I, J_o \subseteq I^{ac} \text{ and } |J_o|=m-j} V_J$$

and

$$(iv) \text{soc}^j(X_{m, I}) = \text{rad}^{m+1-j}(X_{m, I}), \quad 1 \leq j \leq m.$$

To prove this, we use the following:

Theorem 2 (Theorem 2 of [14]). *The radical series of \bar{C} as a kG -module has length $2n+1$. The radical layers are*

$$\text{rad}^j(\bar{C})/\text{rad}^{j+1}(\bar{C}) = \bigoplus_{I \in \mathcal{N} \text{ and } |I_e|-|I_o|+n=j} V_I, \quad 0 \leq j \leq 2n.$$

Moreover,

$$\text{soc}^j(\bar{C}) = \text{rad}^{2n+1-j}(\bar{C}).$$

The crucial observation is that if E is a section of \bar{C} which is a nonsplit extension of V_K by V_L , then E is a nonsplit $k\Omega(f)$ -module if and only if $L = K \cup \{2t-1\}$ for some $t \in N_1$ (see the first para of the proof of b(ii) in §3). Thus either each or none of the simple $k\Omega(f)$ -factors of V_L descends.

In §2, we study the structure of V_I as a $k\Omega(f)$ -module and prove Theorem 1 in §3. We also write the socle structure of \bar{C} as a $k\Omega(f)$ -module for $n = 1, 2, 3, 4$. We view this work as part of the program of determining the module structure of permutation modules for finite groups of Lie type (see [15,16]) and references therein for a few other cases considered in the literature.

2. V_I as $k\Omega(f)$ -modules

2.1

Throughout, we choose the nondegenerate symplectic bilinear form \hbar on $\mathbb{F}_q^4 \times \mathbb{F}_q^4$ to be

$$\hbar((x_1, x_2, x_3, x_4), (y_1, y_2, y_3, y_4)) = x_1y_4 + x_4y_1 + x_2y_3 + y_2x_3.$$

The quadratic forms f_e and f_h on \mathbb{F}_q^4 polarizing to \hbar and of respective Witt indices 1 and 2 are chosen to be

$$f_e(x_1, x_2, x_3, x_4) = x_1x_4 + \alpha^{q+1}x_2^2 + x_2x_3 + \alpha^{q+1}x_3^2; \text{ and}$$

$$f_h(x_1, x_2, x_3, x_4) = x_1x_4 + x_2x_3,$$

where α is an element of $\mathbb{F}_{q^2} \setminus \mathbb{F}_q$ such that $\alpha + \alpha^q = 1$.

This choice of f_e and f_h entails no loss of generality in view of the uniqueness of a non-degenerate symplectic form \hbar on $V(q)$, up to $GL(V(q))$ -equivalence (see Corollary 8.2, p. 587 of [11]); and the uniqueness of a quadratic form on $V(q)$ of a given Witt index and polarizing to \hbar , up to G -equivalence (see Theorem 6, p. 214 of [3] and p. 39 of [4]).

2.2 *The case f_e*

The polynomial $\alpha^{q+1}x_2^2 + x_2x_3 + \alpha^{q+1}x_3^2$ is irreducible over \mathbb{F}_q but factors as $(\alpha x_2 + \alpha^q x_3)(\alpha^q x_2 + \alpha x_3)$ over \mathbb{F}_{q^2} . So, the zero set \mathcal{E} of f_e in $P(3, q)$ is an elliptic quadric and the zero set $\widehat{\mathcal{E}}$ of f_e in $P(3, q^2)$ is a hyperbolic quadric. Since f_e polarizes to \hbar , \mathcal{E} is an ovoid in the generalized quadrangle $W(q)$ defined by \hbar (p. 51 of [2]).

(A) Recall that $\Omega(f_e) \simeq SL(2, q^2)$. We now interpret the $k\Omega(f_e)$ -modules V_i as $kSL(2, q^2)$ -modules. For this, we construct:

- (i) an isomorphism from $SL(2, q^2)$ to $\Omega(f_e)$, following Theorems 15.3.11 and 15.3.18 of [7]; and
- (ii) a graph automorphism τ of G , following pp. 58–60 of [5] (also see Chapter 12 of [20]).

(i)

(a) *The inversive planes $(P(1, q^2), \mathcal{L})$ and (\mathcal{E}, Φ) .* Recall that a subline of the projective line $P(1, q^2)$ over \mathbb{F}_{q^2} is the set of all one-dimensional subspaces of $\mathbb{F}_{q^2}^2$ generated by the \mathbb{F}_q -span of a fixed basis of $\mathbb{F}_{q^2}^2$. Equivalently, it is a nonsingular Hermitian variety $V_A = V(xA(x^{(q)})^t)$ in $P(1, q^2)$ defined by an equation of the form $a_1x_1^{q+1} + bx_1x_2^q + b^qx_1^qx_2 + a_2x_2^{q+1}$, where $x = (x_1, x_2) \in \mathbb{F}_{q^2}^2$, $x^{(q)} = (x_1^q, x_2^q)$ and $A = \begin{pmatrix} a_1 & b \\ b^q & a_2 \end{pmatrix}$ with $a_i \in \mathbb{F}_q, b \in \mathbb{F}_{q^2}, a_1a_2 \neq bb^q$ (Lemma 6.2, p. 138 of [8]). Let \mathcal{L} denote the set of all sublines of $P(1, q^2)$. Then, the incidence structure $(P(1, q^2), \mathcal{L})$, with symmetrized inclusion as incidence is an inversive plane of order q ; that is, a $3 - (q^2 + 1, q + 1, 1)$ design (p. 257 of [2]). This is isomorphic to the incidence system (\mathcal{E}, Φ) , where \mathcal{E} is as above, Φ is the set of all planes in $P(3, q)$ which are not tangent to \mathcal{E} and the incidence again is symmetrized containment (p. 257 of [2]).

(b) *An isomorphism between $(P(1, q^2), \mathcal{L})$ and (\mathcal{E}, Φ) .* To construct this, we follow Theorem 15.3.11, p. 21 of [7]. Let i be the injective map from $P(1, q^2)$ to $P(3, q^2)$ defined by

$$i(P(x_1, x_2)) = P(x_1^{q+1}, x_1x_2^q, x_1^qx_2, x_2^{q+1})$$

and ψ be the involutory projectivity of $P(3, q^2)$ defined by

$$\psi(P(x_1, x_2, x_3, x_4)) = P(x_1, \alpha x_2 + \alpha^q x_3, \alpha^q x_2 + \alpha x_3, x_4).$$

We identify $P(3, q)$ with its image in $P(3, q^2)$ under the map $\mathbb{F}_q a \mapsto \mathbb{F}_{q^2} a$; where $a = (a_1, a_2, a_3, a_4) \in \mathbb{F}_q^4$. We treat \hbar chosen in (2.1) also as a nondegenerate symplectic form on $\mathbb{F}_{q^2}^4$. For x in $P(3, q^2)$, we denote by $\hbar(x, -)$ the plane in $P(3, q^2)$ consisting of all points y of $P(3, q^2)$ such that $\hbar(x, y) = 0$. Since ψ stabilizes \hbar , $\psi(\hbar(x, -)) = \hbar(\psi(x), -)$ for each x in $P(3, q^2)$. Let j be the map from \mathcal{L} to Φ taking V_A to $\hbar(\psi(P(a')), -) \cap P(3, q)$, where $a' = (a_1, b, b^q, a_2) \in \mathbb{F}_{q^2}^4$. These planes are not tangent to \mathcal{E} because f_e takes the nonzero value $a_1 a_2 - b b^q$ on $\psi(P(a'))$ and so $\psi(P(a')) \notin \mathcal{E}$; it is known that the tangent planes to \mathcal{E} in $P(3, q)$ are all of the form $\hbar(x, -) \cap P(3, q)$, $x \in \mathcal{E}$; and the correspondence $\alpha \longleftrightarrow \hbar(\alpha, -) \cap P(3, q)$ between the points of $P(3, q)$ and the planes of $P(3, q)$ is incidence preserving and bijective. The map $\psi \circ i$ from $P(1, q^2)$ to \mathcal{E} and the map j from \mathcal{L} to Φ are both bijective and form an incidence preserving pair; that is, if $p \in V_A \in \mathcal{L}$, then $\psi \circ i(p) \in j(V_A) \in \Phi$.

(c) *The isomorphism from $SL(2, q^2)$ to $\Omega(f_e)$.* The full automorphism group of the incidence structure $(P(1, q^2), \mathcal{L})$ is $P\Gamma L(2, q^2) \simeq SL(2, q^2)\langle \xi \rangle$, where $\xi S \xi^{-1} = S^{(2)} = (s_{ij}^2)$ for all $S = (s_{ij}) \in SL(2, q^2)$ ((1), §4.1, p. 274 of [2]). We give an explicit isomorphism δ from its (unique) subgroup K isomorphic to $SL(2, q^2) \cdot 2$ into $Sp(4, q^2)$ such that $\psi(\text{im } \delta)\psi^{-1}$ equals the stabilizer $O(f_e)$ of \mathcal{E} in G (see Theorem 15.3.18, p. 27 of [7]).

For $S = (s_{ij}) \in SL(2, q^2)$, let P_S denote the projectivity of $P(1, q^2)$ taking $P(x)$ to $P(xS)$. Then, $\langle P_S : S \in SL(2, q^2) \rangle \simeq SL(2, q^2)$. If θ is the involutory projectivity of $P(1, q^2)$ taking $P(x_1, x_2) \mapsto P(x_1^q, x_2^q)$, then $\theta P_S \theta^{-1} = P_{S^{(q)}}$, where S is as above and $S^{(q)} = (s_{ij}^{(q)})$. So $\langle P_S : S \in SL(2, q^2) \rangle \langle \theta \rangle$ is the subgroup of $P\Gamma L(2, q^2)$ of index n .

Now P_S induces a bijection T_S on \mathcal{L} which takes V_A to $V_{SA((S^{(q)})^{-1})^{\text{tr}}}$. With the correspondence $A \leftrightarrow a'$, with a' as above, the map $A \mapsto SA((S^{(q)})^{-1})^{\text{tr}}$ can be written as $a' \mapsto a' R_S$, where

$$R_S = \begin{pmatrix} s_{11}s_{11}^q & s_{11}s_{21}^q & s_{21}s_{11}^q & s_{21}s_{21}^q \\ s_{11}s_{12}^q & s_{11}s_{22}^q & s_{21}s_{12}^q & s_{21}s_{22}^q \\ s_{12}s_{11}^q & s_{12}s_{21}^q & s_{22}s_{11}^q & s_{22}s_{21}^q \\ s_{12}s_{12}^q & s_{12}s_{22}^q & s_{22}s_{12}^q & s_{22}s_{22}^q \end{pmatrix} = S^{\text{tr}} \otimes (S^{(q)})^{\text{tr}}.$$

Let \mathcal{R}_S denote the projectivity of $P(3, q^2)$ taking $P(x_1, x_2, x_3, x_4)$ to $P((x_1, x_2, x_3, x_4)R_S)$. Note that ψ and \mathcal{R}_S both stabilize \hbar ; \mathcal{R}_S fixes the quadratic form $g(x_1, x_2, x_3, x_4) = x_1 x_4 + x_2 x_3$; ψ is a bijection between the varieties $V_{q^2}(g)$ and $V_{q^2}(f_e)$; and $\psi \mathcal{R}_S \psi^{-1}$ is a projectivity of $P(3, q)$ also. So, $\psi \mathcal{R}_S \psi^{-1}$ is in the stabilizer of \mathcal{E} in G . If \mathcal{R}_S is identity, then so is T_S . These facts imply that the map $S \mapsto \psi \mathcal{R}_S \psi^{-1}$ is a monomorphism from $SL(2, q^2)$ to $O(f_e)$. Now, the map $A \mapsto A^{(q)}$ can be written as $a' \mapsto a' \mathcal{R}$, where \mathcal{R} denotes the projectivity of $P(3, q^2)$ taking $P(x_1, x_2, x_3, x_4)$ to $P(x_1, x_3, x_2, x_4)$. Further, \mathcal{R} is in the stabilizer of \mathcal{E} in G , ψ and \mathcal{R} commute and $\mathcal{R} \mathcal{R}_S \mathcal{R}^{-1} = \mathcal{R}_{S^{(q)}}$ for each $S \in SL(2, q^2)$. So, $\langle \psi \mathcal{R}_S \psi^{-1} : S \in SL(2, q^2) \rangle \langle \mathcal{R} \rangle$ is a subgroup of $O(f_e)$ and is isomorphic to $SL_2(q^2) \cdot 2$. Now equality holds by order considerations and the isomorphism follows.

(ii)

We now describe a graph automorphism τ of $Sp(V)$, following pp. 58–60 of [5]. (The argument presented in loc. cit. constructs a graph automorphism for $G = Sp(V(q))$. However the arguments are valid for $Sp(V)$ also.) Let Q denote the nondegenerate quadratic

form on the exterior square $\Lambda^2 V$ of V defined by

$$Q(\sum_{1 \leq i < j \leq 4} \lambda_{ij} e_i \wedge e_j) = \lambda_{12} \lambda_{34} + \lambda_{13} \lambda_{24} + \lambda_{14} \lambda_{23}$$

(and whose zero set in $P(\Lambda^2 V)$ is the well-known *Klein quadric*). Let β denote the polarization of Q and $\gamma = e_1 \wedge e_4 + e_2 \wedge e_3$. Then the restriction of Q to the hyperplane $U = \{x \in \Lambda^2 V: \beta(x, \gamma) = 0\}$ of $\Lambda^2 V$ is a nondegenerate quadratic form; and the restriction of β to U is an alternating form with radical $k\gamma$. The alternating form $\bar{\beta}$ induced by β on $\bar{U} = U/k\gamma$ is nondegenerate. So the symplectic space $(\bar{U}, \bar{\beta})$ is isometric to (V, \hbar) . Let $\bar{p}: \bar{U} \rightarrow V$ be the isometric isomorphism induced by the linear map $p: U \rightarrow V$ defined by

$$\begin{aligned} p(e_1 \wedge e_2) &= e_1, p(e_1 \wedge e_3) = e_2, p(e_2 \wedge e_4) = e_3, \\ p(e_3 \wedge e_4) &= e_4, p(\gamma) = 0. \end{aligned}$$

Then the map taking $g \in \text{Sp}(V)$ to $\bar{p}(\overline{\Lambda^2(g)})\bar{p}^{-1} \in \text{Sp}(V)$ is a graph automorphism τ of $\text{Sp}(V)$ which, on restriction to G , gives a graph automorphism of G .

(B) Let W denote the standard two-dimensional simple $kSL(2, q^2)$ -module. For a non-negative integer i , let W_i denote, as in §1, the $kSL(2, q^2)$ -module whose underlying vector space structure is the same as that of W and the action of $g = (a_{l,m}) \in SL(2, q^2)$ on W_i is the usual action of $g^{(2^i)} = (a_{l,m}^{2^i})$ on W . For any subset I of N , let W_I denote the $kSL(2, q^2)$ -module $\otimes_{i \in I} W_i$. Then, by Steinberg's tensor product theorem, $\{W_I: I \subseteq N\}$ is a complete set of inequivalent simple $kSL(2, q^2)$ -modules. The following decomposition was suggested to the first-named author by Peter Sin.

Lemma 3. For $i \in N_1$,

$$V_{2i}|_{\Omega(f_e)} \simeq W_i \otimes W_{n+i}$$

and

$$V_{2i-1}|_{\Omega(f_e)} \simeq W_i \oplus W_{n+i}.$$

Proof. We prove this by using the above interpretation of V_i as $kSL(2, q^2)$ -modules. The group $H = \langle R_S: S \in SL(2, q^2) \rangle$, where R_S is as in (2.2A.i(c)) and $\Omega(f_e)$ are both isomorphic to $SL(2, q^2)$ and are conjugate in $\text{Sp}(4, k)$ (in fact in $\text{Sp}(4, q^2)$) (see the last paragraph of 2.2.A.i(c)). So, for $I \subseteq N$, V_I considered as a kH -module and V_I considered as a $k\Omega(f_e)$ -module are isomorphic $SL(2, q^2)$ -modules. Let (e_1, e_2, e_3, e_4) be the standard ordered basis for V and (v_1, v_2) be an ordered basis for W . For $S \in SL(2, q^2)$, $S \otimes S^{(q)}$ represents not only the action of S on V with respect to the ordered basis (e_1, e_2, e_3, e_4) of V but also the action of S on $W_0 \otimes W_n$ with respect to the ordered basis $(v_2 \otimes v_2, v_2 \otimes v_1, v_1 \otimes v_2, v_1 \otimes v_1)$ of $W_0 \otimes W_n$. So, $V_0|_{\Omega(f_e)} \simeq W_0 \otimes W_n$. Now, an application of τ^{2i} to V yields the first part of the lemma.

A simple calculation shows that R_S^τ leaves the subspaces $M^1 = ke_4 + ke_1$ and $M^2 = ke_2 + ke_3$ invariant. Further, the matrix representation of R_S^τ on M^1 with respect to the ordered basis (e_4, e_1) is S and it is $S^{(q)}$ on M^2 with respect to the ordered basis (e_3, e_2) . So, an application of τ^{2i-1} to V yields the second part of the lemma. ■

For $I \subseteq N_1$, let \mathcal{N}_I denote the set of $2^{|I|}$ subsets J of N of size $|I|$ such that, for each $t \in I$, only one of t and $n+t$ is in J . Let $\bar{I} = I \cup (n+I)$. Then, for $K \in \mathcal{N}$, Lemma 3 yields

$$V_K|_{\Omega(f_e)} \simeq \left(\bigoplus_{L \in \mathcal{N}_{K_0}} W_L \right) \otimes W_{\bar{K}_e}. \tag{1_e}$$

Notice that each irreducible component in (1_e) determines both the sets K_e and K_o . So we have the following:

COROLLARY 4

Let $K, K' \in \mathcal{N}$ be distinct. Then V_K and $V_{K'}$ are semisimple $k\Omega(f_e)$ -modules with no irreducible factors in common.

2.3 The case f_h

We now study V_I as a $k\Omega(f_h)$ -module. The zero set of f_h in $P(3, q)$ is a quadric of Witt index 2. Since f_h polarizes to \bar{h} , the projective lines contained in this quadric are isotropic with respect to \bar{h} . Further, $\Omega(f_h) = H_1H_2$, $[H_1, H_2] = H_1 \cap H_2 = \langle 1 \rangle$, where H_1 (respectively H_2) $\simeq SL(2, q)$ is the stabilizer of the subspace $ku_1 + ku_2$ (respectively $kv_1 + kv_2$) of V . Here $u_1 = e_1 + e_3, u_2 = e_2 + e_4, v_1 = e_1 + e_2$ and $v_2 = e_3 + e_4$. Further, the action of H_1 (respectively H_2) on $ku_1 + ku_2$ (respectively $kv_1 + kv_2$) with respect to its ordered basis (u_1, u_2) (respectively (v_1, v_2)) is the standard $SL(2, q)$ -action on k^2 .

Let M^1 and M^2 be two copies of the standard 2-dimensional simple $kSL(2, q)$ -module. For $j \in N_1$, define the j th-standard Frobenius twist M_j^i of M^i as in 2.2.B. For $J \subseteq N_1$, define M_J^i as $\otimes_{j \in J} M_j^i$. Further, for I and $J \subseteq N_1$, denote by $M_I^1 \# M_J^2$ the outer tensor product of M_I^1 and M_J^2 . That is, $M_I^1 \# M_J^2$ is the k -vector space $M_I^1 \otimes M_J^2$ with the action of $(h_1, h_2) \in SL(2, q) \times SL(2, q)$ on it given by

$$(h_1, h_2)(m_1 \otimes m_2) = h_1(m_1) \otimes h_2(m_2).$$

Then, $\{M_I^1 \# M_J^2 : I, J \subseteq N_1\}$ is a complete set of pairwise nonisomorphic simple $k(SL(2, q) \times SL(2, q))$ -modules (Theorem 9.14, p. 136 of [9]).

Lemma 5. For $i \in N_1$,

$$V_{2i}|_{\Omega(f_h)} \simeq M_i^1 \# M_i^2$$

and

$$V_{2i-1}|_{\Omega(f_h)} \simeq (M_i^1 \# k) \oplus (k \# M_i^2).$$

Proof. Let $h_i \in H_i$ be represented by the matrix $A_i \in SL(2, q)$ with respect to the ordered bases mentioned above. The matrix $A_1 \otimes A_2$ represents the action of $h = h_1h_2$ on V with respect to its ordered basis (e_1, e_2, e_3, e_4) as well as its action on $(ku_1 + ku_2) \# (kv_1 + kv_2)$ with respect to its ordered basis $(u_1 \otimes v_1, u_2 \otimes v_1, u_1 \otimes v_2, u_2 \otimes v_2)$. So, $V_0|_{\Omega(f_h)} \simeq M^1 \# M^2$. Now, an application of τ^{2i} to V yields the first part of the lemma.

A simple calculation shows that each of the subgroups H_i^τ stabilizes the subspaces $ke_2 + ke_3$ and $ke_1 + ke_4$. Further, the action of H_1^τ (respectively of H_2^τ) on $ke_2 + ke_3$ (respectively on $ke_1 + ke_4$) with respect to the ordered basis (e_2, e_3) (respectively, (e_1, e_4)) is equivalent

to the standard action of $SL(2, q)$ on k^2 and the action on $ke_1 + ke_4$ (respectively on $ke_2 + ke_3$) is trivial. So, $V_1|_{\Omega(f_h)} \simeq (M_1^1 \# k) \oplus (k \# M_1^2)$. An application of $\tau^{2(i-1)}$ to V_1 now yields the second part of the lemma. ■

For any subset I of N_1 , let \mathcal{M}_I denote the set of $2^{|I|}$ ordered partitions (A, B) of I . By Lemma 5, if $K \in \mathcal{N}$, then

$$\begin{aligned} V_K|_{\Omega(f_h)} &\simeq (M_{K_e}^1 \# M_{K_e}^2) \otimes (\oplus_{(A,B) \in \mathcal{M}_{K_0}} M_A^1 \# M_B^2) \\ &\simeq (\oplus_{(A,B) \in \mathcal{M}_{K_0}} M_{K_e \cup A}^1 \# M_{K_e \cup B}^2). \end{aligned} \tag{1_h}$$

Note that for $K \in \mathcal{N}$, K_e and K_o are disjoint. Further K is determined by each irreducible component in (1_h) . So, we have the following:

COROLLARY 6

Let $K, K' \in \mathcal{N}$ be distinct. Then V_K and $V_{K'}$ are semisimple $k\Omega(f_h)$ -modules with no irreducible factors in common.

2.4

The core of the proof of the theorem is in Lemmas 7 and 8 that we now state. Let $f \in \{f_e, f_h\}$ and assume that $n \geq 2$ for Lemmas 7 and 8.

Lemma 7. Let $J, K \in \mathcal{N}$. If the symmetric difference of J and K is not equal to $\{2t - 1\}$ for some $t \in N_1$, then $\text{Ext}_{k\Omega(f)}^1(V_J, V_K) = 0$.

Now, let $J = K \cup \{2t - 1\}$ and $2t - 1 \notin K$. Then, $2t - 2 \notin K$ and, by Theorem, p. 159 of [17], there exists a unique kG -module E , up to isomorphism, which is a nonsplit extension of V_K by V_J .

Lemma 8. $\text{soc}_{k\Omega(f)}(E|_{\Omega(f)}) \simeq V_K|_{\Omega(f)}$. In particular, as a $k\Omega(f)$ -module, E is a nonsplit extension of $V_K|_{\Omega(f)}$ by $V_J|_{\Omega(f)}$.

Proofs of Lemmas 7 and 8 for the cases $f = f_e$ and $f = f_h$ are given separately.

2.4.1 *The case $f = f_e$.* For easy reference, we collect some results due to Alperin and due to Sin. Consider the following condition on $I, J \subseteq N$:

$$I \cup J = (I \cap J) \cup \{r\} \text{ and neither } r \text{ nor } r - 1 \text{ is in } I \cap J. \tag{C_{I,J}}$$

Lemma 9.

- (a) For $I, J \subseteq N$, $\text{Ext}_{kSL(2,q^2)}^1(W_I, W_J)$ as well as $\text{Ext}_{kG}^1(V_I, V_J)$ are both k or both zero according as whether the condition $(C_{I,J})$ holds or not.
- (b) For $i \in N$, $W_i \otimes W_i$ is a uniserial module with composition factors $k \setminus W_{i+1} \setminus k$.

Proof. For (a), see Theorem 3, p. 221 of [1] and Theorem, p. 159 of [17]. For (b), see Lemma 4, p. 224 of [1]. ■

A typographical error about the assumptions on I and J in the statement of part (a) in Theorem 3, p. 221 of [1] has been corrected here (see p. 229 of [1]).

Proof of Lemma 7. In view of the isomorphism $\text{Ext}_{kG}^1(V_J, V_K) \simeq \text{Ext}_{kG}^1(V_{J \cup K}, V_{J \cap K})$, the biadditivity of the map $(M, N) \mapsto \text{Ext}^1(M, N)$ and Lemma 9(a) above, we only need to consider the case when $J = K \cup \{r\}$, $r \notin K$. In this case, the lemma follows from (1_e) and Lemma 9(a). ■

Proof of Lemma 8. From (1_e) ,

$$V_K |_{\Omega(f_e)} \simeq (\oplus_{L \in \mathcal{N}_{K_o}} W_L) \otimes W_{\overline{K_e}}$$

and

$$V_J |_{\Omega(f_e)} \simeq [\oplus_{L \in \mathcal{N}_{K_o}} (W_{L \cup \{t\}} \oplus W_{L \cup \{n+t\}})] \otimes W_{\overline{K_e}}.$$

We need to show that

$$\text{Hom}_{k\Omega(f_e)}(W_{L'} \otimes W_{\overline{K_e}}, E) = 0$$

for each $L' \in \mathcal{N}_{J_o}$. Let D_t denote the unique (up to isomorphism) kG -module which is a nonsplit extension of k by V_{2t-1} (see Lemma 9(a)). Then $E \simeq D_t \otimes V_K$ (Lemma 8, p. 490 of [14]). Since D_t is isomorphic to a submodule of the kG -module $V_{2t-2} \otimes V_{2t-2}$ (see Lemma 2(a), p. 161 of [17]), $D_t \otimes V_K$ embeds in $(V_{2t-2} \otimes V_{2t-2}) \otimes V_K$. Hence it is enough to prove that

$$\text{Hom}_{k\Omega(f_e)}(W_{L'} \otimes W_{\overline{K_e}}, V_{2t-2} \otimes V_{2t-2} \otimes V_K) = 0. \tag{*}$$

Let $L'' \in \mathcal{N}_{K_o}$ and $L' = L \cup \{r\}$ where $L \in \mathcal{N}_{K_o}$, $r = t$ or $n + t$. Since simple $k\Omega(f_e)$ -modules are self-dual, by Lemma 3,

$$\begin{aligned} & \text{Hom}_{k\Omega(f_e)}(W_{L'} \otimes W_{\overline{K_e}}, V_{2t-2} \otimes V_{2t-2} \otimes W_{L''} \otimes W_{\overline{K_e}}) \\ & \simeq \text{Hom}_{k\Omega(f_e)}(W_{L'} \otimes W_{\overline{K_e \cup \{t-1\}}}, W_{L''} \otimes W_{\overline{K_e \cup \{t-1\}}}) = 0. \end{aligned}$$

So $(*)$ holds and the lemma follows. ■

2.4.2 *The case $f = f_h$.* First we recall a useful result due to Jones (Theorem 3, p. 629 of [10]).

Lemma 10. Let R be a Dedekind domain, G_1 and G_2 be finite groups, $G = G_1 \times G_2$, $\Gamma_i = RG_i$, $i = 1, 2$, and $\Gamma = RG$. Let M_i and M'_i be (left) Γ_i -modules. Then, as R -modules,

$$\begin{aligned} \text{Ext}_{\Gamma}^1(M_1 \# M_2, M'_1 \# M'_2) & \simeq \text{Hom}_{\Gamma_1}(M_1, M'_1) \otimes_R \text{Ext}_{\Gamma_2}^1(M_2, M'_2) \\ & \oplus \text{Ext}_{\Gamma_1}^1(M_1, M'_1) \otimes_R \text{Hom}_{\Gamma_2}(M_2, M'_2) \end{aligned}$$

and

$$\text{Hom}_{\Gamma}(M_1 \# M_2, M'_1 \# M'_2) \simeq \text{Hom}_{\Gamma_1}(M_1, M'_1) \otimes_R \text{Hom}_{\Gamma_2}(M_2, M'_2).$$

Proof of Lemma 7. As in the case $f = f_e$, we need only to consider the case when $J = K \cup \{r\}$, where $r \notin K$. If $r = 2t$ for some $t \in N_1$, then (1_h) and Lemma 10 imply $\text{Ext}_{k\Omega(f_h)}^1(V_J, V_K) = 0$. ■

Proof of Lemma 8. From (1_h) ,

$$V_K |_{\Omega(f_h)} \simeq \bigoplus_{(A,B) \in \mathcal{M}_{K_0}} (M_{K_e \cup A}^1 \# M_{K_e \cup B}^2)$$

and

$$V_J |_{\Omega(f_h)} \simeq \bigoplus_{(A,B) \in \mathcal{M}_{K_0}} [(M_{K_e \cup A \cup \{t\}}^1 \# M_{K_e \cup B}^2) \oplus (M_{K_e \cup A}^1 \# M_{K_e \cup B \cup \{t\}}^2)].$$

We need to show that

$$\text{Hom}_{k\Omega(f_h)}((M_{K_e \cup A \cup \{t\}}^1 \# M_{K_e \cup B}^2) \oplus (M_{K_e \cup A}^1 \# M_{K_e \cup B \cup \{t\}}^2), E) = 0$$

for each $(A, B) \in \mathcal{M}_{K_0}$. In view of the discussion regarding D_t in the $f = f_e$ case, we prove that

$$\text{Hom}_{k\Omega(f_h)}((M_{K_e \cup A \cup \{t\}}^1 \# M_{K_e \cup B}^2) \oplus (M_{K_e \cup A}^1 \# M_{K_e \cup B \cup \{t\}}^2), V_{2t-2} \otimes V_{2t-2} \otimes V_K)$$

is zero for each $(A, B) \in \mathcal{M}_{K_0}$. Now if $(A, B), (A', B') \in \mathcal{M}_{K_0}$, then

$$\begin{aligned} & \text{Hom}_{k\Omega(f_h)}((M_{K_e \cup A \cup \{t\}}^1 \# M_{K_e \cup B}^2) \oplus (M_{K_e \cup A}^1 \# M_{K_e \cup B \cup \{t\}}^2), \\ & \quad V_{2t-2} \otimes V_{2t-2} \otimes (M_{K_e \cup A'}^1 \# M_{K_e \cup B'}^2)) \\ & \simeq \text{Hom}_{k\Omega(f_h)}((M_{K_e \cup A \cup \{t\}}^1 \# M_{K_e \cup B}^2) \otimes V_{2t-2} \oplus (M_{K_e \cup A}^1 \# M_{K_e \cup B \cup \{t\}}^2) \otimes V_{2t-2}, \\ & \quad V_{2t-2} \otimes (M_{K_e \cup A'}^1 \# M_{K_e \cup B'}^2)) \\ & \simeq \text{Hom}_{k\Omega(f_h)}((M_{K_e \cup A \cup \{t\}}^1 \# M_{K_e \cup B}^2) \otimes (M_{t-1}^1 \# M_{t-1}^2) \oplus (M_{K_e \cup A}^1 \# M_{K_e \cup B \cup \{t\}}^2) \\ & \quad \otimes (M_{t-1}^1 \# M_{t-1}^2), (M_{t-1}^1 \# M_{t-1}^2) \otimes (M_{K_e \cup A'}^1 \# M_{K_e \cup B'}^2)) \\ & \simeq \text{Hom}_{k\Omega(f_h)}((M_{K_e \cup A \cup \{t-1,t\}}^1 \# M_{K_e \cup B \cup \{t-1\}}^2) \oplus (M_{K_e \cup A \cup \{t-1\}}^1 \# M_{K_e \cup B \cup \{t-1,t\}}^2), \\ & \quad M_{K_e \cup A' \cup \{t-1\}}^1 \# M_{K_e \cup B' \cup \{t-1\}}^2) = 0 \end{aligned}$$

by Lemma 10. Here we have used: (i) the duality between the functors ‘Hom’ and ‘Tensor’, namely for any group X and kX -modules U_i , we have $\text{Hom}_{kX}(U_1, U_2 \otimes U_3) \simeq \text{Hom}_{kX}(U_1 \otimes U_2^*, U_3)$, where U^* is the dual of U ; and (ii) the fact that each simple module for $G = \text{Sp}(4, q)$ is self-dual. So Lemma 8 holds for $f = f_h$ also. ■

3. Proof of Theorem 1

(a) Let $n = 1$. Then $\dim_k(\bar{C}) = 9$ (see, for example, p. 308 of [13]). By Lemma 4, p. 488 of [14] and Frobenius reciprocity (p. 689 of [11]) it follows that V_1, k and V_0 (in descending order) are kG -composition factors of \bar{C} . Semisimplicity of each composition factor of \bar{C} and its multiplicity freeness as a $k\Omega(f)$ -module are proved in Corollaries 4 and 6. For

the remaining part of the proof, we treat the cases separately. Consider the case $f = f_e$. Since $V_0|_{\Omega(f)} \simeq W_0 \otimes W_1$ (Lemma 3) is the Steinberg module (and hence projective) for $\Omega(f)$, we need to prove that the kG -module E which is the unique (up to isomorphism) nonsplit extension of k by V_1 remains nonsplit as a $k\Omega(f)$ -module. Further, since E is isomorphic to a submodule of the kG -module $V_0 \otimes V_0$ (see Lemma 2(a), p. 161 of [17]), $V_1|_{\Omega(f)} \simeq W_0 \oplus W_1$ (Lemma 3) and $W_0 \otimes W_1$ is the Steinberg module for $k\Omega(f)$, by Theorem 1, p. 220 of [1], we have

$$\text{Hom}_{k\Omega(f)}(V_1, V_0 \otimes V_0) = 0.$$

Now we consider the case $f = f_h$. Then M^1, M^2 and $V|_{\Omega(f)} \simeq M^1 \# M^2$ (see Lemma 5) are the Steinberg modules for H_1, H_2 and $\Omega(f)$, respectively. Hence we need to prove that E as a $k\Omega(f)$ -module splits. But this is clear from Lemmas 5 and 10.

(b) Let $n \geq 2$. By Theorem 2, $\{V_I\}_{I \in \mathcal{N}}$ are the kG -composition factors of \bar{C} and they appear with multiplicity one. So, (i) follows from Corollaries 4 and 6.

We now prove (ii). Let V_K and $V_J, K, J \in \mathcal{N}$, be in the i -th and j -th kG -socle layers of $\bar{C}, 1 \leq i < j \leq 2n$. Assume that there is a kG -module E which is a nonsplit extension of V_K by V_J . Then, by Lemma 9(a), $j = i + 1$. Further, E is a nonsplit $k\Omega(f)$ -extension of V_K by V_J if and only if $J = K \cup \{2t - 1\}$ for some $t \in N_1$ (Lemma 7).

Further, if this holds, $\text{soc}_{k\Omega(f)}(E) = V_K$ (Lemma 8). That is, either all or none, of the simple $k\Omega(f)$ -summands of V_J descend. This observation together with the kG -socle structure of \bar{C} yields (ii).

Now, let $m \in \{0, 1, \dots, n - 2, n\}$ and $I \in \mathcal{A}_m$. In what follows, all modules considered are over $k\Omega(f)$. Since \bar{C} is multiplicity free as a $k\Omega(f)$ -module, it has a unique submodule $X_{m,I}$ with head V_K , where K is the unique element of \mathcal{N} with $K_e = I$ and $K_o = I^{\text{ac}}$. First $X_{0,I} = V_{\{2t: t \in I\}}$ (see (1_e) and (1_h)) and is semisimple. Let $m > 0$ and write $X_{m,I}$ as L for brevity. Since $\text{hd}(L)$ is contained in the m -th socle layer of \bar{C} , L is a submodule of $\text{soc}^m(\bar{C})$. For each summand of $\text{rad}^1(L)/\text{rad}^2(L)$ of the form V_J , a nonsplit extension of V_J by $\text{hd}(L)$ appears as a section of L . Further, $\text{hd}(L) = V_K$. So, $J = K \setminus \{2t - 1\}$ for some $t \in I^{\text{ac}}$ (Lemma 8). This proves that $\text{rad}^1(L)$ is contained in the unique submodule L' of $\text{soc}^{m-1}(\bar{C})$ whose head is isomorphic to

$$\bigoplus_{J \in \mathcal{N}, J_e = I, J_o \subseteq I^{\text{ac}} \text{ and } |J_o| = m - 1} V_J. \tag{3.1}$$

The multiplicity freeness of \bar{C} (as a $k\Omega(f)$ -module) implies that $L \cap L' = \text{rad}^1(L)$. We now show that $L' = \text{rad}^1(L)$. Since $(L + L')/L' \simeq L/(L \cap L') = L/\text{rad}^1(L) \simeq V_K$, by uniqueness of $L, L + L' = L$. Thus $L' \subseteq L$ and $L' = L \cap L' = \text{rad}^1(L)$. Thus, $\text{rad}^1(L)/\text{rad}^2(L)$ is the module in (3.1). Now successive application of this argument to $\text{rad}^i(L)/\text{rad}^{i-1}(L)$ yields the statement about the radical structure of $X_{m,I}$ in (iii). Further, (iv) is also clear.

Since $\text{soc}(X_{m,I}) = V_{\{2t: t \in I\}}$ is simple, $X_{m,I}$ is indecomposable. For $I \neq J \in \mathcal{A}_m, X_{m,I} \cap X_{m,J}$ is trivial because their socles are distinct and simple. (In fact, $X_{m,I}$ and $X_{m,J}$ have no composition factors in common.) Let

$$X_m = \bigoplus_{I \in \mathcal{A}_m} X_{m,I}.$$

For distinct $m, m' \in \{0, 1, \dots, n - 2, n\}, X_m$ and $X_{m'}$ have no composition factors in common and every composition factor of \bar{C} occurs in some X_m . Thus sum of X_m 's equals \bar{C} , completing the proof of all parts of (iii) and of Theorem 1. ■

Remark 11.

- (i) If $n = 1$, as a $k\Omega^\tau(f)$ -module, V_0 and V_1 are semisimple and \bar{C} is multiplicity free. Further, if f is of Witt index 1, then \bar{C} is isomorphic to the direct sum of V_0 and a $k\Omega^\tau(f)$ -module U with $\text{hd}(U) \simeq k$ and $\text{rad}(U) \simeq V_1$. If f is of Witt index 2, then \bar{C} is semisimple.
- (ii) If $n \geq 2$, as a $k\Omega^\tau(f)$ -module, then V_I is semisimple for each $I \in \mathcal{N}$ and \bar{C} is multiplicity free and is isomorphic to

$$\bigoplus_{m \in \{0, 1, \dots, n-2, n\}} \left(\bigoplus_{I \in \mathcal{A}_m} U_{m,I} \right),$$

where $U_{m,I}$ is a unique indecomposable module of radical length $m + 1$. Its j -th radical layer $\text{rad}^j(U_{m,I})/\text{rad}^{j+1}(U_{m,I})$ ($0 \leq j \leq m$) is isomorphic to

$$\bigoplus_{J \in \mathcal{N}, J_e=I, J_o \subseteq I^{ac} \text{ and } |J_o|=j} V_J.$$

Moreover, for each $m \in \{0, 1, \dots, n - 2, n\}$,

$$\text{Soc}^j(U_{m,I}) = \text{rad}^{m+1-j}(U_{m,I}).$$

Thus (i) and (ii) follow from an argument similar to the proof of Theorem 1, using Theorem 2 and Lemmas 7 and 8.

Examples. We illustrate the descent of the composition factors V_J in Theorem 2 when considered as a $k\Omega(f)$ -module for the cases $n = 1, 2, 3, 4$. For $J = \{i_1, i_2, \dots\} \subseteq N$, we write V_J as \underline{J} or $\underline{i_1, i_2, \dots}$.

If $n = 1$, then

$$\bar{C} = \underline{0} \oplus (\underline{1}/k).$$

If $n = 2$, then

$$\bar{C} = X_0 \oplus X_2 \equiv [\underline{0} \oplus \underline{2} \oplus \underline{0}, 2] \oplus [\underline{1}, 3/(\underline{1} \oplus \underline{3})/k].$$

If $n = 3$, then

$$\begin{aligned} \bar{C} = X_0 \oplus X_1 \oplus X_3 \equiv & [\underline{0}, \underline{2}, \underline{4} \oplus \underline{0}, \underline{2} \oplus \underline{0}, \underline{4} \oplus \underline{2}, \underline{4}] \oplus [(\underline{0}, \underline{3}/\underline{0}) \\ & \oplus (\underline{2}, \underline{5}/\underline{2}) \oplus (\underline{1}, \underline{4}/\underline{4})] \oplus [\underline{1}, \underline{3}, \underline{5}/(\underline{1}, \underline{3} \oplus \underline{1}, \underline{5} \oplus \underline{3}, \underline{5})/(\underline{1} \oplus \underline{3} \oplus \underline{5})/k]. \end{aligned}$$

If $n = 4$, then

$$\bar{C} = X_0 \oplus X_1 \oplus X_2 \oplus X_4,$$

where

$$X_0 = \bigoplus_{J \subseteq N_1, |J|=3 \text{ or } 4} \underline{2J} \oplus \underline{(0, 4 \oplus 2, 6)},$$

$$X_1 = \bigoplus_{i=0}^3 \underline{(2i, 2i+2, 2i+5)/(2i, 2i+2)},$$

$$X_2 = \underline{(0, 3, 5)/(0, 3 \oplus 0, 5)/0} \oplus \underline{(2, 5, 7)/(2, 5 \oplus 2, 7)/2}$$

$$\oplus \underline{(1, 4, 7)/(1, 4 \oplus 4, 7)/4} \oplus \underline{(6, 1, 3)/(6, 1 \oplus 6, 3)/6},$$

$$X_4 = A_4/A_3/A_2/A_1/A_0,$$

where A_i is the direct sum of \underline{J} as J runs over the subsets of $\{1, 3, 5, 7\}$ of size i .

Acknowledgement

We thank Professor Peter Sin for kindly pointing out an error in Theorem 1 in an earlier version of this paper. The second-named author also thanks Harish-Chandra Research Institute, Allahabad, India for kindly permitting him the use of its library and computing facilities during the period of this work.

References

- [1] Alperin J L, Projective modules for $SL(2, 2^n)$, *J. Pure Appl. Algebra* **15** (1979) 219–234
- [2] Dembowski P, Finite geometries, Classics in Mathematics (Berlin: Springer-Verlag) (1997) (reprint of 1968 edition)
- [3] Dye R H, Interrelations of symplectic and orthogonal groups in characteristic two, *J. Algebra* **59** (1979) 202–221
- [4] Dye R H, On the Arf invariant, *J. Algebra* **53** (1978) 36–39
- [5] Flesner D E, The geometry of subgroups of $PSp_4(2^n)$, *Illinois J. Math.* **19** (1975) 48–70
- [6] Flesner D E, Maximal subgroups of $PSp_4(2^n)$, containing central elations or noncentered skew elations, *Illinois J. Math.* **19** (1975) 247–268
- [7] Hirschfeld J W P, Finite projective spaces of three dimensions (Oxford: Oxford University Press) (1985)
- [8] Hirschfeld J W P, Projective geometries over finite fields (Oxford: Oxford University Press) (1998)
- [9] Huppert B and Blackburn N, Finite groups (Berlin: Springer-Verlag) (1982) vol. 2
- [10] Jones A, Integral representations of direct product of groups, *Canad. J. Math.* **15** (1963) 625–630
- [11] Lang S, Algebra, Third edition (New York: Addison-Wessley) (1999)
- [12] Payne S E and Thas J A, Finite generalized quadrangles (Boston: Advance Publishing Program, Pitman) (1984)
- [13] Sastry N S N, Codes and generalized polygons, in: Proc. of Seminar on combinatorics and applications in honour of Prof. S S Shrikhande (Calcutta: Indian Statistical Institute) (1982) pp. 303–310
- [14] Sastry N S N and Sin P, The code of regular generalized quadrangle of even order, *Proc. Symp. Pure Math.* **63** (1998) 485–496
- [15] Sastry N S N and Sin P, The binary code associated with nondegenerate quadrics of a symplectic space of even order, *J. Combin. Theory (A)* **94** (2001), 1–14

- [16] Sastry N S N and Sin P, On the doubly transitive permutation representations of $Sp(2n, \mathbb{F}_2)$, *J. Algebra* **257** (2002) 509–527
- [17] Sin P, Extension of simple modules for $Sp_4(2^n)$ and $Sz(2^n)$, *Bull. London Math. Soc.* **24** (1992) 159–164
- [18] Steinberg R, Representations of Algebraic groups, *Nagoya J. Math.* **22** (1963) 33–56
- [19] Steinberg R, Lectures on Chevalley Groups, Mimeographed Notes (New Haven, Conn.: Yale Univ. Math. Dept) (1968)
- [20] Taylor D E, The geometry of the classical groups, sigma series in pure mathematics (Berlin: Helderman-Verlag) (1992) vol. 9