

Euler characteristic and quadrilaterals of normal surfaces

TEJAS KALELKAR

Stat-Math Unit, Indian Statistical Institute, Bangalore 560 009, India
E-mail: tejas@isibang.ac.in

MS received 4 October 2006; revised 12 September 2007

Abstract. Let M be a compact 3-manifold with a triangulation τ . We give an inequality relating the Euler characteristic of a surface F normally embedded in M with the number of normal quadrilaterals in F . This gives a relation between a topological invariant of the surface and a quantity derived from its combinatorial description. Secondly, we obtain an inequality relating the number of normal triangles and normal quadrilaterals of F , that depends on the maximum number of tetrahedrons that share a vertex in τ .

Keywords. Euler characteristic; normal surfaces.

1. Introduction

Let M be a compact 3-manifold with a triangulation τ . The triangulation τ may be a pseudo-triangulation. A pseudo triangulation is a simplicial-complex where each 3-cell is a tetrahedron whose interior is embedded, but the boundary may not be injective. A surface F in M is said to be normal if it is properly embedded in M , it intersects each simplex transversally and it intersects each tetrahedron in a disk that can be isotoped, by an isotopy that is invariant on each simplex, to one of the 7 disk types shown in figure 1.

Normal surfaces were first introduced by Kneser [7] in 1929 and later developed by Schubert [11] and Haken [1,2] for use in an algorithm for recognising the unknot. Haken began the construction of an algorithm for solving the homeomorphism problem for a large class of 3-manifolds, the details of which were completed by Jaco and Oertel [4] and Hemion [3] for manifolds containing an incompressible 2-sided surface. Normal surfaces have been used for a variety of tasks by Rubinstein [9,10], Jaco and Tollefson [5] and Jaco, Letscher and Rubinstein [6]. For an extensive study of normal surface theory we refer to [8].

An embedded normal surface is determined uniquely by the number of triangles and quadrilaterals of each type within each tetrahedron. These are called the normal coordinates of the normal surface. There exists a system of homogenous linear equations called the




Figure 1. The 4 normal triangle and 3 normal quadrilateral types.

matching equations which ensure that normal disks in adjacent tetrahedrons glue together to form a surface. As quadrilaterals of different types within the same tetrahedron intersect, the normal coordinates of a normal surface have at most one non-zero coordinate corresponding to the different quadrilateral types, in each tetrahedron. A solution to the matching equations is called admissible if at most one of the coordinates corresponding to the different quadrilateral types is nonzero, in each tetrahedron. The non-negative admissible solutions of the matching equations are in bijection with the set of embedded (not necessarily connected) normal surfaces. Every such solution is a linear combination of a finite set of fundamental solutions with non-negative integer coefficients. The additive structure of the solution space is geometrically realised as the Haken sum of the corresponding surfaces. It is this combinatorial description of normal surfaces that is utilised in solving algorithmic problems.

The goal of this paper is to give a relation between the Euler characteristic of a normal surface, a topological invariant, and the number of normal quadrilaterals in its embedding, obtained from its combinatorial description. Secondly, we get a relation between the number of normal triangles and normal quadrilaterals.

Theorem 1.1. *Let F be a normal surface in a triangulated 3-manifold M . Let Q be the number of normal quadrilaterals in F . Then,*

$$\chi(F) \geq 2 - 7Q.$$

In particular, if F is an oriented, closed and connected normal surface of genus g ,

$$g \leq \frac{7}{2}Q.$$

DEFINITION 1.2

Let F be a normal surface in M . Let t be a normal triangle of F that lies in a tetrahedron Δ . The triangle t is said to link a vertex v of Δ if t separates $\partial\Delta$ into two disks such that the disk containing v has no other vertices of Δ . Similarly, a normal arc α in a face \mathcal{F} is said to link a vertex v of \mathcal{F} if the segment containing v in $\partial\mathcal{F} - \alpha$ has no other vertices of \mathcal{F} .

DEFINITION 1.3

Let $S(v)$ be the boundary of a small ball neighbourhood of v . The sphere $S(v)$ is a normal surface composed of normal triangles linking v , one from each normal isotopy class. This is defined to be the vertex linking sphere linking vertex v .

Remark 1.4. Any closed connected normal surface S in M composed of normal triangles is normally isotopic to a vertex-linking sphere. This is due to the following reasons.

Firstly, all normal triangles in S link the same vertex. If this were not true, there would be normal triangles t_1 and t_2 in S , linking distinct vertices of τ , that intersect in common edges. This is not possible as normal arcs linking different vertices of a face are not normally isotopic. So, let v be the common vertex linked by triangles of S . Then S is a cover of $S(v)$, as $S(v)$ is composed of one triangle from each of the normal isotopy classes of triangles linking vertex v . As $S(v)$ is a sphere and S is a closed connected cover of $S(v)$, the covering projection map is a homeomorphism. Therefore, there is only one triangle of S in each normal isotopy class of triangles linking v and thus, S is normally isotopic to $S(v)$.

This remark is the motivation for Theorem 1.1. The remark is not true for closed connected normal surfaces embedded in 3-complexes. For example, take a triangulation τ of a closed, oriented surface F and let $C(F)$ be the cone of F . Take the triangulation of $C(F)$ obtained by taking the corresponding cone of each triangle of τ . Push F into the interior of this 3-complex so that F is embedded as a closed normal surface composed entirely of normal triangles.

We expect that in Riemannian 3-manifolds, triangles correspond to positive curvature pieces. This would imply that when quadrilaterals have a curvature that is bounded below, the smaller the Euler characteristic, the greater would be the number of quadrilaterals.

Remark 1.5. An inequality relating the Euler characteristic with the total number of normal disks is easy to obtain. This is because the number of normal discs gives a bound on the number of disjoint closed curves on the surface, up to isotopy. This, in turn, gives a lower bound on the Euler characteristic.

Our second theorem, Theorem 1.6, gives a relation between the number of triangles and the number of quadrilaterals of an embedded normal surface. Let A be the union of triangles of an embedded normal surface F , v a vertex of the triangulation τ , N_v the number of tetrahedrons containing v and σ a *strongly connected* (defined in Definition 2.3) component of the triangles of A linking vertex v . By Lemma 2.4, σ has at most N_v triangles. Any two such components are ‘connected’ by quadrilaterals. So we get an upper bound on the number of triangles in terms of the number of quadrilaterals and the maximum number of tetrahedrons linking a vertex of τ .

Theorem 1.6. *Let F be a normal surface in M , no component of which is a vertex-linking sphere. Let T and Q be the number of normal triangles and normal quadrilaterals of F respectively. Let N be the maximum number of tetrahedrons linking a vertex of τ . Then,*

$$T \leq 4NQ.$$

2. An inequality relating the Euler characteristic and the number of quadrilaterals of normal surfaces

In this section we give a proof of Theorem 1.1. Let F be a normal surface in M . Let Q be the number of normal quadrilaterals in F . Let A (respectively B) be the union of normal triangles (respectively normal quadrilaterals) of F . A and B are 2-complexes which are subsets of the surface F . Define the boundary of a 2-complex σ to be the union of edges of σ that do not intersect $\text{int}(\sigma)$.

An outline of the proof is as follows. We suitably modify the 2-complexes A and B , at singular points, to get surfaces A' and B' that intersect in a disjoint union of circles with $A' \cup B' = F$. We then show that the connected components of A' are planar surfaces. We, therefore, get a lower bound on the Euler characteristic of A' in terms of the number of components $|\partial A'|$ of $\partial A'$. The boundaries $\partial A'$ and $\partial B'$ are equal. We introduce weights and get an inequality relating $|\partial B'|$ with the weight $w(\partial B)$ of ∂B . Now as $w(\partial B) \leq w(B) = 4Q$ we obtain an upper bound on $|\partial A'|$ in terms of the number of quadrilaterals. Thus, we get a lower bound on the Euler characteristic of A' in terms of the number of quadrilaterals.

For a bound on the Euler characteristic of B' , we take an increasing union of quadrilaterals and use the fact that each new quadrilateral intersects the union of the preceding stage in a circle or at most 4 disjoint segments. This gives us a lower bound in terms of the number of quadrilaterals.

As the Euler characteristic of F is the sum of Euler characteristics of A' and B' , we get a lower bound for the Euler characteristic of F in terms of Q .

DEFINITION 2.1

Let D_n be the collection of normal discs of F . Let D'_n be the collection of triangles obtained by taking the barycentric subdivision of D_n . Then, the regular neighbourhood of a point w in F is defined as $N(w) = \text{int}(\cup\{D'_n : w \in D'_n\})$.

DEFINITION 2.2

A point w in A is said to be a singular point if $N(w)$ is not homeomorphic to an open set in the upper half-plane $\{(x, y) \in \mathbb{R}^2 \mid y \geq 0\}$. Let ω be the set of singular points of A and $N(\omega) = \cup\{N(w) : w \in \omega\}$.

Let $A' = A - N(\omega)$, $B' = B \cup \overline{N(\omega)}$. Therefore, A' and B' are compact surfaces with $F = A' \cup B'$ and $A' \cap B'$ a disjoint union of circles. The surface A' is the closure of the interior of A , while B' is the closure of a neighbourhood of B in F .

DEFINITION 2.3

A subcomplex σ of A is said to be strongly connected if for any two triangles t and t' in σ , there exists a sequence of triangles $\{t_i\}_{i=1}^n$ with $t = t_1$ and $t' = t_n$, such that for all i , t_i and t_{i+1} intersect in at least one edge. A subcomplex consisting of a single triangle is taken to be strongly connected. It is easy to see that A can be decomposed as a union of maximal strongly connected components.

Lemma 2.4. The interior of a strongly connected component σ of A is a planar surface. Let N_v be the number of tetrahedrons that contain the vertex v . Let N be the maximum N_v over all vertices v in the triangulation τ . Then there are at most N triangles in σ .

Proof. All triangles in the strongly connected component σ of A link the same vertex. Otherwise, there would be a common normal arc shared by a normal triangle of F linking a vertex v and a normal triangle of F linking a distinct vertex w . This is a contradiction as normal arcs in a face, linking different vertices are not normally isotopic. So we can assume all triangles of σ link the same vertex v .

To each normal triangle of σ we can associate a height from the vertex v . Given a height h from the vertex v , each tetrahedron has at most one triangle of σ at height h . So, in all, there are at most N_v triangles at height h . Triangles linking v at different heights from v do not intersect. So, as σ is connected, all triangles of σ can be assumed to be at the same height. Thus, there are at most N_v triangles in σ .

Furthermore, the union of all possible normal triangles linking v at the height h (from v) gives a closed connected normal surface. Hence by Remark 1.4, it is a vertex linking sphere $S(v)$. As σ is composed of triangles at height h from v , $\sigma \subset S(v)$. So the interior of σ is a planar surface. □

Lemma 2.5. Let $|A'|$ denote the number of components of A' . Then,

$$\chi(A') \geq 2|A'| - 4Q.$$

Proof.

Claim 1. $\chi(A') = 2|A'| - |\partial A'|$. Let $\{A_n\}_n$ be the strongly connected components of A . Then, $\text{int}(A') = \text{int}(A) = \cup_n \text{int}(A_n)$. By Lemma 2.4, the interior of A_n are planar surfaces. So, as A' is a surface (not just a 2-complex), A' is itself a disjoint union of connected planar surfaces.

A connected planar surface S with b boundary components has $\chi(S) = 2 - b$. As A' is a disjoint union of connected planar surfaces $\chi(A') = 2|A'| - |\partial A'|$ as claimed.

DEFINITION 2.6

Let σ be a 2-complex. Define $w(e)$, weight of an edge $e \in \sigma$, to be

$$w(e) = \begin{cases} 1, & \text{if } e \subset \partial\sigma \\ 2, & \text{otherwise.} \end{cases}$$

Define $w(\sigma)$, the weight of σ , to be the total weight of all edges of σ . For ψ a 1-complex, define $w(\psi)$ to be the total number of edges of ψ . From the definition, it is clear that $w(\sigma) \geq w(\partial\sigma)$.

Claim 2. $4Q \geq |\partial A'|$. The boundary of B is a union of simple closed curves $\{c_i\}$ that pairwise intersect in singular points. The set $\{c_i\}$ is in bijection with the connected components of $\partial B'$. Therefore, $w(\partial B) = \sum w(c_i) \geq |\partial B'|$.

As $4Q = w(B) \geq w(\partial B)$ and $\partial B' = \partial A'$, we get the inequality $4Q \geq |\partial A'|$ as claimed.

Using Claims 1 and 2, we get the desired relation

$$\chi(A') \geq 2|A'| - 4Q.$$

□

Lemma 2.7. For $B \neq \phi$,

$$\chi(B') \geq \begin{cases} |\omega| - 3Q, & \text{when } |\omega| > 0 \\ 4 - 3Q, & \text{when } |\omega| = 0 \end{cases}.$$

Proof. Let $B'_0 = \overline{N(\omega)}$. Let B'_n be a union of n quadrilaterals of B' such that $B'_{n+1} \supset B'_n$. For a quadrilateral q such that $B'_{n+1} = B'_n \cup q$, $B'_n \cap q$ is either a circle or a disjoint union of at most 4 arcs (or points). So, $\chi(B'_n \cap q) \leq 4$, and we get

$$\begin{aligned} \chi(B'_{n+1}) &= \chi(B'_n) + \chi(q) - \chi(B'_n \cap q) \\ &\geq (\chi(B'_n) + 1) - 4. \end{aligned}$$

As $\overline{N(\omega)}$ is a disjoint union of discs,

$$\chi(B'_0) = |\omega|.$$

Therefore,

$$\chi(B') = \chi(B'_Q) \geq \begin{cases} |\omega| - 3Q, & \text{when } |\omega| > 0 \\ 1 - 3(Q - 1), & \text{when } |\omega| = 0 \end{cases}.$$

□

Proof of Theorem 1.1. When $B = \phi$, the surface F is composed of normal triangles and so, is a union of vertex linking spheres. Therefore as $A \neq \phi$, $\chi(F) = 2|A| \geq 2$.

Assume $B \neq \phi$, as $A' \cap B'$ is a disjoint union of circles. Therefore using Lemmas 2.5 and 2.7 we get

$$\begin{aligned} \chi(F) &= \chi(A' \cup B') \\ &= \chi(A') + \chi(B') \\ &\geq \begin{cases} (2|A'| - 4Q) + (|\omega| - 3Q), & \text{when } |\omega| > 0 \\ (2|A'| - 4Q) + 4 - 3Q, & \text{when } |\omega| = 0 \end{cases} \end{aligned}$$

When $|A'| = 0$ the surface is composed entirely of quadrilaterals, so that $B \neq \phi$ and $|\omega| = 0$. So, we get the desired relation

$$\chi(F) \geq 2 - 7Q.$$

□

3. An inequality relating the number of triangles and quadrilaterals of normal surfaces

In this section we prove an inequality relating the number of triangles and quadrilaterals of a normal surface.

Proof of Theorem 1.6. Let $F(v)$ be the union of normal triangles of F linking the vertex v . Let $\{F(v)_n\}_n$ be the strongly connected components of $F(v)$.

Consider a graph Γ whose vertices are the normal quadrilaterals of F and the non-empty $F(v)_n$. Call the former vertices Q -vertices and the latter as S -vertices. Let there be an edge of Γ joining an S -vertex and a Q -vertex for every edge shared by the corresponding $F(v)_n$ and the corresponding quadrilateral. Let there also be edges of Γ between Q -vertices corresponding to every edge shared by the corresponding quadrilaterals. For a quadrilateral in the pseudo triangulation we could have simple loops corresponding to quadrilaterals that are not injective on its edges.

As each quadrilateral has 4 sides, the degree of each Q -vertex is 4 and as there are no vertex-linking spheres, there are no isolated S -vertices. As each edge of Γ has at least one vertex incident on a Q -vertex (no edges between S -vertices) therefore, $4Q = \text{total degree of } Q \text{ vertices} \geq \text{total degree of } S\text{-vertices} \geq S$, where Q and S are the number of Q -vertices and S -vertices respectively.

By Lemma 2.4 each $F(v)_n$ has at most N triangles, where N is the maximum number of tetrahedrons linking a vertex in the triangulation of M . Therefore, $NS \geq T$, where T is the number of triangles in F . Therefore, we get the required relation,

$$T \leq 4NQ.$$

Remark 3.1. This proof also holds for a normal surface F embedded in a 3-complex, if we assumed that no component of F was composed solely of normal triangles.

□

Acknowledgements

The author would like to thank Siddhartha Gadgil for useful discussions and advice. The CSIR-SPM Fellowship is acknowledged for financial support.

References

- [1] Haken Wolfgang, Theorie der Normalflächen, *Acta Math.* **105** (1961) 245–375
- [2] Haken Wolfgang, Über das Homöomorphieproblem der 3-Mannigfaltigkeiten. I, *Math. Z.* **80** (1962) 89–120
- [3] Hemion Geoffrey, The classification of knots and 3-dimensional spaces (Oxford Science Publications, Oxford University Press) (1992)
- [4] Jaco William and Oertel Ulrich, An algorithm to decide if a 3-manifold is a Haken manifold, *Topology* **23(2)** (1984) 195–209
- [5] Jaco William and Tollefson Jeffrey L, Algorithms for the complete decomposition of a closed 3-manifold, *Illinois J. Math.* **39(3)** (1995) 358–406
- [6] Jaco William, Letscher David and Rubinstein J Hyam, Algorithms for essential surfaces in 3-manifolds, *Topology and Geometry: Commemorating SISTAG*, Contemporary Mathematics, no. 314 (RI, Providence: Amer. Math. Soc.) (2002) 107–124
- [7] Kneser Hellmuth, Geschlossene Flächen in dreidimensionalen Mannigfaltigkeiten, *Jahresbericht der Deut. Math. Verein.* **38** (1929) 248–260
- [8] Matveev Sergei, Algorithmic Topology and Classification of 3-Manifolds (Berlin, Heidelberg: Springer-Verlag) (2003) vol. 9
- [9] Rubinstein J Hyam, An algorithm to recognise the 3-sphere, Proceedings of the International Congress of Mathematicians (Zurich) (1994) vol. 1 (Birkhauser) (1995) pp. 601–611
- [10] Rubinstein J Hyam, Polyhedral minimal surfaces, Heegaard splittings and decision problems for 3-dimensional manifolds, *Geometric Topology* (GA: Athens) (1993) AMS/IP Stud. Adv. Math., vol. 2, Amer. Math. Soc. (1997) pp. 1–20
- [11] Schubert H, Bestimmung der Primfactor Zerlegung von Verkettungen, *Math. Z.* **76** (1961) 116–148