

Multiple decomposability of probabilities on contractible locally compact groups

WILFRIED HAZOD and KATRIN KOSFELD

Faculty of Mathematics, Technische Universität Dortmund, Vogelpothsweg 87,
D-44227 Dortmund, Germany
E-mail: Wilfried.Hazod@math.uni-dortmund.de;
Katrin.Kosfeld@math.uni-dortmund.de

MS received 30 March 2011; revised 15 November 2011

Abstract. Operator decomposable probabilities on vector spaces – generalizing (semi-)stable and self-decomposable laws – are well known. Here we are concerned with *multiple* decomposability on locally compact groups. In fact, as it turned out that contraction properties play an essential role, throughout we concentrate on contractible locally compact groups.

Keywords. Contractible locally compact group; group norm; multiple decomposable probability; logarithmic moment.

1. Introduction

On vector spaces, operator-decomposability of probabilities – a common setup for operator semi-stable, stable and self-decomposable laws – were the subject of various investigations. In analogy to the Urbanik classes in the case of self-decomposability, nested classes of multiple operator decomposable laws and their properties were investigated. For the background the reader is referred to [12, 31, 32] for finite and infinite dimensional real vector spaces, and [16] and the references therein for multiple decomposability. In case of locally compact groups we mention [27] and [23] (see also [6]), and for multiple decomposability on p -adic groups and vector spaces, [18] and the literature mentioned there. (For some aspects of multiple self-decomposability, i.e., decomposability with respect to a continuous one-parameter group of automorphisms, on groups see [4].)

Let \mathbb{G} be a locally compact group and $\tau \in \text{Aut}(\mathbb{G})$. A probability $\mu \in \mathcal{M}^1(\mathbb{G})$ is called τ -decomposable – in symbols: $\mu \in L_0(\tau)$ – if there exists a cofactor $\nu \in \mathcal{M}^1(\mathbb{G})$ such that $\mu = \nu \star \tau(\mu)$. Put $L_{-1}(\tau) := \mathcal{M}^1(\mathbb{G})$. μ is called n -times τ -decomposable, $n \geq 0$, – in symbols: $\mu \in L_n(\tau)$ – if μ is τ -decomposable with a cofactor $\nu \in L_{n-1}(\tau)$. (For details, see Definition 5.1 below.)

A continuous convolution semigroup $(\mu_t)_{t \geq 0}$, of probabilities is τ -decomposable if for all $t > 0$, μ_t is τ -decomposable with cofactor ν_t , where the cofactors $(\nu_t)_{t \geq 0}$ form a continuous convolution semigroup. Analogously, $(\mu_t)_{t \geq 0}$ is n -times τ -decomposable if it is τ -decomposable and the cofactor semigroup is $(n - 1)$ -times τ -decomposable.

This motivates the following definition: Let \mathcal{A} denote the generating functional of $(\mu_t)_{t \geq 0}$, $\langle \mathcal{A}, f \rangle = \frac{d^+}{dt} |_{t=0} \langle \mu_t, f \rangle$, evaluated for $f \in \mathcal{D}(\mathbb{G})$, the Bruhat–Schwartz test

function space. Let $\mathcal{GF}(\mathbb{G})$ denote the cone of generating functionals. \mathcal{A} is τ -decomposable – in symbols: $\mathcal{A} \in \Lambda_0(\tau)$ – if for some generating functional $\mathcal{A}^{(1)}$ we have $\mathcal{A} = \mathcal{A}^{(1)} + \tau(\mathcal{A})$. Put $\Lambda_{-1}(\tau) := \mathcal{GF}(\mathbb{G})$. Analogously, \mathcal{A} is n -times τ -decomposable ($n \in \mathbb{Z}_+$) – in symbols: $\mathcal{A} \in \Lambda_n(\tau)$ – if $\mathcal{A} \in \Lambda_0(\tau)$, $\mathcal{A} = \mathcal{A}^{(n)} + \tau(\mathcal{A})$, with $\mathcal{A}^{(n)} \in \Lambda_{n-1}(\tau)$. Hence there exist generating functionals $\mathcal{A}^{(i)}$ such that $\mathcal{A}^{(i)} = \mathcal{A}^{(i-1)} + \tau(\mathcal{A}^{(i)})$ for $1 \leq i \leq n+1$ with $\mathcal{A}^{(n+1)} := \mathcal{A}$. Thus $\mathcal{A} = \mathcal{A}^{(0)} + \sum_1^{n+1} \tau(\mathcal{A}^{(i)})$. (For more details, see Definition 6.5.)

Recall that for non-Abelian groups, generating functionals of continuous convolution semigroups play the role of logarithms of Fourier transforms, the second characteristic functions. For details the reader is referred to Ch IV, § 4.2 of [10], [3, 7, 29].

Throughout we have to suppose that \mathbb{G} is contractible and $\tau \in \text{Aut}(\mathbb{G})$ is contracting, a natural condition. In complete analogy to the vector space case, we prove that n -times τ -decomposable laws are representable as infinite convolution products, $\mu = \star_{j \geq 0} \tau^j(\nu)$ (for $n = 0$) resp. $\mu = \star_{j_n \geq 0} \left(\dots \left(\star_{j_0 \geq 0} \tau^{\sum_0^n j_i}(\nu) \right) \right)$ (for $n \geq 1$). (cf. Proposition 5.2). Under commutativity assumptions – and if a technical condition is satisfied (trivially satisfied e.g. in the case of homogeneous groups) – we obtain, re-arranging the iterated convolution products: μ is n -times τ -decomposable iff $\mu = \star_{k \geq 0} \tau^k \left(\nu^{\binom{n+k}{n}} \right)$, and this is the case iff ν possesses logarithmic moments of order $n+1$.

In case of continuously embeddable probabilities we obtain, again in analogy to the vector space situation – at least for connected resp. totally disconnected contractible groups – characterizations of multiple τ -decomposable generating functionals of continuous convolution semigroups which correspond to characterizations of the Lévy–Khinchin form in the case of real and p -adic vector spaces obtained in [16] and [18] respectively.

Most of the results are part of the second named author's thesis [13].

The paper is organized as follows: First, in § 2 we state the main results, Theorems 2.1 and 2.2. For their proofs we need several tools which are of independent interest. These tools, group norms and automorphism norms, resp. convergence of infinite convolution products, are collected in §§ 3 and 4. The proofs of the main results, and some slightly more general results follow in §§ 5 and 6 in a sequence of propositions. We discuss briefly the possibility of analogous investigations for certain classes of hypergroups in an [Appendix](#).

2. Main results

Throughout let \mathbb{G} denote a contractible locally compact group with contracting automorphism $\tau \in \text{Aut}(\mathbb{G})$. μ has a finite logarithmic moment of order n (in symbols, $\mu \in \mathcal{P}_{\log^n}$) if for a random variable X with distribution μ , $\mathbb{E}((\log_+(|X|))^n) = \int_{\mathbb{G}} (\log_+ |x|)^n d\mu(x) < \infty$, where $|\cdot|$ denotes a group norm on \mathbb{G} . (cf. Definition 3.1). As mentioned before, μ is n -times τ -decomposable iff μ has a representation as $(n+1)$ -times iterated convolution product. To be allowed to re-arrange convolution products (in case $n > 0$) we define $\mathcal{S} \subseteq \mathcal{M}^1(\mathbb{G})$ to be a commutative sub-semigroup with $\tau(\mathcal{S}) \subseteq \mathcal{S}$ and $\nu \in \mathcal{S}$.

Theorem 2.1. *Let $\mu, \nu \in \mathcal{S}$, $n \in \mathbb{Z}_+$.*

(a) $\mu \in \mathcal{M}^1(\mathbb{G})$ is n -times τ -decomposable iff

$$\mu = \star_{j_n \geq 0} \left(\star_{j_{n-1} \geq 0} \dots \left(\star_{j_0 \geq 0} \tau^{\sum_1^n j_i}(\nu) \right) \dots \right). \quad (2.1)$$

And if the right invariance group is trivial, $H_r(\mu) := \{x : \mu \star \varepsilon_x = \mu\} = \{e\}$, then

$$\mu = \star_{j=0}^{\infty} \tau^j \left(\nu^{\binom{n+j}{n}} \right). \quad (2.2)$$

- (b1) If $\nu \in \mathcal{P}_{\log_+^{n+1}}$, the convolution products (2.1), (2.2) and all rest products $\lambda_k := \star_{j=k+1}^{\infty} \tau^j \left(\nu^{\binom{n+j}{n}} \right)$, $k \in \mathbb{Z}_+$, converge and $\lambda_k \xrightarrow{k \rightarrow \infty} \varepsilon_e$.
- (b2) Conversely, assume that the products (2.1) or (2.2) are converging, and again $H_r(\mu) = \{e\}$, then $\nu \in \mathcal{P}_{\log_+^{n+1}}$.

If \mathbb{G} is a homogeneous group hence aperiodic, all invariance groups are trivial, and in the case $n \geq 0$, the condition $H_r(\mu) = \{e\}$ is trivially satisfied. And also for general locally compact contractible groups, for $n = 0$, the condition $H_r(\mu) = \{e\}$ is superfluous (cf. Proposition 4.9).

Let μ be embedded into a continuous convolution semigroup (μ_t) , and let \mathcal{A} denote the generating functional. Let (multiple) τ -decomposability of continuous convolution semigroups resp. of generating functionals be defined as before. In this case we obtain

Theorem 2.2.

(a) Let \mathbb{G} be a connected contractible group, hence a homogeneous group, and let $\mathcal{G} \cong \mathbb{R}^d$ denote the Lie algebra. Let $\tau \in \text{Aut}(\mathbb{G})$ and let $\overset{\circ}{\tau} \in \text{Aut}(\mathcal{G}) \subseteq \text{GL}(\mathbb{R}^d)$ denote the differential. Hence $\exp^{-1} \circ \tau = \overset{\circ}{\tau} \circ \exp^{-1}$. Let $\overset{\circ}{\mathcal{A}}$ denote the image of \mathcal{A} under \exp^{-1} . Then we have:

\mathcal{A} is n -times τ -decomposable (on \mathbb{G}) iff $\overset{\circ}{\mathcal{A}}$ is n -times τ -decomposable (on the vector space \mathcal{G}). Equivalently, if the continuous convolution semigroup $(\lambda_t) \subseteq \mathcal{M}^1(\mathcal{G})$ generated by $\overset{\circ}{\mathcal{A}}$ is n -times $\overset{\circ}{\tau}$ -decomposable.

This is the case iff the Lévy–Khinchin representation of $\overset{\circ}{\mathcal{A}}$ has the form given in Theorem 3.1 of [16].

(b) Let \mathbb{G} be a contractible totally disconnected locally compact group. Then there exist $p \in \mathbb{N}$, $p > 1$ (not necessarily a prime power), a homeomorphism $\psi : \mathbb{G} \rightarrow \mathcal{G} := \mathbb{Q}_p$, the additive group of p -adic numbers, and $\overset{\circ}{\tau} \in \text{Aut}(\mathbb{Q}_p)$, such that $\psi \circ \tau = \overset{\circ}{\tau} \circ \psi$. Again for a generating functional \mathcal{A} , let $\overset{\circ}{\mathcal{A}} := \psi(\mathcal{A})$ denote the image under ψ . Then we have:

\mathcal{A} is n -times τ -decomposable (on \mathbb{G}) iff $\overset{\circ}{\mathcal{A}}$ is n -times $\overset{\circ}{\tau}$ -decomposable (on \mathcal{G}). Equivalently, if the continuous convolution semigroup $(\lambda_t) \subseteq \mathcal{M}^1(\mathcal{G})$ generated by $\overset{\circ}{\mathcal{A}}$ is n -times $\overset{\circ}{\tau}$ -decomposable.

This is the case iff the Lévy–Khinchin representation of $\overset{\circ}{\mathcal{A}}$ on \mathbb{Q}_p has the form given in Theorems 4.3 and 5.2 of [18].

3. Group norms and automorphism norms on contractible groups

Let \mathbb{G} denote a locally compact group, $\text{Aut}(\mathbb{G})$ the group of topological automorphisms endowed with the topology of uniform convergence on compact subsets.

DEFINITION 3.1

A group norm on \mathbb{G} is a continuous sub-additive functional $|\cdot| : \mathbb{G} \rightarrow \mathbb{R}_+$ such that

- (i) $|x| = |x^{-1}|$ for all $x \in \mathbb{G}$,
- (ii) $|x| = 0$ iff $x = e$,
- (iii) $x_n \rightarrow e$ iff $|x_n| \rightarrow 0$.

Thus $d(x, y) := |x^{-1}y|$ defines a right-invariant metric generating the topology of \mathbb{G} .

Remark 3.2. In the case of contractible locally compact groups we shall show that group norms always exist (see Proposition 3.10). Note that for homogeneous groups (homogeneous) norms are usually defined by a weaker condition $|xy| \leq C(|x| + |y|)$ for some $C \geq 1$. However it is known that sub-additive homogeneous norms, i.e., norms with $C = 1$, exist (cf. [8]). Hence here we restrict to sub-additive functionals.

Note furthermore, that group norms are frequently called *pseudonorms* (generating the topology of \mathbb{G}), cf. [33].

Obviously we have: Let $|\cdot|_i$, $i = 1, 2$, and $|\cdot|$ be group norms. Then the following functionals are also group norms:

- (1) $|\cdot|_* := \max_{i=1,2} |\cdot|_i$,
- (2) $|\cdot|_0 := |\cdot|_1 + |\cdot|_2$,
- (3) $|\cdot|_\sigma := |\sigma(\cdot)|$ for $\sigma \in \text{Aut}(\mathbb{G})$ and
- (4) $\log(|\cdot| + 1)$.

In the sequel we make use of the following examples:

Example 3.3. Let \mathbb{G} be a *connected* contractible group, therefore a homogeneous group. In particular, \mathbb{G} is a simply connected nilpotent Lie group admitting a continuous one-parameter group of *dilations* $(\delta_t)_{t>0} \subseteq \text{Aut}(\mathbb{G})$, such that $\delta_t \delta_s = \delta_{t \cdot s}$ and $\delta_t(x) \rightarrow e$, $t \rightarrow 0$, for all $x \in \mathbb{G}$. Then there exists a group norm satisfying $|\delta_t(x)| = t \cdot |x|$ for $t > 0$, $x \in \mathbb{G}$. Thus $\mathbb{G} \setminus \{e\} \cong \mathbb{R}_+^\times \otimes \{x : |x| = 1\}$. Such a norm – a sub-additive homogeneous norm – will be called *adapted to the dilations*.

Any two homogeneous norms adapted to (δ_t) are equivalent. For details see §2.7 of [7] and the literature mentioned there, in particular §2.16 (references and comments to chapter II), R 2.7.IV.

Example 3.4. Let \mathbb{G} be *totally disconnected* and assume that there exist compact open subgroups U_n , $n \in \mathbb{Z}$ such that $U_{n+1} \triangleleft U_n$, $n \in \mathbb{Z}$, $\bigcup U_n = \mathbb{G}$, $\bigcap U_n = \{e\}$. ((U_n) is called filtration by compact open subgroups.) Fix $0 < \alpha < 1$. Define $|x| := |x|_\alpha := \alpha^k$ if $x \in U_k \setminus U_{k+1}$, and $|e| := 0$. Then $|\cdot|$ is a group norm, *adapted to* (U_n) . (In fact, let $k \leq \ell$, $x \in U_k$, $y \in U_\ell$. Then $xy \in U_k U_\ell \subseteq U_k$. Hence $|xy| \leq \alpha^k$. i.e., $|xy| \leq \max\{|x|, |y|\} \leq |x| + |y|$.) The other assertions are immediately verified.

Finite dimensional vector spaces are (commutative) homogeneous groups, with homothetical transformations $H_t : x \mapsto t \cdot x$ as dilations. And vector space norms are homogeneous norms. As on vector spaces, homogeneous norms on a homogeneous group (with respect to fixed dilations) are equivalent. In contrast, this is not the case for group norms (adapted to τ) as the following example shows:

Example 3.5. Let as in Examples 3.4 and 3.8 (below), $(U_n)_{n \in \mathbb{Z}}$ be a filtration of \mathbb{G} with compact neighbourhoods of e , i.e., $U_n \supseteq U_{n+1}$, $\bigcup U_n = \mathbb{G}$, $\bigcap U_n = \{e\}$. Let $|\cdot|$ be a

group norm, and assume for some $\alpha \in (0, 1)$ that $\alpha^{n+1} \leq |x| \leq \alpha^n$ for $x \in U_n \setminus U_{(n+1)}$. Consider the group norm $|x|_1 := \log(|x| + 1)$. For $x_n \in U_{-(n+1)} \setminus U_{-n}$, $n \in \mathbb{N}$, we have $|x_n|, |x_n|_1 \rightarrow \infty$, but $|x_n|_1/|x_n| \geq \alpha^{-n+1}/\log(1 + \alpha^{-n}) \rightarrow \infty$. Hence the norms are not equivalent.

As easily verified, such filtrations exist for the groups considered in Examples 3.3 and 3.4.

DEFINITION 3.6

An automorphism norm corresponding to a group norm $|\cdot|$ is a sub-multiplicative functional $\|\cdot\| : \text{Aut}(\mathbb{G}) \rightarrow \mathbb{R}_+$ satisfying $|\sigma(x)| \leq \|\sigma\| \cdot |x|$ ($x \in \mathbb{G}, \sigma \in \text{Aut}(\mathbb{G})$).

$\|\sigma\|_0 := \sup_{x \neq e} |\sigma(x)|/|x|$ is obviously an automorphism norm. It will be called canonical automorphism norm (with respect to $|\cdot|$). If not otherwise defined, $\|\cdot\|$ will always denote a canonical automorphism norm.

$\tau \in \text{Aut}(\mathbb{G})$ is called contracting if $\tau^n(x) \xrightarrow{n \rightarrow \infty} e$ for all x . As is well known, if τ is contracting then it is compactly contracting, i.e., for all compact $K \subseteq \mathbb{G}$ we have $\tau^n(K) \rightarrow \{e\}$ (cf. Lemma 1.4(iv) of [30], [35]).

We continue the previous Examples 3.3 and 3.4.

Example 3.7. Let \mathbb{G} be a contractible *connected* group as in Example 3.3 with contracting $\tau \in \text{Aut}(\mathbb{G})$. Then there exist a group of dilations (δ_t) with $\tau\delta_t = \delta_t\tau$ for all t . (cf. Remark 2.1.13(a) of [7]). Let $|\cdot|$ denote a (sub-additive homogeneous) norm adapted to the dilations. For short, *adapted to τ* . Then, as easily verified, for all automorphisms $\sigma \in \text{Cent}((\delta_t))$, in particular for $\sigma = \tau$, we have for the canonical norm:

- (1) $\|\sigma\| = \sup_{x \neq e} |\sigma(x)|/|x| = \max_{|x|=1} |\sigma(x)| = \max_{|x| \leq 1} |\sigma(x)|$.
- (2) τ is (compactly) contracting, hence $\tau^n \{x : |x| = 1\} \rightarrow \{e\}$.
- (3) Therefore $\|\tau^m\| \rightarrow 0$, hence there exists $m_0 \in \mathbb{N}$ with $\|\tau^m\| < 1$ for all $m \geq m_0$, and
- (4) $\|\tau^{-1}\| > 1$.

Example 3.8. Let, as in Example 3.4, \mathbb{G} be *totally disconnected* and contractible with contracting $\tau \in \text{Aut}(\mathbb{G})$. There exists a filtration (U_n) by open compact subgroups such that in addition $\tau U_n = U_{n+1}$, $n \in \mathbb{Z}$. We define for $0 < \alpha < 1$ the norm as in Example 3.4. Then by construction, $|\tau^n(x)| = \alpha^n \cdot |x|$, $x \in \mathbb{G}$. Therefore, for the canonical automorphism norm, we obtain $\|\tau^n\| = \alpha^n$. In particular, $\|\tau\| = \alpha < 1$ and $\|\tau^{-1}\| = \alpha^{-1} > 1$.

Lemma 3.9. Let \mathbb{G} be a contractible locally compact group with contracting $\tau \in \text{Aut}(\mathbb{G})$. Let $|\cdot|$ be a group norm and assume for some $m_0 \geq 1$, $\|\tau^m\| < 1$ for $m \geq m_0$. Then there exist $0 < r < R < 1$, $m_1 \geq m_0$, such that for $N \geq m_1$,

$$r^N \cdot |x| \leq |\tau^N(x)| \leq R^N \cdot |x|, \tag{3.1}$$

$$R^{-N} \cdot |x| \leq |\tau^{-N}(x)| \leq r^{-N} \cdot |x|. \tag{3.2}$$

Proof. Let $N \geq m_0$, $N = k \cdot m_0 + j$, $0 \leq j < m_0$. Then

$$|\tau^N(x)| \leq \|\tau^{m_0}\|^k \cdot |\tau^j(x)| \leq (\|\tau^{m_0}\|^{1/m_0})^k \cdot \|\tau^j\| \cdot |x| \leq C \cdot R_0^N \cdot |x|$$

for $R_0 := \|\tau^{m_0}\|^{1/m_0} < 1$ and $C \geq \max_{0 \leq j < m_0} (\|\tau^j\| R_0^{-j}) \geq 1$.

Hence $|\tau^N(x)| \leq (C^{1/N} \cdot R_0)^N |x|$. For sufficiently large $N \geq m_1 (\geq m_0)$ we have $C^{1/N} \cdot R_0 \leq C^{1/m_1} \cdot R_0 =: R_1 < 1$. Thus the right-hand side of (3.1) holds for $1 > R \geq R_1$. The left-hand side, as immediately seen, is valid for $0 < r \leq r_1 := \|\tau^{-1}\|^{-1}$ and all $N \in \mathbb{Z}_+$.

Again the right-hand side of (3.2) follows immediately for all $r^{-1} \geq r_1^{-1} := \|\tau^{-1}\| > 1$ and all $N \in \mathbb{Z}_+$. And almost verbatim as in the proof of (3.1), the left-hand side follows for $1 < R^{-1} \leq R_1^{-1}$.

Whence the assertion follows. □

According to Examples 3.7 and 3.8 the assumptions of Lemma 3.9 are satisfied for connected, resp. totally disconnected contractible groups. In that case the growth conditions (3.1) and (3.2) hold. Now let \mathbb{G} be a contractible locally compact group with contracting $\tau \in \text{Aut}(\mathbb{G})$. Then $\mathbb{G} = \mathcal{N} \otimes \mathbb{D}$, where \mathcal{N} is connected and \mathbb{D} totally disconnected with contracting $\rho \in \text{Aut}(\mathcal{N})$ and $\sigma \in \text{Aut}(\mathbb{D})$ respectively, such that $\tau = \rho \otimes \sigma$ (cf. Proposition 4.2, Corollary 4.3 of [30], Theorem 3.1.2 of [7]). According to Examples 3.7 and 3.8, there exist group-norms $|\cdot|_{\mathcal{N}}$ and $|\cdot|_{\mathbb{D}}$ on \mathcal{N} and \mathbb{D} adapted to ρ and σ respectively.

Then $|\cdot|$, defined by $|(x, y)| := |x|_{\mathcal{N}} + |y|_{\mathbb{D}}$, for $(x, y) \in \mathcal{N} \otimes \mathbb{D} = \mathbb{G}$ is a group norm on \mathbb{G} , adapted to τ . Hence we have

PROPOSITION 3.10

Let \mathbb{G} be a contractible locally compact group with contracting $\tau \in \text{Aut}(\mathbb{G})$. There exists a group norm $|\cdot|$ on \mathbb{G} , with corresponding automorphism norm satisfying the growth conditions (3.1) and (3.2).

In particular, \mathbb{G} is σ -compact, metrizable, hence second countable.

Proof. As mentioned above, \mathbb{G} splits as a direct product $\mathbb{G} = \mathcal{N} \otimes \mathbb{D}$ and also $\tau = \rho \otimes \sigma \cdot \rho$ and σ are both contracting on \mathcal{N} and \mathbb{D} respectively and satisfy the growth conditions (3.1) and (3.2) for $1 > R \geq R_i$ and $0 < r \leq r_i \leq R_i, i = 1, 2$. In case of \mathcal{N} , (3.1) and (3.2) hold for $R \geq R_1$ and $N \geq m_1 \geq 0$ (cf. Examples 3.7 and 3.8.)

With the group norm on \mathbb{G} defined as before and immediately verified, τ satisfies (3.1) and (3.2), with $1 > R \geq \max\{R_1, R_2\}, 0 < r \leq \min\{r_1, r_2\}$ and $N \geq m_1$.

\mathbb{G} is metrizable as $|\cdot|$ defines a metric, and σ -compact, as τ is compactly contracting. Whence the last assertion follows. □

For later use (in §6) we note:

PROPOSITION 3.11

(a) *Let \mathbb{G} be a connected contractible group with contracting $\tau \in \text{Aut}(\mathbb{G})$, therefore a (homogeneous, hence nilpotent and simply connected) Lie group of dimension d . Let $\mathcal{G} \cong \mathbb{R}^d$ denote the Lie algebra and let $\mathring{\tau} \in \text{Aut}(\mathcal{G}) \subseteq \text{GL}(\mathbb{R}^d)$ denote the differential of τ . $\exp^{-1} : \mathbb{G} \rightarrow \mathcal{G}$ is a C^∞ -homeomorphism satisfying*

$$\exp^{-1} \circ \tau = \mathring{\tau} \circ \exp^{-1}.$$

(b) *Let \mathbb{G} be a totally disconnected contractible group with contracting $\tau \in \text{Aut}(\mathbb{G})$. Then there exists $p \in \mathbb{N} \setminus \{1\}, \mathring{\tau} \in \text{Aut}(\mathcal{G})$ and a homeomorphism $\psi : \mathbb{G} \rightarrow \mathcal{G} \cong \mathbb{Q}_p$ satisfying*

$$\psi \circ \tau = \mathring{\tau} \circ \psi.$$

Here $\mathbb{G} = \mathbb{Q}_p$ denotes the additive group of p -adic numbers. (p is not assumed to be a prime power here.)

Proof.

(a) The assertion is obvious since \mathbb{G} is a simply connected nilpotent Lie group, hence homeomorphic to \mathbb{G} .

(b) Let (U_n) be a filtration by open compact subgroups adapted to τ . Put $p := \text{ord}(U_0/U_1)$. Proposition 4.2, Corollary 4.3 of [30] (resp. Remarks 3.1.8–3.1.10 of [7]) contains explicit constructions of the homeomorphism ψ and of $\overset{\circ}{\tau}$. In fact, the construction there is the following:

Let $\mathbb{Z}_p := \{0, \dots, p-1\}$ and let $\mathcal{P} \subseteq \bigotimes_{\mathbb{Z}} \mathbb{Z}_p$ denote the set of sequences $\{\vec{x} := (x(k))_{k \in \mathbb{Z}} : x(k) \in \mathbb{Z}_p, x(k) = 0 \forall k \leq K(\vec{x})\}$. Every subset $\Lambda_k := \{\vec{x} : x(j) = 0 \forall j \leq k\}$ is compact with respect to the product topology. And endowed with the topology that makes all cosets $y \cdot \Lambda_k := \{y(k)x(k) : \vec{x} \in \Lambda_k\}$ compact open, \mathcal{P} is a totally disconnected space. If we define the group operation co-ordinate-wise, \mathcal{P} becomes a locally compact group, which will be called Λ . This group is contractible, since the shift σ , defined by $(\sigma(\vec{x}))(k) := x(k+1)$, is a contracting automorphism with the property $\sigma(\Lambda_n) = \Lambda_{n+1}$. Furthermore, there exists a homeomorphism $\phi_1 : \mathbb{G} \rightarrow \Lambda$ such that $\phi_1 \circ \tau = \sigma \circ \phi_1$ and $\phi_1(U_n) = \Lambda_n$ for all n .

Let \mathbb{Q}_p denote the additive group of the ring of p -adic numbers, let $H_p : r \mapsto p \cdot r$ denote the homothetical transformation on \mathbb{Q}_p . We have $H_p \in \text{Aut}(\mathbb{Q}_p)$, in fact, H_p is contracting. If Δ_0 denotes the p -adic integers, then $\Delta_k := H_{p^k}(\Delta_0)$, $k \in \mathbb{Z}$, is a filtration of \mathbb{Q}_p with $p = \text{ord}(\Delta_k/\Delta_{k+1})$. (Note that \mathbb{Q}_p may be represented as the space \mathcal{P} defined above, with a different algebraic structure (see for e.g. Definition 10.2–Theorem 10.10 of [9]). H_p operates as shift on this product space.) Hence there exists a homeomorphism $\phi_2 : \mathbb{Q}_p \rightarrow \Lambda$ such that $\phi_2 \circ H_p = \sigma \circ \phi_2$ and $\phi_2(\Delta_k) = \Lambda_k$ for all k .

Hence with the composition $\psi : \mathbb{G} \xrightarrow{\phi_1} \Lambda \xrightarrow{\phi_2^{-1}} \mathbb{Q}_p$ and $\overset{\circ}{\tau} := H_p$ the assertion follows. (In fact, in Proposition 3.7 of [30] and Example 3.1.8 of [7], it is supposed that p is a prime power, hence \mathbb{Q}_p is a field. But the construction does not depend on this assumption.) \square

4. Convergence of convolution products on contractible groups

Recall that for locally compact groups some versions of Lévy's equivalence theorem are available. For our purposes we need the following versions:

PROPOSITION 4.1

Let \mathbb{G} be a locally compact group. (\mathbb{G} is assumed to be second countable, as random variables are involved. A superfluous condition, if \mathbb{G} is contractible).

(a) Let \mathbb{G} be aperiodic, i.e., without non-trivial compact subgroups. Then the classical version of the equivalence theorem holds:

If v_i , $i \geq 0$, ν , are probabilities and X_i , $i \geq 0$, Y are independent \mathbb{G} -valued random variables with distributions $X_i(P) = v_i$, $i \geq 0$, and $Y(P) = \mu$. Then the following assertions are equivalent:

- (i) $\prod_{i=0}^N X_i := X_0 \dots X_N \xrightarrow{N \rightarrow \infty} Y$ P -a.s.
- (ii) $\prod_{i=0}^N X_i \xrightarrow{N \rightarrow \infty} Y$ P -stochastically.

(iii) $\mu_N := \star_{i=0}^N v_i \xrightarrow{N \rightarrow \infty} \mu = Y(P)$ weakly.

(iii') For all $k \geq -1$, the products $\lambda_{k,N} := \star_{i=k+1}^N v_i$ are weakly convergent to λ_k (for $N \rightarrow \infty$) with $\lambda_{-1} = \mu$, and moreover, $\lambda_k = \star_{i=k+1}^\infty v_i \rightarrow \varepsilon_e$ (for $k \rightarrow \infty$).

(b) If \mathbb{G} is not aperiodic then still (i) \Rightarrow (ii) \Rightarrow (iii') \Rightarrow (iii) and (iii') \Rightarrow (i) hold.

For (iii) \Rightarrow (i) additional conditions are needed:

Let $H_r(\lambda) := \{x \in \mathbb{G} : \lambda \star \varepsilon_x = \lambda\}$ denote the right invariance group of $\lambda \in \mathcal{M}^1(\mathbb{G})$. If we assume in addition $H_r(\mu) = \{e\}$, then (iii) \Rightarrow (i).

Note that as \mathbb{G} is not supposed to be Abelian we have to define products in a fixed order: $\prod_0^N x_i := x_0 x_1 \cdots x_N$ for $x_i \in \mathbb{G}$, and analogously $\star_0^N v_i := v_0 \star \cdots \star v_N$ for probabilities v_i .

Proof. For any second countable locally compact group \mathbb{G} , the implications (i) \Rightarrow (ii) \Rightarrow (iii) are obvious. To prove (ii) \Rightarrow (iii') note that a.s. or stochastic convergence of $\prod_{i=0}^\infty X_i$ imply for any $k \in \mathbb{Z}_+$ convergence of $\prod_{i=k}^\infty X_i$ and moreover, $\prod_{i=k}^\infty X_i \xrightarrow{k \rightarrow \infty} e$. Thus (iii') follows.

Claim. (iii) implies that $\{\lambda_{k,n}\}_{0 \leq k \leq n}$ is relatively compact. A crucial condition in Theorems 2.2.16 and 2.2.19 of [10].

We have for $N, M > 0$: $\mu_{N+M} = \mu_N \star \lambda_{N,N+M}$. According to the shift compactness theorem (Ch. III, Theorems 2.1, 2.2 of [22], Theorem 1.2.21 of [10]), the set $\{\lambda_{N,N+M}\}$ is relatively compact. Let \mathcal{L} denote the set of accumulation points (for $N \rightarrow \infty$). Let $\lambda \in \mathcal{L}$ and assume, for some subsequence, $\lambda_{k_N, n_N} \rightarrow \lambda$. Then, for $K \in \mathbb{N}$, $k_N \geq K$ we have $\lambda_{K, n_N} = \lambda_{K, k_N} \star \lambda_{k_N, n_N}$.

(iii') \Rightarrow (i). By assumption (iii'), $\lambda_{K, n_N} \rightarrow \lambda_K := \star_{j=K+1}^\infty v_j$. Whence $\lambda_K = \lambda_K \star \lambda$ follows, i.e. $\text{supp } \lambda \subseteq H_r(\lambda_K)$. Again by assumption (iii'), $\lambda_K \rightarrow \varepsilon_e$, hence $\varepsilon_e = \varepsilon_e \star \lambda$, i.e. $\lambda = \varepsilon_e$ and $\mathcal{L} = \{\varepsilon_e\}$. According to Theorem 2.2.16 of [10], (i) follows.

Assume (iii), and assume further that \mathbb{G} is aperiodic. As mentioned before, $\{\lambda_{k,n}\}$ is relatively compact. Thus, according to Theorem 2.2.19 of [10], (i) follows.

Assume (iii), and in addition, $H_r(\mu) = \{e\}$. As before, for $\lambda \in \mathcal{L}$ we obtain $\mu = \mu \star \lambda$. Again $\text{supp}(\lambda) \subseteq H_r(\mu)$ and thus, by assumption, $\lambda = \varepsilon_e$. Again, according to Theorem 2.2.16 of [10], (i) follows. □

Note that $H_r(\mu) = \{e\} \Rightarrow H_r(\lambda_k) = \{e\}$ and, if the measures v_i belong to a commutative sub-semigroup \mathcal{S} of $\mathcal{M}^1(\mathbb{G})$, $H_r(\lambda_{k,n}) = \{e\}$.

For special products considered in the sequel, $\mu = \star_{i \geq 0} \tau^i(v)$, the condition $H_r(\mu) = \{e\}$ is superfluous.

COROLLARY 4.2

Let \mathbb{G} be a contractible locally compact group with contracting $\tau \in \text{Aut}(\mathbb{G})$.

Let $X_i, i \geq 0, Y$ be independent random variables with distribution $v = X_i(P), i \geq 0, \mu := Y(P)$. Then, with the notations as above, $\mathcal{L} = \{\varepsilon_e\}$ and hence

$$\prod_{i=0}^N \tau^i(X_i) \rightarrow Y \text{ a.s.} \quad \text{iff} \quad \star_{i=0}^N \tau^i(v) \rightarrow \mu \text{ weakly.}$$

Proof. It suffices to show that condition (i) of Proposition 4.1 is satisfied. If $\mu_N := \star_{j=0}^N \tau^j(v) \xrightarrow{N \rightarrow \infty} \mu$. Then for all K , $\lambda_{K,N} := \star_{j=K+1}^N \tau^j(v) \xrightarrow{N \rightarrow \infty} \tau^{K+1}(\mu) = \lambda_K$. As τ is contracting, $\tau^{K+1}(\mu) = \lambda_K \rightarrow \varepsilon_e$ follows. $\lambda \in \mathcal{L}$, let $\lambda_{K_n, N_n} \rightarrow \lambda$. As $\lambda_{K_n} = \lambda_{K_n, N_n} \star \lambda_{N_n} \rightarrow \varepsilon_e$ and $\lambda_{N_n} \rightarrow \varepsilon_e$, it follows that $\lambda_{K_n, N_n} \rightarrow \varepsilon_e$, thus $\lambda = \varepsilon_e$, by the shift compactness theorem (Ch. III, Theorems 2.1, 2.2 of [22], Theorem 1.2.21 of [10]). Hence (iii') and thus (i) (in Proposition 4.1) follow. \square

Remark 4.3. We define *ad hoc*: A infinite convolution product $\mu = \star_{j \geq 0} v_j$ converges strongly if all rest products $\lambda_k := \star_{j > k} v_j$ are convergent ($k \geq -1$) and in addition $\lambda_k \xrightarrow{k \rightarrow \infty} \varepsilon_e$, i.e. if condition (iii') in Proposition 4.1 is satisfied. A product of convolution powers $\mu = \star_{j \geq 0} v_j^{k(j)}$ converges strongly if it converges strongly in lexicographical order.

- (a) As shown in Proposition 4.1, (i) \Leftrightarrow (iii'), strong convergence implies that the equivalence theorem holds.
- (b) Moreover, we define *ad hoc* for a fixed group norm $|\cdot|$: If (X_j) are independent \mathbb{G} -valued random variables, with distributions $X_i(P) = v_i$, then $\prod X_j$ is a.s. absolutely convergent if $\sum |X_i| < \infty$ a.s. Analogously we define for independent arrays $(X_{i,j})$ with distributions $X_{i,j}(P) = v_j$ that the iterated products $\prod_{j \geq 0} \prod_{i=1}^{k(j)} X_{i,j}$ converge a.s. absolutely if $\sum_j \sum_{i=1}^{k(j)} |X_{i,j}| < \infty$ a.s.

A.s. absolute convergence implies strong convergence. In fact, a.s. *absolute* convergence implies a.s. convergence of all partial products and re-arranged products of the above products, and hence weak convergence of the convolution products $\star_{j \geq 0} v_j$ resp. $\star_{j \geq 0} v_j^{k(j)}$. Moreover, $\sum_{j > K} \sum_{i=0}^{k(j)} |X_{i,j}| \rightarrow 0$ a.s. implies that the rest products $\star_{j > K} v_j^{k(j)}$ converge to ε_e ($K \rightarrow \infty$).

As is well known, on vector spaces and also on homogeneous groups, convolution products $\star_{k+1}^\infty \tau^j(v)$ converge iff logarithmic moments exist. As in the case of homogeneous groups (cf. [6, 7]) we define logarithmic moments in terms of group norms. (For vector spaces, see for e.g. [11].)

DEFINITION 4.4

Let \mathbb{G} be a locally compact group with a group norm $|\cdot|$. Let (Ω, Σ, P) be a probability space. Let X be a \mathbb{G} -valued random variable with distribution $X(P) = v$. X resp. v possesses finite logarithmic moments of order n , symbolically $v \in \mathcal{P}_{\log_+^n}$, if $\mathbb{E}((\log_+ |X|)^n) = \int_{\mathbb{G}} (\log_+ |x|)^n dv(x) < \infty$, equivalently, if $\int (\log_+(1 + |x|))^n dv(x) < \infty$.

Obviously, we have: Let $f : \Omega \rightarrow \mathbb{R}_+$ be measurable, $\psi \in C^1(\mathbb{R})$ strictly increasing with integrable ψ' . Then

$$\int_{\Omega} \psi \circ f dP = \int_{\mathbb{R}_+} P \{f > y\} \psi'(y) dy.$$

Applying this observation to the functions $t \mapsto \log_+(t)$ resp. $t \mapsto \log(1 + t)$ we obtain

Lemma 4.5. $\nu \in \mathcal{P}_{\log_+^{n+1}}$ iff one of the following equivalent conditions is satisfied:

$$\begin{aligned} \mathbb{E}(\log(1 + |X|)^{n+1}) < \infty &\Leftrightarrow \int_0^\infty P(|X| > e^y - 1)y^n dy < \infty \\ &\Leftrightarrow \int_0^\infty \nu\{|\cdot| > e^y - 1\}y^n dy < \infty \\ &\Leftrightarrow \mathbb{E}(\log_+(|X|)^{n+1}) < \infty \\ &\Leftrightarrow \int_1^\infty P(|X| > e^y)y^n dy < \infty \\ &\Leftrightarrow \int_1^\infty \nu\{|\cdot| > e^y\}y^n dy < \infty. \end{aligned}$$

Easy calculations show as follows:

Lemma 4.6. Let X be a \mathbb{G} -valued random variable and $|\cdot|$ a group norm. Then,

$$\int_1^\infty P\{\log_+ |X| > y\}y^n dy < \infty \Rightarrow \sum_1^\infty P\{\log_+ |X| > j\beta\}j^n < \infty$$

for all $\beta > 0$. And if for some $\beta > 0$,

$$\sum_1^\infty P\{\log_+ |X| > j\beta\}j^n < \infty, \text{ then } \int_1^\infty P\{\log_+ |X| > y\}y^n dy < \infty.$$

(Note that $f_1 : y \mapsto P(\log_+ |X| > y)$ is decreasing and $f_2 : [1, \infty) \ni y \mapsto y^n$ is increasing. This yields upper and lower estimates for the Riemann sums approximating $\int_1^\infty f_1 \cdot f_2 dy$.)

PROPOSITION 4.7

Let \mathbb{G} be a contractible locally compact group with contracting $\tau \in \text{Aut}(\mathbb{G})$ and group norm $|\cdot|$ satisfying the growth condition (3.1). Let, for fixed $n \in \mathbb{N}$, $k(j)$, $j \geq 1$, be a sequence in \mathbb{N} such that $k(j)/j^n$ is bounded. (E.g., mostly in the sequel we have $k(j) = \binom{n+j}{n}$, as throughout in the sequel). Then we have

$$\nu \in \mathcal{P}_{\log_+^{n+1}} \Rightarrow \mu := \star_{j \geq 0} \tau^j(\nu^{k(j)}) \text{ is weakly convergent.}$$

Moreover, let $(X_{i,j}) : 1 \leq i \leq k(j)$, $j \geq 0$, $(X_i) : i \geq 0$, be i.i.d. arrays with distributions $X_{i,j}(P) = \nu$, $X_i(P) = \nu$. Then

$$\sum_{j \geq 0} \sum_{i=1}^{k(j)} |\tau^j(X_{i,j})| < \infty \text{ a.s. and } \sum_{j \geq 0} k(j) |\tau^j(X_j)| < \infty \text{ a.s.} \tag{4.1}$$

In particular, $\prod_{j \geq 0} \prod_{i=1}^{k(j)} \tau^j(X_{i,j})$ is a.s. convergent and, re-arranged products and partial products are convergent.

(For real vector spaces see for e.g. Lemma 3.6.5 of [11], for $n = 0$, or [16] for $n \geq 1$.)

Proof.

(1) Assume first that the growth condition (3.1) holds true with $m_1 = 1$.

Let $X, X_i, X_{i,j}$ be i.i.d. random variables with distribution ν . Assume $\nu \in \mathcal{P}_{\log_+^{n+1}}$ for some (fixed) $n \geq 0$. As mentioned before, in Lemma 4.5, we have $\int_0^\infty P(\log_+ |X| > y) y^n dy < \infty$. And, according to Lemma 4.6, for any $\alpha > 0$ this is equivalent to

$$\begin{aligned} \sum_{j=0}^{\infty} j^n P(\log_+ |X| > j\alpha) < \infty &\Leftrightarrow \sum_{j=0}^{\infty} j^n P(|X| > (e^\alpha)^j) < \infty \\ &\Leftrightarrow I := \sum_{j=0}^{\infty} \sum_{i=1}^{j^n} P(|X_{i,j}| > (e^\alpha)^j) < \infty. \end{aligned}$$

The growth condition (3.1) yields $r^n \cdot |x| \leq |\tau^n x| \leq R^n \cdot |x| \forall x$ and hence for all $\delta > 0$,

$$\{|\tau^n x| \geq r^n \delta\} \supseteq \{|x| \geq \delta\} \supseteq \{|\tau^n x| \geq R^n \delta\}.$$

Inserting $\delta = e^{\alpha j}$ yields

$$\begin{aligned} P(|X_{i,j}| \geq (e^\alpha)^j) &\leq P(|\tau^j(X_{i,j})| \geq (r \cdot e^\alpha)^j), \\ P(|X_{i,j}| \geq (e^\alpha)^j) &\geq P(|\tau^j(X_{i,j})| \geq (R \cdot e^\alpha)^j) \end{aligned}$$

and thus

$$\begin{aligned} \sum_{j=0}^{\infty} \sum_{i=1}^{j^n} P(|\tau^j(X_{i,j})| \geq (R e^\alpha)^j) \\ \leq I \leq \sum_{j=0}^{\infty} \sum_{i=1}^{j^n} P(|\tau^j(X_{i,j})| \geq (r e^\alpha)^j). \end{aligned}$$

Hence, with $\delta_\alpha := R \cdot e^\alpha$ we obtain $\sum_{j=0}^{\infty} \sum_{i=1}^{j^n} P(|\tau^j(X_{i,j})| \geq \delta_\alpha^j) < \infty$. Put, for $1 \leq i \leq j^n$, $j \geq 1$ and fixed $\alpha > 0$, $A_{i,j} := \{|\tau^j(X_{i,j})| \geq \delta_\alpha^j\}$. Then, by the lemma of Borel–Cantelli, $P(A^*) = 0$, where $A^* := \limsup A_{k,\ell}$ (in lexicographical order), i.e., considering $\mathbb{C}A^*$, we obtain: For P -almost all ω there exists $j(\omega)$ such that for all $i \leq j^n$, $j \geq j(\omega)$ we have: $\omega \in \mathbb{C}A_{i,j}$, i.e., $|\tau^j(X_{i,j}(\omega))| < \delta_\alpha^j$.

Hence, for small α , such that $\delta_\alpha = R e^\alpha < 1$,

$$\begin{aligned} \sum_{j=0}^{\infty} \sum_{i=1}^{j^n} |\tau^j(X_{i,j})| &= \sum_{j=0}^{j(\omega)} \sum_{i=1}^{j^n} |\tau^j(X_{i,j})| + \sum_{j>j(\omega)} \sum_{i=1}^{j^n} |\tau^j(X_{i,j})| \\ &\leq c_1(\omega) + \sum_{j \geq 0} \sum_{i=1}^{j^n} \delta_\alpha^j \\ &= c_1(\omega) + \sum_{j \geq 0} j^n \delta_\alpha^j < \infty \quad P - \text{a.s.} \end{aligned}$$

with some constant $c_1(\omega)$.

Analogously, $\sum_{j \geq 0} j^n |\tau^j(X_{i,j})| \leq c_2(\omega) + \sum_{j \geq 0} j^n \delta_\alpha^j < \infty$ P -a.s.

(2) In the case $m_1 > 1$ the proof runs analogously and the above estimates hold true only for $j \geq m_1$. Hence in the estimates for the sums $\sum_{j=0}^{\infty} \sum_{i=1}^{j^n} |\tau^j(X_{i,j})|$ etc., additional additive constants (depending on ω) occur.

A.s. absolute convergence yields that $\prod_{j \geq 0} \prod_{i=1}^{j^n} \tau^j(X_{i,j})$ as well as all partial products and re-arranged products converge a.s. (Remark 4.3). In particular, weak convergence of $\star_{j \geq 0} \tau^j(v^{k(j)})$ follows.

By assumption, $k(j) \leq C \cdot j^n$, whence $\sum_{j=0}^{\infty} \sum_{i=1}^{k(j)} |\tau^j(X_{i,j})| < \infty$ P -a.s. follows. \square

Converse results. For homogeneous groups we obtain as in the case of vector spaces (in analogy to e.g. [16, 17]):

PROPOSITION 4.8

Let \mathbb{G} be a homogeneous group with contracting $\tau \in \text{Aut}(\mathbb{G})$ and corresponding group norm $|\cdot|$. Let again $k(j) \approx j^n$, e.g. $k(j) = \binom{n+j}{n}$. Then we have

$$\star_{j \geq 0} \tau^j(v^{k(j)}) \text{ converges weakly} \Rightarrow v \in \mathcal{P}_{\log_+^{n+1}}.$$

Proof. \mathbb{G} is a homogeneous group, hence aperiodic. According to the equivalence theorem, Proposition 4.1, weak convergence of the products $\star_{j \geq 0} \tau^j(v^{k(j)})$ implies a.s. convergence of $\prod_{j \geq 0} \prod_{i=1}^{k(j)} \tau^j(X_{i,j})$ for iid random variables with distribution $X_{i,j}(P) = \nu$. Hence we obtain

$$P \left(\bigcap_{n \geq 0} \left(\bigcup_{j \geq n, i \leq k(j)} \{|\tau^j X_{i,j}| > \delta\} \right) \right) = 0 \text{ for all } \delta > 0.$$

According to the lemma of Borel–Cantelli, it follows that for all $\delta > 0$,

$$I := \sum_{j=0}^{\infty} \sum_{i=0}^{j^n} P(|\tau^j X_{i,j}| > \delta) = \sum_{j \geq 0} j^n P(|\tau^j X_{i,j}| > \delta) < \infty$$

since the variables $X_{i,j}$ are identically distributed and $k(j) \approx j^n$. The growth condition (3.1) yields (for all i)

$$\{|\tau^j X_{i,j}| > \delta\} = \{|\tau^j X_{i,j}| > r^j r^{-j} \delta\} \supseteq \{|X_{i,j}| > r^{-j} \delta\}.$$

We have $r^{-1} > 1$. Choose $\delta > 1$ and put $\gamma := r^{-1} \delta$, $\gamma_j := \delta^{1/j} r^{-1} \leq \gamma$. Since $P(|X| > \gamma_j^j) \geq P(|X| > \gamma^j)$, $j \in \mathbb{Z}_+$, it follows that

$$\infty > I \geq \sum_{j \geq 0} j^n P(|X_{i,j}| > \delta r^{-j} = \gamma_j^j) = \sum_{j \geq 0} j^n P(|X_{i,j}| > \gamma^j),$$

and hence (with $\beta = \log \gamma \geq 0$)

$$\infty > I \geq \sum_{j \geq 0} j^n P(\log |X_{i,j}| > \beta j).$$

Thus, by Lemma 4.6, $\int_1^\infty P(\log_+ |X| > y) y^n dy < \infty$, whence $\nu \in \mathcal{P}_{\log_+^{n+1}}$. \square

The proof of Proposition 4.8 shows that the following more general result holds true.

PROPOSITION 4.9

Let \mathbb{G} be a contractible locally compact group with contracting $\tau \in \text{Aut}(\mathbb{G})$, and let $|\cdot|$ be a group norm adapted to τ and satisfying (3.1). Let $k(j) \approx \binom{n+j}{n}$. Let $(X_{i,j})_{i,j \geq 0}$ be independent random variables with distribution ν . Assume that convergence of the convolution product $\mu = \star_{j=0}^\infty \tau^{k(j)}(\nu)$ implies a.s. convergence of $\prod_{j=0}^\infty \prod_{i=0}^{k(j)} \tau^j(X_{i,j})$. Then we have $\mu \in \mathcal{P}_{\log_+^{n+1}}$.

The condition is satisfied in the case $n = 0$ (Corollary 4.2), or if the product converges strongly (in the sense of Remark 4.3) or if $H_r(\mu) = \{e\}$, (or if \mathbb{G} is a homogeneous group, as in Proposition 4.8).

Proof. Analogous to the proof of Proposition 4.8, \mathbb{G} was supposed to be a homogeneous group just in order to apply the equivalence theorem, Proposition 4.1, (iii) \Rightarrow (i). Note that for $n = 0$, weak and a.s. convergence are equivalent by Corollary 4.2. Analogously, if strong convergence or $H_r(\mu) = \{e\}$ or, more generally, $\mathcal{L} = \{\varepsilon_e\}$ is assumed, a.s. convergence follows by Remark 4.3 (resp. Proposition 4.1). \square

Remark 4.10.

(a) We have shown by the preceding Propositions 4.7–4.8 that for homogeneous groups, $\forall n \in \mathbb{Z}_+$, for any group norm – not only homogeneous norm – satisfying a growth condition (3.1) the existence of logarithmic moments $\nu \in \mathcal{P}_{\log_+^{n+1}}$ characterizes weak convergence $\star_{j \geq 0} \tau^j \left(\nu \binom{n+j}{n} \right)$. This generalizes Proposition 3.4, Theorem 3.5 of [6], Proposition 2.14.24 of [7], for the case $n \geq 0$. In particular, this proves that $\mathcal{P}_{\log_+^{n+1}}$ does not depend on the particular chosen group norm.

Analogously, for $n = 0$, Proposition 4.9 generalizes the above mentioned results in [6, 7], to general contractible locally compact groups. Again this implies that \mathcal{P}_{\log_+} does not depend on the particular chosen group norm.

(b) A.s. absolute convergence, $\sum_{j \geq 0} \sum_{i=1}^{\binom{n+j}{n}} |\tau^j(X_{i,j})| < \infty$ a.s., implies that weak convergence of re-arranged products and of partial products of $\star_j \tau^j \left(\nu \binom{n+j}{n} \right)$. (cf. Propositions 4.7, eq. (4.1)) We therefore obtain:

(b1) Let \mathbb{G} be contractible with contracting $\tau \in \text{Aut}(\mathbb{G})$. Assume that $\nu_{(k)} := \star_{j \geq 0} \tau^j \left(\nu \binom{k+j}{k} \right)$ is convergent for all $k \leq n - 1$. Let $(Z_j^{(k)})$ be the corresponding random variables with distributions $Z_j^{(k)}(P) = \nu_{(k)}$. Then we have by Corollary 4.2,

resp. Proposition 4.8: $\prod_{j \geq 0} \tau^j(Z_j^{(k)})$ converges a.s. absolutely for all $k \leq n-1$, i.e., $\sum_{j \geq 0} |\tau^j(Z_j^{(k)})| < \infty$ a.s.

(b2) Furthermore, let $(Z^{(n)})$, $(Z_j^{(n-1)})$, $(Z_{i,j}^{(n-2)})$, ... be sequences of independent random variables with distributions $\nu_{(n)}$, $\nu_{(n-1)}$, ... Then

$$\begin{aligned} Z^{(n)} &= \prod_{j \geq 0} \tau^j(Z_j^{(n-1)}) = \prod_{j \geq 0} \tau^j \prod_{i \geq 0} \tau^i(Z_{j,i}^{(n-2)}) \\ &= \dots = \prod_{j_n \geq 0} \tau^{j_n} \left(\prod_{j_{n-1} \geq 0} \tau^{j_{n-1}} \left(\dots \left(\prod_{j_0 \geq 0} \tau^{j_0} (Z_{j_n, \dots, j_0}^{(0)}) \dots \right) \right) \right) \\ &= \prod_{j_n \geq 0} \left(\prod_{j_{n-1} \geq 0} \left(\dots \left(\prod_{j_0 \geq 0} \tau^{\sum_0^n j_i} (Z_{j_n, \dots, j_0}^{(0)}) \dots \right) \right) \right). \end{aligned}$$

(b3) If in (b2), \mathbb{G} is a homogeneous group or if one of the conditions in Proposition 4.9 is satisfied, then $\nu \in \mathcal{P}_{\log^+_{n+1}}$ and Proposition 4.7 a.s. convergence of the random variables $\sum_{j_n, \dots, j_0} |\tau^{\sum j_i} (X_{j_n, \dots, j_0})|$ follows.

5. Multiple τ -decomposability of probabilities

In this section, let \mathbb{G} be a contractible locally compact group with contracting $\tau \in \text{Aut}(\mathbb{G})$. Furthermore, let $|\cdot|$ be a group norm adapted to τ satisfying the growth condition (3.1).

DEFINITION 5.1

(a) $\mu \in \mathcal{M}^1(\mathbb{G})$ is τ -decomposable – in symbols, $\mu \in L_0(\tau)$ – with cofactor $\nu \in \mathcal{M}^1(\mathbb{G})$ if $\mu = \nu \star \tau(\mu)$.

In terms of random variables, a random variable X with distribution μ is τ -decomposable if there exists a random variable Y , such that (X, Y) are independent and $X \stackrel{d}{=} Y \cdot \tau(X)$ (equality of distributions).

(b) Put $L_{-1}(\tau) := \mathcal{M}^1(\mathbb{G})$. $\mu \in \mathcal{M}^1(\mathbb{G})$ is n -times τ -decomposable – in symbols, $\mu \in L_n(\tau)$ – if, μ is τ -decomposable, $\mu = \nu_{(n)} \star \tau(\mu)$, with cofactor $\nu_{(n)} \in L_{n-1}(\tau)$.

Therefore, $\mu = \nu_{(n)} \star \tau(\mu)$, $\nu_{(n)} = \nu_{(n-1)} \star \tau(\nu_{(n)})$, ..., $\nu_{(1)} = \nu \star \tau(\nu_{(1)})$. Hence $\mu = \nu \star \tau(\nu_{(1)}) \star \dots \star \tau(\nu_{(n)}) \star \tau(\mu)$.

If the contraction condition is not supposed, μ is usually called *strongly* τ -decomposable if in addition $\tau^k(\nu) \rightarrow \varepsilon_e$. Here, as we always restrict to contracting τ , ‘strongly’ is superfluous.

PROPOSITION 5.2

(a) μ is τ -decomposable with cofactor ν iff

$$\mu = \star_{j=0}^{\infty} \tau^j(\nu) := \lim_{N \rightarrow \infty} \nu \star \dots \star \tau^N(\nu).$$

(b) μ is n -times τ -decomposable iff

$$\mu = \star_{j_n \geq 0} \left(\star_{j_{n-1} \geq 0} \left(\dots \star_{j_0 \geq 0} \tau^{\sum_0^n j_i}(\nu) \dots \right) \right).$$

Proof.

- (a) \Rightarrow . Obviously, since for all N we have $\star_0^N \tau^j(v) \star \tau^{N+1}(\mu) = \mu$ and, since τ is contracting, $\tau^{N+1}(\mu) \rightarrow \varepsilon_e$. The assertion follows by the shift compactness theorem [10, 22].
 \Leftarrow . Assume $\mu = \star_0^\infty \tau^k(v)$. Hence $\mu = v \star \star_1^\infty \tau^k(v) = v \star \tau(\mu)$. (For homogeneous groups see [6, 7].)
- (b) Follows by inserting in (a) successively $\nu_{(i)} = \star_0^\infty \tau^k(\nu_{(i-1)})$, $\nu_{(0)} = \nu$. \square

Hence according to Propositions 4.7, 4.8, 4.9 we immediately obtain:

PROPOSITION 5.3

$\nu \in \mathcal{M}^1(\mathbb{G})$ is the cofactor of a τ -decomposable law iff $\nu \in \mathcal{P}_{\log_+}$.

We mention two results concerning the structure of τ -decomposable laws. The proofs are left to the reader.

PROPOSITION 5.4

- (a) The set of τ -decomposable laws is weakly closed.
 (b) If μ_i are τ -decomposable with cofactors ν_i and if $\mu_i, \nu_i, i = 1, 2$, belong to a commutative sub-semigroup $\mathcal{S} \subseteq \mathcal{M}^1(\mathbb{G})$, with $\tau(\mathcal{S}) \subseteq \mathcal{S}$, then $\mu_1 \star \mu_2$ is τ -decomposable with cofactor $\nu_1 \star \nu_2$.

The following lemma will be essential in the sequel. Since compact subgroups are not excluded, a careful discussion of convergence and re-arranging of iterated convolution products seems to be necessary.

Lemma 5.5. Let, as before, \mathbb{G} be contractible with contracting $\tau \in \text{Aut}(\mathbb{G})$, let $\mathcal{S} \subseteq \mathcal{M}^1(\mathbb{G})$ be a commutative sub-semigroup with $\tau(\mathcal{S}) \subseteq \mathcal{S}$. Let $\mu, \nu \in \mathcal{S}$. Let $n \in \mathbb{N}$. If $n > 1$ we assume in addition $H_r(\mu) = \{e\}$. Then the following assertions are equivalent:

$$\mu = \star_{j_n \geq 0} \left(\dots \left(\star_{j_0 \geq 0} \tau^{\sum_0^n j_i}(\nu) \right) \dots \right), \quad (5.1)$$

$$\mu = \star_{j \geq 0} \tau^j(\nu^{k_n(j)}) \quad \text{with} \quad k_n(j) = \binom{n+j}{n}. \quad (5.2)$$

Proof.

(5.1) \Rightarrow (5.2). We proceed by induction on n . Let $n = 1$. Then $\mu = \star_{j \geq 0} \tau^j(\star_{i \geq 0} \tau^i(\nu)) = \star_{j \geq 0} \tau^j(\nu_{(1)})$ and $\nu_{(1)} := \star_{i \geq 0} \tau^i(\nu)$ are both τ -decomposable. Furthermore, $\nu, \nu_{(1)} \in \mathcal{P}_{\log_+}$ (Propositions 4.7, 4.9). For all $L, M \in \mathbb{N}$ we have

$$\mu = \star_{i=0}^L \tau^i(\nu_{(1)}) \star \tau^{L+1}(\mu), \quad (5.3)$$

$$\nu_{(1)} = \star_{i=0}^M \tau^i(\nu) \star \tau^{M+1}(\nu_{(1)}). \quad (5.4)$$

Hence for all $L \in \mathbb{N}$, inserting successively (5.4) in (5.3) with $M = L - \ell$ we obtain (in view of $\nu \in \mathcal{S}$):

$$\begin{aligned} \mu &= \star_{\ell=0}^L \tau^\ell \left(\star_{j=0}^{L-\ell} \tau^j(\nu) \star \tau^{L-\ell+1}(\nu_{(1)}) \right) \star \tau^{L+1}(\mu). \\ &=: \lambda_L^{(1)} \star \tau^{L+1} \left(\nu_{(1)}^{L+1} \right) \star \tau^{L+1}(\mu), \end{aligned}$$

where $\lambda_L^{(1)} = \star_{\ell=0}^L \tau^\ell \left(\star_{j=0}^{L-\ell} \tau^j(\nu) \right) = \star_{i=0}^L \left(\star_{j=0}^i \tau^j(\nu) \right) = \star_{i=0}^L \tau^i(\nu^{i+1})$ and $\tau^{L+1}(\mu) \rightarrow \varepsilon_e$. We have to show that $\lambda_L^{(1)} \rightarrow \mu$, equivalently, $\tau^{L+1}(\nu_{(1)}^{L+1}) \rightarrow \varepsilon_e$. Hence $\mu = \star_0^\infty \tau^i(\nu^{i+1})$, i.e. (5.2) for $n = 1$.

Since $\nu_{(1)} \in \mathcal{P}_{\log_+}$, for a i.i.d. sequence (Y_j) of random variables with distribution $\nu_{(1)}$ we obtain $\sum |\tau^j(Y_j)| < \infty$ a.s. (Proposition 4.7, eq. (4.1)). τ satisfies the growth condition (3.1), $\|\tau^k\| \leq R^k$, $k \geq m_1$, for $0 < R < 1$. Let $C := \max \|\tau^k\|$ (≥ 1). Let $L+1 = N_1 + N_2$, $N_i > m_1$. Then we obtain

$$\begin{aligned} \left| \tau^{L+1} \left(\prod_0^{L+1} Y_j \right) \right| &\leq \sum_0^{N_1} |\tau^{L+1}(Y_j)| + \sum_{N_1+1}^{L+1} |\tau^{L+1}(Y_j)| \\ &\leq R^{N_2} \sum_0^{N_1} |\tau^{N_1}(Y_j)| + \sum_{N_1+1}^{L+1} |\tau^{L+1}(Y_j)| \\ &\leq R^{N_2} \cdot C \cdot \left(\sum_0^\infty |\tau^j(Y_j)| \right) \\ &\quad + C \cdot \left(\sum_{N_1+1}^\infty |\tau^j(Y_j)| \right) \xrightarrow{(N_1, N_2) \rightarrow \infty} 0 \quad \text{a.s.} \end{aligned}$$

In fact, we have $|\tau^{N_1}(Y_j)| \leq \|\tau^{N_1-j}\| |\tau^j(Y_j)| \leq C \cdot |\tau^j(Y_j)|$ for $j \leq N_1$, and $|\tau^{L+1}(Y_j)| \leq \|\tau^{L+1-j}\| |\tau^j(Y_j)| \leq C \cdot |\tau^j(Y_j)|$ for $N_1 < j \leq L+1$.

Let the assertion be proved for $k \leq n$. We assumed that $H_r(e) = \{e\}$, hence Proposition 4.9 is applicable. Let $\nu_{(0)} = \nu, \dots, \nu_{(i)} = \star_{k \geq 0} \tau^k(\nu_{(i-1)})$, $\nu_{(n+1)} = \mu$. By assumption, $\mu = \star_{j=0}^\infty \tau^j(\nu_{(n)})$ and by induction hypothesis, $\nu_{(n)} = \star_{j=0}^\infty \tau^j(\nu^{k_n(j)})$ with $k_n(j) = \binom{n+j}{n}$. Hence $\mu \in \mathcal{P}_{\log_+}$ and $\nu_{(n)} \in \mathcal{P}_{\log_+^{n+1}}$.

Hence for all arrays $(X_{i,j})$ of i.i.d. random variables with distribution ν we have a.s. absolute convergence, $\sum_{j \geq 0} \sum_{i=0}^{k_n(j)} |\tau^j(X_{i,j})| < \infty$ a.s. Furthermore, for all $L > 0$, we have

$$\mu = \star_{\ell=0}^L \tau^\ell(\nu_{(1)}) \star \tau^{L+1}(\mu) =: \mu_L \star \tau^{L+1}(\mu)$$

and (since all measures belong to \mathcal{S})

$$\mu_L = \star_{\ell=0}^L \tau^\ell \left(\star_{j=0}^{L-\ell} \tau^j(\nu^{k_n(j)}) \right) \star \star_{\ell=0}^L \tau^\ell \left(\star_{j>L-\ell} \nu^{k_n(j)} \right) =: \lambda_L^{(n)} \star \beta_L.$$

As in the case $n = 1$, we observe

$$\lambda_L^{(n)} = \star_{\ell=0}^L \star_{j=0}^{\ell} \tau^{\ell} (\nu^{k_n(j)}) = \star_{\ell=0}^L \tau^{\ell} (\nu^{\sum_{i=0}^{\ell} k_n(i)}) = \star_{\ell=0}^L \tau^{\ell} (\nu^{k_{n+1}(\ell)}).$$

Hence we have to show that $\beta_L \rightarrow \varepsilon_e$. Then $\lambda_L^{(n)} \rightarrow \mu$, hence (5.2) is proved.

Let Z_L be distributed according to β_L . Since $\nu \in \mathcal{S}$, we may assume that $Z_L = \prod_{\ell=0}^L \tau^{\ell} \left(\prod_{j=L-\ell+1}^{\infty} \tau^j \left(\prod_{i=0}^{k_n(j)} X_{i,j} \right) \right)$.

We have $|Z_L| \leq \sum_{\ell=0}^L C R^{\ell} \left(\sum_{j>L-\ell} \sum_{i=0}^{k_n(j)} |\tau^j(X_{i,j})| \right) =: U_L$. Put for $K \in \mathbb{Z}_+$, $f_K := \sum_{j \geq K} \sum_{i=0}^{k_n(j)} |\tau^j(X_{i,j})|$. Recall that $f_K \searrow 0$ a.s., in particular, $f_K \leq f_0 < \infty$ a.s. As in the case $n = 1$, let $L = N_1 + N_2$ with $N_i > m_1$ and decompose $U_L = C \cdot \sum_{\ell=0}^{N_1} R^{\ell} f_{L-\ell} + C \cdot \sum_{\ell=N_1+1}^L R^{\ell} f_{L-\ell} =: U_{N_1}^L + V_{N_1}^L$. In view of $U_{N_1}^L \leq C \cdot (\sum_{\ell \geq 0} R^{\ell}) \cdot f_{N_2}$ and $V_{N_1}^L \leq C \cdot (\sum_{\ell \geq N_1+1} R^{\ell}) \cdot f_0$ we obtain $U_L \xrightarrow{(N_1, N_2) \rightarrow \infty} 0$ a.s., hence $Z_L \rightarrow e$ a.s., and thus $\beta_L \rightarrow \varepsilon_e$. The assertion (5.1) \Rightarrow (5.2) is proved. Note furthermore that (5.2) implies $\nu \in \mathcal{P}_{\log_+^{n+2}}$ (Proposition 4.9).

(5.2) \Rightarrow (5.1). Assume $\mu = \star_{j \geq 0} \tau^j (\nu^{k_n(j)})$. As before, $\nu \in \mathcal{P}_{\log_+^{n+1}}$ follows, and for all i.i.d. arrays $(X_{i,j})$ with distribution ν we have a.s. absolute convergence, $\sum_{j \geq 0} \sum_{i=0}^{k_n(j)} |\tau^j(X_{i,j})| < \infty$ a.s.

Let (Y_{j_0, \dots, j_n}) be i.i.d. random variables with distribution ν . Let, for fixed j , $\{X_{i,j} : 1 \leq i \leq k_n(j)\}$ denote the re-arranged random variables $\{Y_{j_0, \dots, j_n} : \sum_0^n j_i = j\}$. Then $\sum_{j_n} \dots \sum_{j_0} |\tau^{\sum_0^n j_i} (Y_{j_0, \dots, j_n})| = \sum_{j \geq 0} \sum_{i=0}^{k_n(j)} |\tau^j(X_{i,j})| < \infty$ and $\sum_{\{j_n, \dots, j_0\} : \sum_0^n j_i > N} |\tau^{\sum_0^n j_i} (Y_{j_0, \dots, j_n})| \xrightarrow{N \rightarrow \infty} 0$ a.s.

Hence for all $(M_0, \dots, M_n) \in \mathbb{Z}_+^{n+1}$ the random variables $U_{M_0, \dots, M_n} := \prod_{j_n=0}^{M_n} \dots \prod_{j_0=0}^{M_0} \tau^{\sum_0^n j_i} (Y_{j_0, \dots, j_n})$ are distributed according to $\lambda_{M_0, \dots, M_n} := \star_{j_n=0}^{M_n} \dots \star_{j_0=0}^{M_0} \tau^{\sum_0^n j_i} (\nu)$.

Furthermore, we have $\sum_{(j_0, \dots, j_n) \notin \otimes_1^n \{M_0, \dots, M_i\}} |\tau^{\sum_0^n j_i} (Y_{j_0, \dots, j_n})| \rightarrow 0$ a.s. if $(M_0, \dots, M_n) \rightarrow \infty$ (since $\sum_0^n j_i > \min\{M_0, \dots, M_n\}$ if $(j_0, \dots, j_n) \notin \otimes_1^n \{M_0, \dots, M_i\}$). Thus U_{M_0, \dots, M_n} is a.s. convergent, to some random variable U say, for any sequence $(M_0, \dots, M_n) \rightarrow \infty$ (co-ordinate-wise). Hence $\lambda_{M_0, \dots, M_n} \rightarrow \mu'$. The Cauchy property yields that μ' , the distribution of U , does not depend on the particular sequence $(M_0, \dots, M_n) \rightarrow \infty$. Therefore, $\mu' = \star_{j_0} \dots \star_{j_n} \tau^{\sum_0^n j_i} (\nu)$ follows, i.e. (5.2) holds. But then by the step (5.1) \Rightarrow (5.2), $\mu' = \mu$ follows.

The proof is complete. \square

Putting things together we obtain the following result which contains Theorem 2.1.

PROPOSITION 5.6

Let \mathbb{G} be contractible with contracting $\tau \in \text{Aut}(\mathbb{G})$. Let $\mathcal{S} \subseteq \mathcal{M}^1(\mathbb{G})$ be a commutative sub-semigroup with $\tau\mathcal{S} \subseteq \mathcal{S}$ and assume in the sequel that ν and all cofactors belong to \mathcal{S} .

Consider the following conditions:

- (a) μ is n -times τ -decomposable, $\mu = v_{(n)} \star \tau(\mu), \dots, v_{(1)} = v \star \tau(v_{(1)})$.
 (b1) $v \in \mathcal{P}_{\log_+^{n+1}}$.
 (b2) $\mu = \star_{k \geq 0} \tau^k \left(v \binom{n+k}{n} \right)$ is convergent.
 (b3) $v_{(k)} = \star_{j \geq 0} \tau^j \left(v \binom{k+j}{k} \right)$ is convergent $\forall k \leq n+1$, $v_{(n+1)} = \mu$.
 (c) For an i.i.d. sequence $X_{i,j}$ with distribution $X_{i,j}(P) = v$, the products $\prod_{j \geq 0} \prod_{i=1}^{\binom{n+j}{n}} \tau^j X_{i,j}$ converge a.s. absolutely, i.e., $\sum_{j \geq 0} \sum_{i=1}^{\binom{n+j}{n}} |\tau^j X_{i,j}| < \infty$ a.s.

Then it follows that (b1) \Rightarrow (b2), (b3), and (c) (with cofactors $v_{(k)}$ as in (b3)), moreover, (b1) \Rightarrow (a).

And if, in addition, $H_r(\mu) = \{e\}$, then (a) \Rightarrow (b2) and (b2) \Rightarrow (b1).

The condition $H_r(\mu) = \{e\}$ can be replaced by any other condition of Proposition 4.9 providing a.s. convergence.

Proof. For (b1) \Rightarrow (b2), (b3), (c), see Proposition 4.7 and Remarks 4.3 and 4.10. For (c) \Rightarrow (b1), see Remark 4.10(b).

(b1) \Rightarrow (a) will be proved by induction on n :

$n = 1$: Let $v \in \mathcal{P}_{\log_+^2}$. Then the products $\mu = \star_{j \geq 0} \tau^j(v^{j+1})$ and $v_{(1)} = \star_{j \geq 0} \tau^j(v)$ are convergent (since (b1) and hence (b2) and (c) hold). Therefore, by Lemma 5.5, ($n = 1$), $\mu = \star_{j \geq 0} \tau^j \left(\star_{i \geq 0} \tau^i(v) \right)$, i.e. (a) follows by Proposition 5.2.

Let (b1) \Rightarrow (a) be proved for $k \leq n$. Let $v \in \mathcal{P}_{\log_+^{n+2}}$, hence (b2), (b3), (c) (by Proposition 4.7). (b2) yields $\mu = \star_{k \geq 0} \tau^k \left(v \binom{n+1+k}{n+1} \right)$ and $v_{(n)} = \star_{i \geq 0} \tau^i \left(v \binom{n+i}{n} \right)$ converge. Using $\sum_{i=0}^k \binom{n+i}{n} = \binom{n+1+k}{n+1}$, hence $v \binom{n+1+k}{n+1} = \star_{i=0}^k v \binom{n+i}{n}$, we obtain (observing that according to (c) and Remark 4.3, re-arrangement of convolution products is allowed):

$$\begin{aligned} \mu &= \star_{k \geq 0} \tau^k \left(v \binom{n+1+k}{n+1} \right) = \star_{k \geq 0} \star_{i=0}^k \tau^k \left(v \binom{n+i}{n} \right) \\ &= \star_{j \geq 0} \tau^j \left(\star_{i \geq j} \tau^i \left(v \binom{n+i}{n} \right) \right) = \star_{j \geq 0} \tau^j(v_{(n)}). \end{aligned}$$

By Proposition 5.2, μ is τ -decomposable with – per induction hypothesis n -times decomposable – cofactor $v_{(n)}$. Hence (a) holds.

If $H_r(\mu) = \{e\}$, the converse (a) \Rightarrow (b2) follows by Lemma 5.5, and (b2) \Rightarrow (b1) follows by Propositions 4.8 and 4.9. \square

6. Multiple τ -decomposability of continuous convolution semigroups and of generating functionals

It is well known that τ -decomposable laws are closely related to limit laws of triangular arrays of random variables of the form $\tau^k(X_j)$. However, as already mentioned in the pioneer work [14], even on the real line these arrays are in general not infinitesimal, hence the limits need not be infinitely divisible. Infinitely divisible τ -decomposable laws – called τ -semi-self-decomposable – on vector spaces have a considerable rich structure,

see for e.g. the investigations [15–18], and the literature mentioned there. Here we discuss briefly the possibility to carry over this concept to the group case.

Throughout we assume again \mathbb{G} to be contractible and $\tau \in \text{Aut}(\mathbb{G})$ to be contracting. If μ is τ -decomposable and if we assume in addition embeddability of μ into a continuous convolution semigroup, more generally, into a continuous hemigroup, then there exist infinitesimal triangular arrays converging to μ . We note a slight generalization of §3, Proposition 2 of [23].

PROPOSITION 6.1

Let τ be contracting and μ be τ -decomposable, $\mu = \nu \star \tau(\mu)$. Assume furthermore that ν is embeddable into a continuous hemigroup $(\nu(s, t))_{s \leq t}$ of probabilities with $\nu = \nu(0, 1)$. Then there exists an uniformly infinitesimal triangular array of the form $\lambda_{i,n} = \tau^n \left(\gamma_i^{(n)} \right)$, $1 \leq i \leq k_n$, such that

$$\star_{i=1}^{k_n} \lambda_{i,n} = \tau^n \left(\star_{i=1}^{k_n} \gamma_i^{(n)} \right) \xrightarrow{n \rightarrow \infty} \mu.$$

Here $\gamma_i^{(n)}$ are probabilities, embeddable into hemigroups for all n, i .

Proof. Choose $k_n \nearrow \infty$ such that $k_{n+1} - k_n \nearrow \infty$ with $k_0 = 0$, e.g. $k_n := n^2$. Put $\alpha_i := \nu \left(\frac{i-1-k_{j-1}}{k_j-k_{j-1}}, \frac{i-k_{j-1}}{k_j-k_{j-1}} \right)$ for $k_{j-1} < i \leq k_j$, $j \geq 0$, and $\gamma_i^{(n)} := \tau^{j-n-1}(\alpha_i)$. μ is τ -decomposable, hence $\star_{j=0}^{\infty} \tau^j(\nu) = \mu$. Therefore,

$$\begin{aligned} \star_{i=1}^{k_n} \lambda_{i,n} &= \tau^n \left(\star_{j=1}^{k_n} \gamma_j^{(n)} \right) = \tau^n \left(\star_{j=1}^n \left(\star_{i=k_{j-1}+1}^{k_j} \tau^{j-1-n}(\alpha_i) \right) \right) \\ &= \tau^n \left(\star_{j=1}^n \tau^{j-n-1} \left(\star_{i=k_{j-1}+1}^{k_j} \alpha_i \right) \right) = \left(\star_{j=1}^n \tau^{j-1} \left(\star_{i=k_{j-1}+1}^{k_j} \alpha_i \right) \right). \end{aligned}$$

By the hemigroup property $\star_{i=k_{j-1}+1}^{k_j} \alpha_i = \star_{i=k_{j-1}+1}^{k_j} \nu \left(\frac{i-1-k_{j-1}}{k_j-k_{j-1}}, \frac{i-k_{j-1}}{k_j-k_{j-1}} \right) = \nu(0, 1) = \nu$, hence $\star_{i=1}^{k_n} \lambda_{i,n} = \tau^n \left(\star_1^{k_n} \gamma_j^{(n)} \right) = \star_{j=0}^{n-1} \tau^j(\nu) \rightarrow \mu$.

To prove infinitesimality, put $\Gamma := \{\nu(s, t) : 0 \leq s \leq t \leq 1\}$. Γ is uniformly tight, hence for any neighbourhood U of e and any $\varepsilon > 0$ we have $\tau^n(\gamma)(\mathbb{G} \setminus U) < \varepsilon$ for sufficiently large n , for $n \geq N_\varepsilon$ say, and all $\gamma \in \Gamma$.

(1) For $n \geq 2N_\varepsilon$ and $j \leq \lfloor \frac{n}{2} \rfloor$, hence $n - j \geq N_\varepsilon$, if $k_{j-1} \leq i < k_j$. We have

$$\tau^n \left(\gamma_i^{(n)} \right) (\mathbb{G} \setminus U) = \tau^{n-j} \left(\nu \left(\frac{i-1-k_{j-1}}{k_j-k_{j-1}}, \frac{i-k_{j-1}}{k_j-k_{j-1}} \right) \right) (\mathbb{G} \setminus U) < \varepsilon.$$

(2) By continuity of the hemigroup there exists a $\delta > 0$ such that for $0 \leq t - s < \delta$ we have $\nu(s, t)(\mathbb{G} \setminus U) < \varepsilon$. By assumption, $k_n - k_{n-1} \nearrow \infty$, hence for sufficiently large n , $n \geq L_\delta$ say, we have $k_j - k_{j-1} > 1/\delta$. For $n \geq 2L_\delta$ and $j \leq \lfloor \frac{n}{2} \rfloor$ we obtain $\nu \left(\frac{i-1-k_{j-1}}{k_j-k_{j-1}}, \frac{i-k_{j-1}}{k_j-k_{j-1}} \right) (\mathbb{G} \setminus V) < \varepsilon$ for any neighbourhood V of the unit such that $V \subseteq \bigcap_{k \geq 0} \tau^{-k}(U)$ (cf. [30], Lemma 3.1.3(d) of [7]).

Hence for all $n \geq 2L_\delta$, $j \geq \lceil \frac{n}{2} \rceil$, $k_{j-1} \leq i < k_j$ we have

$$\begin{aligned} \tau^n \left(\gamma_i^{(n)} \right) (\mathbb{G} \setminus U) &= \tau^{n-j} \left(v \left(\frac{i-1-k_{j-1}}{k_j-k_{j-1}}, \frac{i-k_{j-1}}{k_j-k_{j-1}} \right) \right) (\mathbb{G} \setminus U) \\ &\leq v \left(\frac{i-1-k_{j-1}}{k_j-k_{j-1}}, \frac{i-k_{j-1}}{k_j-k_{j-1}} \right) (\mathbb{G} \setminus V) \\ &< \varepsilon, \end{aligned}$$

whence the assertion follows by (1) and (2). \square

DEFINITION 6.2

$\mu \in \mathcal{M}^1(\mathbb{G})$ is τ -semi-self-decomposable if there exist $\nu_n \in \mathcal{M}^1(\mathbb{G})$, $n \in \mathbb{N}$, and $b_n \in \text{Aut}(\mathbb{G})$ such that for $k_n \rightarrow \infty$ we have

- (i) $b_n^{-1} b_{n+1} \rightarrow \tau$,
- (ii) $b_n^{-1} \left(\begin{smallmatrix} k_n \\ \star \\ j=1 \end{smallmatrix} v_j \right) \rightarrow \mu$,

\star denoting convolution in revised order, $\star_1^n v_i := \nu_n \star \dots \star \nu_1$, and

- (iii) the array $(\lambda_{n,i} := b_n^{-1}(v_j))_{1 \leq i \leq k_n}$ is uniformly infinitesimal.

Let $\tilde{\mathcal{L}}_s(\tau)$ denote the set of limits of row products $b_n^{-1} \left(\begin{smallmatrix} k_n \\ \star \\ i=0 \end{smallmatrix} v_i \right)$ of arrays satisfying (i) and (ii), and let $\mathcal{L}_s(\tau)$ denote the set of τ -semi-self-decomposable laws.

We obviously have

PROPOSITION 6.3

$\mu \in \tilde{\mathcal{L}}_s(\tau)$ iff μ is τ -decomposable.

Proof.

\Rightarrow . By assumption, $b_{n+1}^{-1} b_n \rightarrow \tau$ and $b_{n+1}^{-1} \left(\begin{smallmatrix} k_{n+1} \\ \star \\ j=1 \end{smallmatrix} v_j \right) \rightarrow \mu$. Hence

$$\begin{aligned} b_{n+1}^{-1} \left(\begin{smallmatrix} k_{n+1} \\ \star \\ j=1 \end{smallmatrix} v_j \right) &= b_{n+1}^{-1} \left(\begin{smallmatrix} k_{n+1} \\ \star \\ j=k_{n+1} \end{smallmatrix} v_j \right) \star (b_{n+1}^{-1} b_n) b_n^{-1} \left(\begin{smallmatrix} k_n \\ \star \\ 1 \end{smallmatrix} v_i \right) \\ &\rightarrow v \star \tau(\mu). \end{aligned}$$

(The terms on the left-hand side and the second terms on the right-hand side converge both to μ resp. $\tau(\mu)$ by assumption, hence the first terms on the right are relatively compact according to the shift compactness theorem, with accumulation point v .)

\Leftarrow . We have, for any n ,

$$\mu = \lim_n \star_{j=0}^n \tau^{n-j}(v) = \lim_n \star_{j=0}^n \tau^j(v).$$

The assertion follows with $b_n := \tau^{-n}$ and $v_j = \tau^{-j}(v)$, $0 \leq j \leq n$. \square

Obviously the array $(\tau^n(v_j))$ is in general not uniformly infinitesimal.

On vector spaces, limits of *uniformly infinitesimal* triangular arrays are infinitely divisible, hence embeddable and vice versa. On (non-Abelian) groups embeddability of limits of commutative infinitesimal triangular arrays is only known for particular classes of groups under restrictive conditions, see [2, 19–21, 25, 26, 28] and the references therein. In the following we assume again $\mathcal{S} \subseteq \mathcal{M}^1(\mathbb{G})$ to be a commutative sub-semigroup with $\tau(\mathcal{S}) \subseteq \mathcal{S}$. If we assume in addition the arrays $\lambda_{n,j} = (b_n^{-1}(v_j))_{1 \leq j \leq k_n, n \geq 1}$ to be infinitesimal, under suitable conditions and under the assumption that all measures belong to \mathcal{S} , we obtain that τ -semi-self-decomposable laws are embeddable into continuous convolution semigroups.

To motivate the next definitions we consider first the example of real vector spaces:

Example 6.4. Let $\mathbb{G} = \mathbb{R}^d$, $\tau \in \text{GL}(\mathbb{R}^d)$. Let μ be infinitely divisible, hence embeddable into a uniquely determined continuous convolution semigroup $(\mu_t)_{t \geq 0}$ with $\mu = \mu_1$.

- (1) If μ_t is τ -decomposable for all $t \geq 0$, $\mu_t = \nu(t) \star \tau(\mu_t)$, then it is easily seen that the cofactors $\{\nu(t) : t \geq 0\}$ form a continuous convolution semigroup. Hence we write $\nu(t) = \nu_t$. Note that the cofactors are uniquely determined. (In fact, this follows by uniqueness of embedding and by Lévy's uniqueness and continuity theorem for Fourier transforms on \mathbb{R}^d .)
- (2) Assume that $\mu = \mu_1$ is embeddable and τ -decomposable, and moreover, that the cofactor is embeddable, i.e. there exist continuous convolution semigroups $(\mu_t)_{t \geq 0}$, $(\nu_t)_{t \geq 0}$ such that $\mu_1 = \nu_1 \star \tau(\mu_1)$. Again, it follows by uniqueness of embedding that $\mu_t = \nu_t \star \tau(\mu_t)$ for all $t \geq 0$.
- (3) Furthermore, if $\mu = \nu \star \tau(\mu)$ and if the cofactor ν is embeddable into a continuous convolution semigroup (ν_t) then μ is embeddable. In fact, $(\mu_t = \lim_n \star_{k=0}^n \tau^k(\nu_t))_{t \geq 0}$ is a continuous convolution semigroups, since the set of infinitely divisible laws is closed, and $\mu_1 = \mu$. Hence by (2), we obtain: a τ -decomposable law μ is embeddable iff the cofactor is embeddable.
- (4) In case $n = 1$, i.e. $\mu_t = \nu_t \star \tau(\mu_t)$ and $\nu_t = \lambda_t \star \tau(\nu_t)$ (for all t), $\mu_t = \lambda_t \star \tau(\nu_t) \star \tau(\mu_t) \forall t \geq 0$.

Analogously we obtain for n -times decomposability (for all t): $\mu_t =: \nu_t^{(n+1)} = \nu_t^{(n)} \star \tau(\nu_t^{(n-1)}) \star \dots \star \tau(\nu_t^{(0)})$, and $\nu_t^{(i)} = \nu_t^{(i-1)} \star \tau(\nu_t^{(i)})$, $1 \leq i \leq n+1$.

- (5) If we consider in (4) Fourier transforms resp. their logarithms, the '*second characteristic functions*', we obtain: $\hat{\nu}_t^{(n+1)} = \hat{\mu}_t = e^{t \cdot L}$, and $\hat{\nu}_t^{(i)} = e^{t \cdot L^{(i)}}$, $i = 1, \dots, n+1$. Thus $L = L^{(0)} + \sum_{i=1}^{n+1} L^{(i)} \circ \tau^*$, where $L =: L^{(n+1)}$, $L^{(i)} = L^{(i-1)} + L^{(i)} \circ \tau^*$, $1 \leq i < n+1$ (τ^* denoting the adjoint of $\tau \in \text{GL}(\mathbb{R}^d)$.)

This is a motivation for the following definition of (multiple) decomposability of continuous convolution semigroups on groups \mathbb{G} . All continuous convolution semigroups are assumed to have trivial idempotents, $\mu_0 = \varepsilon_e$.

DEFINITION 6.5

Let \mathbb{G} be a contractible group with contracting $\tau \in \text{Aut}(\mathbb{G})$.

- (a) $(\mu_t)_{t \geq 0}$ is called τ -decomposable continuous convolution semigroup if there exists a continuous convolution semigroup of 'cofactors' $(\nu_t)_{t \geq 0}$ such that $\mu_t = \nu_t \star \tau(\mu_t)$ for all $t \geq 0$ (Note that $\mu_0 = \varepsilon_e \Rightarrow \nu_0 = \varepsilon_e$.)

- (b) $(\mu_t)_{t \geq 0}$ is n -times τ -decomposable if all μ_t are τ -decomposable and if there exists a continuous convolution semigroup of cofactors $(\mu_t^{(n)})_{t \geq 0}$, which is $(n - 1)$ -times τ -decomposable.
- (c) Recall that continuous convolution semigroups (μ_t) are identified with their generating functionals $\mathcal{A} = \frac{d^+}{dt} |_{t=0} \mu_t$ evaluated at $f \in \mathcal{D}(\mathbb{G})$, the Schwartz–Bruhat function space. Let $\mathcal{GF}(\mathbb{G})$ denote the cone of generating functionals.

In view of (b) we define:

$\mathcal{A} =: \mathcal{A}^{(0)} \in \mathcal{GF}(\mathbb{G})$ is τ -decomposable – in symbols, $\mathcal{A} \in \Lambda_0(\tau)$ – if $\mathcal{A} = \mathcal{B} + \tau(\mathcal{A})$, for some $\mathcal{B} \in \mathcal{GF}(\mathbb{G})$.

Put $\Lambda_{-1} := \mathcal{GF}(\mathbb{G})$ and define analogously for $n \in \mathbb{N}$: $\mathcal{A} =: \mathcal{A}^{(n+1)}$ is n -times τ -decomposable – in symbols, $\mathcal{A} \in \Lambda_n(\tau)$ – if \mathcal{A} is τ -decomposable, $\mathcal{A} = \mathcal{A}^{(n)} + \tau(\mathcal{A})$, where $\mathcal{A}^{(n)} \in \Lambda_{n-1}$. Hence there exist $\mathcal{A}^{(i)} \in \mathcal{GF}(\mathbb{G})$, $i = 1, \dots, n + 1$, such that $\mathcal{A}^{(i)} = \mathcal{A}^{(i-1)} + \tau(\mathcal{A}^{(i)})$, therefore $\mathcal{A} = \mathcal{A}^{(0)} + \sum_{i=1}^{n+1} \tau(\mathcal{A}^{(i)})$.

Note that (c) is in analogy to the vector space case considered in Example 6.4(4). In fact, generating functionals replace the second characteristic functions. (For more details on continuous convolution semigroups and generating functionals, see [3, 7, 10, 29].)

Here we used the following notations: For $\mathcal{A} \in \mathcal{GF}(\mathbb{G})$ and $\sigma \in \text{Aut}(\mathbb{G})$, $\sigma(\mathcal{A}) \in \mathcal{GF}(\mathbb{G})$ is defined by $\langle \sigma(\mathcal{A}), f \rangle := \langle \mathcal{A}, f \circ \sigma \rangle$ for test functions $f \in \mathcal{D}(\mathbb{G})$.

More generally, let \mathbb{G}_i be locally compact groups, let $\psi : \mathbb{G}_1 \rightarrow \mathbb{G}_2$ be a homeomorphism which respects the differentiable structure, i.e. $f \circ \psi \in \mathcal{D}(\mathbb{G}_1)$ for all $f \in \mathcal{D}(\mathbb{G}_2)$. Then, $\psi(\mathcal{A}) \in \mathcal{GF}(\mathbb{G}_2)$ for all $\mathcal{A} \in \mathcal{GF}(\mathbb{G}_1)$. (Where again $\psi(\mathcal{A})$ is defined by $\langle \psi(\mathcal{A}), f \rangle := \langle \mathcal{A}, f \circ \psi \rangle$ for test functions $f \in \mathcal{D}(\mathbb{G}_2)$.) In fact, $\psi(\mathcal{A}) \in \mathcal{GF}(\mathbb{G}_2)$ follows immediately by E. Siebert’s characterization of generating functionals (Satz 5 of [29], Theorem 4.5.8 of [10]).

Remark 6.6. As mentioned before, if (μ_t) is a continuous convolution semigroup with generating functional \mathcal{A} , then n -times decomposability of (μ_t) implies n -times decomposability of \mathcal{A} . The converse is true if the involved measures $\mu_t, \nu_s^{(i)}$ belong to a commutative sub-semigroup \mathcal{S} with $\tau(\mathcal{S}) \subseteq \mathcal{S}$.

The n -times decomposable continuous convolution semigroups and generating functionals (in Lévy Khinchin representation of the second characteristic functions) were investigated for real and p -adic vector spaces and characterized in various ways (cf. [16, 18]). We use these results to obtain characterizations for at least the main examples, *connected* resp. *totally disconnected* contractible groups. First we mention the obvious result.

Lemma 6.7. *We assume that as above a homeomorphism $\psi : \mathbb{G}_1 \rightarrow \mathbb{G}_2$ exists respecting the differentiable structures, and in addition there exist $\tau \in \text{Aut}(\mathbb{G}_1)$, $\rho \in \text{Aut}(\mathbb{G}_2)$ such that $\psi \circ \tau = \rho \circ \psi$. Then we have*

- (a) *If τ is contracting (on \mathbb{G}_1), then so is ρ (on \mathbb{G}_2).*
- (b) *If $\mathcal{A} \in \mathcal{GF}(\mathbb{G}_1)$ is (n -times) τ -decomposable then also $\psi(\mathcal{A}) \in \mathcal{GF}(\mathbb{G}_2)$ is (n -times) ρ -decomposable, $n \geq 0$.*

The above mentioned results imply Proposition 6.8 below, from which Theorem 2.2 follows. We use the notations introduced in Proposition 3.11. Let \mathbb{G} be a contractible group, $\mathcal{A} \in \mathcal{GF}(\mathbb{G})$ a generating functional.

PROPOSITION 6.8

(a) Let \mathbb{G} be a connected contractible locally compact group, i.e., a (homogeneous) Lie group, of dimension d , say. Let $\mathcal{G} \cong \mathbb{R}^d$ denote the Lie algebra, \exp the exponential mapping and for $\tau \in \text{Aut}(\mathbb{G})$ let $\overset{\circ}{\tau} \in \text{GL}(\mathcal{G}) \subseteq \text{GL}(\mathbb{R}^d)$ denote the differential. For $A \in \mathcal{GF}(\mathbb{G})$ let $\overset{\circ}{A} := \exp^{-1}(A) \in \mathcal{GF}(\mathcal{G})$ and $(\lambda_t) \subseteq \mathcal{M}^1(\mathcal{G})$ the corresponding continuous convolution semigroup. Then the following assertions are equivalent:

- (i) A is n -times τ -decomposable (in $\mathcal{GF}(\mathbb{G})$).
- (ii) $\overset{\circ}{A}$ is n -times $\overset{\circ}{\tau}$ -decomposable (in $\mathcal{GF}(\mathcal{G}) \cong \mathcal{GF}(\mathbb{R}^d)$).
- (iii) The continuous convolution semigroup $(\lambda_t) \subseteq \mathcal{M}^1(\mathcal{G})$ generated by $\overset{\circ}{A}$ is n -times $\overset{\circ}{\tau}$ -decomposable.
- (iv) The Lévy–Khinchin representation of $\overset{\circ}{A} \in \mathcal{GF}(\mathcal{G}) \cong \mathcal{GF}(\mathbb{R}^d)$ has the structure described in Theorem 3.1 of [16].

(b) Let \mathbb{G} be a totally disconnected contractible locally compact group, with contracting $\tau \in \text{Aut}(\mathbb{G})$ (cf. Examples 3.4, 3.8). Let $\mathcal{G} \cong \mathbb{Q}_p$ denote the tangent object described in Example 3.8, Proposition 3.11(b), with corresponding homeomorphism $\psi : \mathbb{G} \rightarrow \mathbb{Q}_p$, and let $\overset{\circ}{\tau} \in \text{Aut}(\mathbb{Q}_p)$ such that $\psi \circ \tau = \overset{\circ}{\tau} \circ \psi$. For $A \in \mathcal{GF}(\mathbb{G})$ let $\overset{\circ}{A} := \psi(A)$ and $(\lambda_t) \subseteq \mathcal{M}^1(\mathcal{G})$ the corresponding continuous convolution semigroup. Then the following assertions are equivalent:

- (i) A is n -times τ -decomposable (in $\mathcal{GF}(\mathbb{G})$).
- (ii) $\overset{\circ}{A}$ is n -times $\overset{\circ}{\tau}$ -decomposable (in $\mathcal{GF}(\mathcal{G})$).
- (iii) The continuous convolution semigroup $(\lambda_t) \subseteq \mathcal{M}^1(\mathcal{G})$ generated by $\overset{\circ}{A}$ is n -times $\overset{\circ}{\tau}$ -decomposable.
- (iv) The Lévy–Khinchin representation of $\overset{\circ}{A} \in \mathcal{GF}(\mathcal{G}) = \mathcal{GF}(\mathbb{Q}_p)$ has the structure described in Theorems 4.3, 5.2 of [18]. (Note that on totally disconnected groups generating functionals are defined by the Lévy measures, which are characterized in [18].)

Proof.

- (a) $\psi := \exp^{-1} : \mathbb{G} \rightarrow \mathcal{G} \cong \mathbb{R}^d$ satisfies the assumptions of the Lemma 6.7 since \mathbb{G} is a simply connected nilpotent Lie group. The equivalences of (i) \Leftrightarrow (ii) resp. (ii) \Leftrightarrow (iii) follow by Lemma 6.7(b) resp. Remark 6.6, whereas (iii) \Leftrightarrow (iv) is proved in Theorem 3.1 of [16].
- (b) By Example 3.8 and Lemma 6.7 the equivalences (i)–(iii) follow as in (a). The equivalence (iii) \Leftrightarrow (iv) follows from Theorems 4.3 and 5.2 of [18]. (In fact, in [18] the result is proved for vector spaces over the field of p -adic numbers, hence for prime powers p . But, at least for dimension 1, the proof also works for the additive group of the ring of p -adic numbers, p not necessarily a prime power.) \square

Appendix: Multiple decomposability on matrix-cone hypergroups

The investigations before are not limited to the case of groups. If we stay in the locally compact set-up, hypergroups are natural candidates. For the background the reader is

referred to the monograph of Bloom and Heyer [1]. Recently Voit [34] and Rösler [24] investigated new classes of hypergroup structures on matrix cones with 'group like' properties. In [5] some basic probabilistic aspects of these hypergroups were investigated. It turned out that most of the tools needed for the investigations above in §§ 4–6 are also available for these hypergroups: There exist norms and automorphisms, in particular, contractible automorphisms. Furthermore, shift compactness theorems (Proposition 1.12, Theorem 1.10 of [5]), and a version of the equivalence theorem (Theorem 2.20 of [5]) are proved. Moreover, the convolution is commutative and Hermitean, hence by Lévy's continuity theorem for hypergroups convergence of infinite convolution products, triangular arrays are easier to handle. The τ -decomposable laws (the case $n = 0$) and their co-factors are characterized (Corollary 2.21–Proposition 2.23 of [5]). Altogether, most of the results for multiple decomposable laws obtained before have a counterpart (at least) on matrix cone hypergroups. We do not go into details here. The investigations will be continued and published elsewhere.

References

- [1] Bloom W and Heyer H, *Harmonic Analysis of Probability Measures on Hypergroups* (1995) (Berlin-New York: Walter de Gruyter)
- [2] Guivarc'h Y and Shah R, Asymptotic properties of convolution operators and limit distributions of triangular arrays on locally compact groups, *Trans. Am. Math. Soc.* **357** (2005) 3683–3729
- [3] Hazod W, *Stetige Faltungshalbgruppen von Wahrscheinlichkeitsmaßen und erzeugende Distributionen*, Lecture Notes in Mathematics 595 (1977) (Springer)
- [4] Hazod W, Multiple self-decomposable laws on vector spaces and on groups: The existence of background driving processes, in: *Statistical Inference, Economic Analysis and Matrix Algebra* edited by B Schipp and W Krämer (2008) (Physica Verlag) pp. 295–307
- [5] Hazod W, Probability on matrix-cone hypergroups: Limit theorems and structural properties, *J. Appl. Anal.* **15** (2009) 205–245
- [6] Hazod W and Scheffler H-P, Strongly τ -decomposable and selfdecomposable laws on simply connected nilpotent groups, *Monatsh. Math.* **128** (1999) 269–282
- [7] Hazod W and Siebert E, *Stable Probability Measures on Euclidean Spaces and on Locally Compact Groups, Structural Properties and Limit Theorems*, Mathematics and its Applications (2001) (Kluwer A.P.) vol. 531
- [8] Hebisch W and Sikora A, A smooths subadditive homogeneous norm on a homogeneous group, *Studia Math.* **96** (1990) 231–236
- [9] Hewitt E and Ross K A, *Abstract Harmonic Analysis I* (1970) (Berlin-Heidelberg-New York: Springer)
- [10] Heyer H, *Probability Measures on Locally Compact Groups* (1977) (Berlin-Heidelberg-New York: Springer)
- [11] Jurek Z and Mason D, *Operator Limit Distributions in Probability Theory* (1993) (J. Wiley Inc.)
- [12] Krakowiak W, Zero-one-laws for A -decomposable measures on Banach spaces, *Bull. Pol. Acad. Sci. Math.* **33** (1985) 85–90
- [13] Kosfeld K, *Zerlegbarkeitseigenschaften von Verteilungen auf lokalkompakten Gruppen*. Dissertation, Technische Universität Dortmund (2009), Electronic version available: <http://hdl.handle.net/2003/27013>
- [14] Loève M, Nouvelles classes de lois limites, *Bull. Soc. Math. Fr.* **73** (1945) 107–206
- [15] Maejima M and Naito Y, Semi-self-decomposable distributions and a new class of limit laws, *Probab. Theory Related Fields* **112** (1998) 13–31

- [16] Maejima M, Sato K I and Watanabe T, Operator semi-self-decomposability, (C, Q) -decomposability and related nested classes, *Tokyo J. Math.* **22** (1999) 473–509
- [17] Maejima M, Sato K I and Watanabe T, Completely operator semi-self-decomposable distributions, *Tokyo J. Math.* **23** (2000) 235–253
- [18] Maejima M and Shah R, Operator semistable, operator semi-selfdecomposable probability measures and related nested classes on p -adic vector spaces, *Monatsh. Math.* **151** (2007) 293–318
- [19] Neuenschwander D, Limits of commutative triangular systems on simply connected step 2-nilpotent Lie groups, *J. Theoret. Probab.* **5** (1992) 217–222
- [20] Neuenschwander D, Triangular systems on discrete subgroups of simply connected nilpotent Lie groups. *Publ. Math. Debrecen* **47** (1995) 329–333
- [21] Neuenschwander D, Triangular systems of probability measures on simply connected nilpotent Lie groups and discrete subgroups of exponential Lie groups, *C.R. Acad. Sci. Paris* **333** (2001) 1029–1034
- [22] Parthasarathy K R, Probability Measures on Metric Spaces (1967) (Academic Press)
- [23] Raja C R E, Operator semi-self-decomposable probability measures and related nested subclasses of measures on vector spaces, *Monatsh. Math.* **142** (1992) 351–361
- [24] Rösler M, Bessel convolutions on matrix cones, *Compos. Math.* **143** (2007) 749–779
- [25] Shah R, Limits of commutative triangular systems on real and p -adic groups, *Math. Proc. Camb. Philos. Soc.* **120** (1996) 181–192
- [26] Shah R, The central limit theorem on locally compact groups, *Israel J. Math.* **110** (1999) 189–218
- [27] Shah R, Self-decomposable measures on simply connected nilpotent Lie groups, *J. Theoret. Probab.* **13** (2000) 65–83
- [28] Shah R, Limits of commutative triangular systems on locally compact groups, *Proc. Indian Math. Soc.* **111** (2001) 49–63
- [29] Siebert E, Über die Erzeugung von Faltungshalbgruppen auf beliebigen lokalkompakten Gruppen, *Math. Z.* **131** (1973) 313–333
- [30] Siebert E, Contractive automorphisms on locally compact groups, *Math. Z.* **191** (1986) 73–90
- [31] Siebert E, Strongly operator-decomposable probability measures on separable Banach spaces, *Math. Nachr.* **154** (1991) 315–326
- [32] Siebert E, Operator decomposability of Gaussian measures on separable Banach spaces, *J. Theoret. Probab.* **5** (1992) 333–347
- [33] Tkačenko M, Introduction to topological groups, *Topology Appl.* **86** (1998) 179–231
- [34] Voit M, Bessel convolutions on matrix cones: Algebraic properties and random walks, *J. Theoret. Probab.* **22** (2009) 741–771
- [35] Wang J S P, The Mautner peneomen on p -adic Lie groups, *Math. Z.* **185** (1984) 403–412