

Outer-2-independent domination in graphs

MARCIN KRZYWKOWSKI^{1,2,*}, DOOST ALI MOJDEH³
and MARYEM RAOOFI⁴

¹Department of Pure and Applied Mathematics, University of Johannesburg,
Johannesburg, South Africa

²Faculty of Electronics, Telecommunications and Informatics, Gdansk University
of Technology, Gdansk, Poland

³Department of Mathematics, University of Mazandaran, Babolsar, Iran

⁴Department of Mathematics, Fereydun Kenar Education, Mazandaran, Iran

*Corresponding Author.

E-mail: marcin.krzywkowski@gmail.com; damojdeh@ipm.ir

MS received 21 March 2014; revised 21 August 2014

Abstract. We initiate the study of outer-2-independent domination in graphs. An outer-2-independent dominating set of a graph G is a set D of vertices of G such that every vertex of $V(G) \setminus D$ has a neighbor in D and the maximum vertex degree of the subgraph induced by $V(G) \setminus D$ is at most one. The outer-2-independent domination number of a graph G is the minimum cardinality of an outer-2-independent dominating set of G . We show that if a graph has minimum degree at least two, then its outer-2-independent domination number equals the number of vertices minus the 2-independence number. Then we investigate the outer-2-independent domination in graphs with minimum degree one. We also prove the Vizing-type conjecture for outer-2-independent domination and disprove the Vizing-type conjecture for outer-connected domination.

Keywords. Outer-2-independent domination; domination; outer-connected domination; Vizing's conjecture; cartesian product of graphs.

Mathematics Subject Classification. 05C69.

1. Introduction

Let $G = (V, E)$ be a graph. We denote the number of vertices of G by n and the number of edges by m . Thus $|V(G)| = n$ and $|E(G)| = m$. By complement of G (denoted by \bar{G}), we mean a graph which has the same vertices as G , and the two vertices of \bar{G} are adjacent if and only if they are not adjacent in G . By the neighborhood of a vertex v of G , we mean the set $N_G(v) = \{u \in V(G) : uv \in E(G)\}$. The degree of a vertex v , denoted by $d_G(v)$, is the cardinality of its neighborhood. By a leaf we mean a vertex of degree one, while a support vertex is a vertex adjacent to a leaf. We denote the set of leaves of a graph G by $L(G)$. Let $\delta(G)$ ($\Delta(G)$, respectively) mean the minimum (maximum, respectively) degree among all vertices of G . We denote the complete graph on n vertices by K_n . We

denote the path (cycle, respectively) on n vertices by P_n (C_n , respectively). A wheel W_n , where $n \geq 4$, is a graph with n vertices, formed by connecting a vertex to all vertices of a cycle C_{n-1} . The distance between two vertices of a graph is the number of edges in a shortest path connecting them. The eccentricity of a vertex is the greatest distance between it and any other vertex. The diameter of a graph G , denoted by $\text{diam}(G)$, is the maximum eccentricity among all vertices of G . By $K_{p,q}$, we denote a complete bipartite graph, the partite sets of which have cardinalities p and q . Generally, let K_{t_1, t_2, \dots, t_k} denote the complete multipartite graph with vertex set $S_1 \cup S_2 \cup \dots \cup S_k$, where $|S_i| = t_i$ for positive integers $i \leq k$. By a star we mean the graph $K_{1,m}$ where $m \geq 2$. We say that a subset of $V(G)$ is independent if there is no edge between any two vertices of this set. Generally, for positive integers k , a subset S of $V(G)$ is k -independent if the maximum degree of the subgraph induced by the vertices of S is at most $k - 1$. The k -independence number of a graph G , denoted by $\alpha^k(G)$, is the maximum cardinality of a k -independent subset of the set of vertices of G . The clique number of G , denoted by $\omega(G)$, is the number of vertices of a greatest complete graph which is a subgraph of G . By G^* we denote the graph obtained from G by removing all isolated vertices, leaves and support vertices. The cartesian product of graphs G and H , denoted by $G \square H$, is a graph such that $V(G \square H) = V(G) \times V(H)$ and $E(G \square H) = \{(u_1, v_1)(u_2, v_2) : u_1 = u_2 \text{ and } v_1 v_2 \in E(H) \text{ or } v_1 = v_2 \text{ and } u_1 u_2 \in E(G)\}$.

A subset $D \subseteq V(G)$ is a dominating set of G if every vertex of $V(G) \setminus D$ has a neighbor in D , while it is an outer-connected dominating set of G , if additionally, either $D = V(G)$ or $G - D$ is connected. The domination (outer-connected domination, respectively) number of a graph G , denoted by $\gamma(G)$ ($\tilde{\gamma}_c$, respectively), is the minimum cardinality of a dominating (outer-connected dominating, respectively) set of G . Outer-connected domination was introduced by Cyman [1], and further studied, for example, in [2, 4]. For a comprehensive survey of domination in graphs, see [3].

A subset $D \subseteq V(G)$ is an outer-2-independent dominating set, abbreviated O2IDS, of G if D is dominating and $\Delta(G - D) \leq 1$. The outer-2-independent domination number of G , denoted by $\tilde{\gamma}_{2i}(G)$, is the minimum cardinality of an outer-2-independent dominating set of G . An outer-2-independent dominating set of G of minimum cardinality is called a $\tilde{\gamma}_{2i}(G)$ -set.

We initiate the study of outer-2-independent domination in graphs. We show that if a graph has minimum degree at least two, then its outer-2-independent domination number equals the number of vertices minus the 2-independence number. Then we investigate the outer-2-independent domination in graphs with minimum degree one. We find the outer-2-independent domination numbers for several classes of graphs. Next we study the influence of removing or adding vertices and edges. Then we prove some lower and upper bounds, and we characterize the extremal graphs. We also give Nordhaus–Gaddum type inequalities. We prove the Vizing-type conjecture for outer-2-independent domination. We also disprove the Vizing-type conjecture for outer-connected domination.

2. General graphs

We begin with the following straightforward observation.

Observation 1. For every graph G , $\tilde{\gamma}_{2i}(G) \geq \gamma(G)$.

Let us observe that for any non-negative integer there exists a graph such that the difference between its outer-2-independent domination and domination numbers equals that non-negative integer.

Observation 2. For every integer $n \geq 3$, $\tilde{\gamma}_{2i}(K_n) = n - 2 = \gamma(K_n) + n - 3$.

We now give the outer-2-independent domination numbers of paths and cycles.

Observation 3. For every integer $n \geq 3$, $\tilde{\gamma}_{2i}(P_n) = \tilde{\gamma}_{2i}(C_n) = \lfloor (n + 2)/3 \rfloor$.

We now give the outer-2-independent domination numbers of complete multipartite graphs.

PROPOSITION 4

Let $n_1 \leq n_2 \leq \dots \leq n_t$ (where $t \geq 3$) be positive integers. We have

$$\tilde{\gamma}_{2i}(K_{n_1, n_2, \dots, n_t}) = \begin{cases} \sum_{j=1}^{t-1} n_j & \text{if } n_t \geq 2; \\ t - 2 & \text{if } n_t = 1. \end{cases}$$

Proof. Let V_1, V_2, \dots, V_t be the partite sets of the graph K_{n_1, n_2, \dots, n_t} , where $|V_i| = n_i$. If $n_t = 1$, then $n_i = 1$ for $1 \leq i \leq t$. Thus G is a complete graph K_t and $\tilde{\gamma}_{2i}(G) = t - 2$. Let $n_t \geq 2$. If one vertex from partite 1 to partite $t - 1$ is not in an O2IDS, then this vertex with the vertices of partite t induce a subgraph with $\Delta \geq 2$. Thus all vertices from partite 1 to partite $t - 1$ must be in any O2IDS. Now if we say S consists of the vertices of 1 to $t - 1$ partite sets, then S has minimum cardinality. Thus $\tilde{\gamma}_{2i}(K_{n_1, n_2, \dots, n_t}) = \sum_{j=1}^{t-1} n_j$. \square

We have the following lower bound on the outer-2-independent domination number of a graph in terms of its clique number.

Observation 5. For every graph G , $\tilde{\gamma}_{2i}(G) \geq \omega(G) - 2$.

Let us observe that the above bound is tight. For $n \geq 3$ we have $\tilde{\gamma}_{2i}(K_n) = n - 2 = \omega(K_n) - 2$.

Now, let us observe that for any non-negative integer there exists a graph such that the difference between its outer-2-independent domination number and clique number equals the non-negative integer.

Observation 6. For every positive integer k , $\tilde{\gamma}_{2i}(P_{3k}) = \omega(P_{3k}) + k - 2$.

We now prove that if a graph has no leaves and isolated vertices, then its outer-2-independent domination number equals the number of vertices minus the 2-independence number.

PROPOSITION 7

Let G be a graph. If $\delta(G) \geq 2$, then $\tilde{\gamma}_{2i}(G) = n - \alpha^2(G)$.

Proof. Let D be a maximum 2-independent set of G . Every vertex of D has at most one neighbor in the set D . Since $\delta(G) \geq 2$, every vertex of D has a neighbor in $V(G) \setminus D$. Let us observe that $V(G) \setminus D$ is a dominating set of the graph G . Consequently, it is an outer-2-independent dominating set of G . Therefore $\tilde{\gamma}_{2i}(G) \leq n - \alpha^2(G)$. On the other hand, we have $\alpha^2(G) \geq n - \tilde{\gamma}_{2i}(G)$ as the complement of every outer-2-independent dominating set is a 2-independent set. \square

3. Graphs with minimum degree one

Throughout this section we only consider connected graphs with minimum degree one.

We have the following relation between the outer-2-independent domination number of a graph and the 2-independence number of the graph obtained from it by removing all leaves and support vertices.

Lemma 8. For every graph $G \neq K_2$ with l leaves, $\tilde{\gamma}_{2i}(G) = n - \alpha^2(G^*) - l$.

Proof. Let us observe that there exists a $\tilde{\gamma}_{2i}(G)$ -set that does not contain any leaf. Let D be such a set. We have $V(G) \setminus (D \cup L(G)) \subseteq V(G^*)$. The set $V(G) \setminus (D \cup L(G))$ is 2-independent, thus $\alpha^2(G^*) \geq |V(G) \setminus (D \cup L(G))| = n - \tilde{\gamma}_{2i}(G) - l$. Now let D^* be a maximum 2-independent set of the graph G^* . Let us observe that in the graph G every vertex of D^* has a neighbor in the set $V(G) \setminus (D^* \cup L(G))$. This implies that $|V(G) \setminus (D^* \cup L(G))|$ is an outer-2-independent dominating set of the graph G . Therefore $\tilde{\gamma}_{2i}(G) \leq |V(G) \setminus (D^* \cup L(G))| = n - \alpha^2(G^*) - l$. This implies that $\tilde{\gamma}_{2i}(G) = n - \alpha^2(G^*) - l$. \square

We have the following bounds on the outer-2-independent domination number of a graph.

Observation 9. For every graph G , $1 \leq \tilde{\gamma}_{2i}(G) \leq n$.

We now characterize the graphs attaining the bounds from the previous observation.

PROPOSITION 10

Let G be a graph. Then

- (i) $\tilde{\gamma}_{2i}(G) = 1$ if and only if G has a universal vertex v such that each component of $G - v$ is K_1 or K_2 ;
- (ii) $\tilde{\gamma}_{2i}(G) = n$ if and only if $G = \overline{K_n}$.

Proof. First assume that G has a universal vertex, say v , such that $G - v$ consists of only isolated vertices and paths on two vertices. Let us observe that $\{v\}$ is an outer-2-independent dominating set of the graph G , as the vertex v dominates the whole graph, and the set $V(G) \setminus \{v\}$ is 2-independent. Thus $\tilde{\gamma}_{2i}(G) = 1$.

Now assume that G is a graph such that $\tilde{\gamma}_{2i}(G) = 1$. Let v be a vertex that forms an outer-2-independent dominating set of G . The vertex v is universal, as all vertices of G are dominated by it. The set $V(G) \setminus \{v\}$ is 2-independent, thus $G - v$ consists of isolated vertices and paths on two vertices.

Obviously, $\tilde{\gamma}_{2i}(\overline{K_n}) = n$. Now let G be any graph such that $\tilde{\gamma}_{2i}(G) = n$. Suppose that some two vertices of G , say x and y , are adjacent. Let us observe that $V(G) \setminus \{x\}$ is an O2IDS of the graph G . Therefore $\tilde{\gamma}_{2i}(G) \leq n - 1$, a contradiction. This implies that G consists of isolated vertices. \square

COROLLARY 11

For every graph G with at least two vertices, $\tilde{\gamma}_{2i}(G) \leq n - 1$.

3.1 Bounds

We have the following upper bound on the outer-2-independent domination number of a graph.

PROPOSITION 12

For every graph G with s support vertices, we have

$$\tilde{\gamma}_{2i}(G) \leq \frac{\Delta(G) \cdot (n - l) + s}{\Delta(G) + 1}.$$

Proof. By Lemma 8, we have $\tilde{\gamma}_{2i}(G) = n - \alpha^2(G^*) - l$. We now get

$$\alpha^2(G^*) \geq \alpha(G^*) \geq \gamma(G^*) \geq \frac{|V(G^*)|}{\Delta(G^*) + 1} \geq \frac{n - s - l}{\Delta(G) + 1}. \quad \square$$

We have the following upper bound on the outer-2-independent domination number of a graph in terms of its diameter.

PROPOSITION 13

If G is a graph of diameter D , then $\tilde{\gamma}_{2i}(G) \leq n - \lfloor 2(d + 1)/3 \rfloor$.

Proof. Let v_0, v_1, \dots, v_d be a diametrical path in G . If $d \in \{3k - 1, 3k\}$, then let $S = \{v_{3i}, v_{3i+2} : 0 \leq i \leq (d - 1)/3\}$. If $d = 3k + 1$, then let $S = \{v_{3i}, v_{3i+2} : 0 \leq i \leq d/3 - 1\} \cup \{v_d\}$. Let us observe that $V(G) \setminus S$ is an O2IDS of the graph G . \square

The bound in the previous proposition is tight, as $\tilde{\gamma}_{2i}(P_n) = \lfloor (n + 2)/3 \rfloor = n - \lfloor 2n/3 \rfloor = n - \lfloor 2(d + 1)/3 \rfloor$.

We have the following bounds on the outer-2-independent domination number of a graph in terms of its order and size.

PROPOSITION 14

For every graph G ,

$$n - 2 - \sqrt{n(n - 3) - 2m + 5} \leq \tilde{\gamma}_{2i}(G) \leq n - 2 + \sqrt{n(n - 3) - 2m + 5}.$$

Proof. Let D be a $\tilde{\gamma}_{2i}(G)$ -set. Let t denote the number of edges between the vertices of D and the vertices of $V(G) \setminus D$. Obviously, $m \leq (n - |D| - 1)/2 + t + |E(G[D])|$. Since G has at least one leaf, we have $t \leq |D| \cdot (|V(G) \setminus D| - 1) + 1$. Obviously, $|E(G[D])| \leq |D| \cdot (|D| - 1)/2$. Now simple calculations imply the result. \square

We now study the influence of the removal of a vertex of a graph on its outer-2-independent domination number.

PROPOSITION 15

Let G be a graph. For every vertex v of G , $\tilde{\gamma}_{2i}(G) - 1 \leq \tilde{\gamma}_{2i}(G - v) \leq \tilde{\gamma}_{2i}(G) + d_G(v) - 1$.

Proof. Let D be a $\tilde{\gamma}_{2i}(G)$ -set. If $v \notin D$, then observe that D is an O2IDS of the graph $G - v$. Now assume that $v \in D$. Let us observe that $D \cup N_G(v) \setminus \{v\}$ is an O2IDS of the graph $G - v$. Therefore $\tilde{\gamma}_{2i}(G - v) \leq |D \cup N_G(v) \setminus \{v\}| \leq |D \setminus \{v\}| + |N_G(v)| = \tilde{\gamma}_{2i}(G) + d_G(v) - 1$.

Now let D' be any $\tilde{\gamma}_{2i}(G - v)$ -set. It is easy to see that $D' \cup \{v\}$ is an O2IDS of the graph G . Thus $\tilde{\gamma}_{2i}(G) \leq \tilde{\gamma}_{2i}(G - v) + 1$. \square

Let us observe that the bounds from the previous proposition are tight. For the lower bound, let $G = K_n$, where $n \geq 4$. We have $\tilde{\gamma}_{2i}(K_n) = n - 2 = n - 3 + 1 = \tilde{\gamma}_{2i}(K_{n-1}) + 1 = \tilde{\gamma}_{2i}(K_n - v) + 1$. For the upper bound, let G be a star $K_{1,m}$. We denote the central vertex by v . We have $G - v = mK_1$. We get $\tilde{\gamma}_{2i}(G - v) = \tilde{\gamma}_{2i}(mK_1) = m = 1 + m - 1 = \tilde{\gamma}_{2i}(G) + d_G(v) - 1$.

We now study the influence of the removal of an edge of a graph on its outer-2-independent domination number.

PROPOSITION 16

Let G be a graph. For every edge e of G ,

$$\tilde{\gamma}_{2i}(G - e) \in \{\tilde{\gamma}_{2i}(G) - 1, \tilde{\gamma}_{2i}(G), \tilde{\gamma}_{2i}(G) + 1\}.$$

Proof. Let D be a $\tilde{\gamma}_{2i}(G)$ -set, and let $e = xy$ be an edge of G . If $x, y \notin D$ or $x, y \in D$, then it is easy to observe that D is an O2IDS of the graph $G - e$. Now assume that exactly one of those vertices, say x , belongs to the set D . Let us observe that $D \cup \{y\}$ is an O2IDS of $G - e$. Thus $\tilde{\gamma}_{2i}(G - e) \leq \tilde{\gamma}_{2i}(G) + 1$. Now let D' be a $\tilde{\gamma}_{2i}(G - e)$ -set. If some of the vertices x and y belong to the set D' , then D' is an O2IDS of the graph G . If none of the vertices x and y belongs to the set D' , then it is easy to observe that $D' \cup \{x\}$ is an O2IDS of the graph G . Therefore $\tilde{\gamma}_{2i}(G) \leq \tilde{\gamma}_{2i}(G - e) + 1$. \square

Let us observe that all values from the previous proposition can be achieved. For the lowest value, let G be a graph obtained from K_4 by removing an edge. Let e be an edge of G such that $G - e \neq C_4$. We have $\tilde{\gamma}_{2i}(G - e) = 1 = \tilde{\gamma}_{2i}(G) - 1$. For the middle value, let e be an edge of K_3 . We have $\tilde{\gamma}_{2i}(G - e) = 1 = \tilde{\gamma}_{2i}(G)$. For the highest value, let e be the edge of K_2 . We have $\tilde{\gamma}_{2i}(G - e) = 2 = \tilde{\gamma}_{2i}(G) + 1$.

Similarly, we have the following proposition concerning the influence of adding an edge on the outer-2-independent domination number of a graph.

PROPOSITION 17

Let G be a graph. If $e \notin E(G)$, then

$$\tilde{\gamma}_{2i}(G + e) \in \{\tilde{\gamma}_{2i}(G) - 1, \tilde{\gamma}_{2i}(G), \tilde{\gamma}_{2i}(G) + 1\}.$$

3.2 Nordhaus–Gaddum type inequalities

We now give Nordhaus–Gaddum type inequalities for the sum of the outer-2-independent domination number of a graph and its complement.

Theorem 18. *For every graph G , $n - 2 \leq \tilde{\gamma}_{2i}(G) + \tilde{\gamma}_{2i}(\bar{G}) \leq 2n$, with equality in the upper bound if and only if $G = K_1$.*

Proof. Obviously, $\tilde{\gamma}_{2i}(G) \leq n$ and $\tilde{\gamma}_{2i}(\bar{G}) \leq n$. Thus $\tilde{\gamma}_{2i}(G) + \tilde{\gamma}_{2i}(\bar{G}) \leq 2n$. Now assume that $\tilde{\gamma}_{2i}(G) \leq n - 3$. Let D be a $\tilde{\gamma}_{2i}(G)$ -set. Let us observe that for any triple vertices of $V(G) \setminus D$, at least one of them is adjacent in the graph \bar{G} to both the two remaining ones. Let \bar{D} be a $\tilde{\gamma}_{2i}(\bar{G})$ -set. At most two vertices of every triple of vertices of $V(G) \setminus D$ do not belong to the set \bar{D} as its complement is 2-independent. This implies that at most two vertices of $V(G) \setminus D$ do not belong to the set \bar{D} . Therefore $|\bar{D}| \geq |V(G) \setminus D| - 2$. We now get $\tilde{\gamma}_{2i}(G) + \tilde{\gamma}_{2i}(\bar{G}) = |D| + |\bar{D}| \geq |D| + |V(G) \setminus D| - 2 = n - 2$.

Assume that $\tilde{\gamma}_{2i}(G) + \tilde{\gamma}_{2i}(\bar{G}) = 2n$. Then $\tilde{\gamma}_{2i}(G) = \tilde{\gamma}_{2i}(\bar{G}) = n$, and by Proposition 10, $G = \bar{G} = \bar{K}_n$. This is only possible if $n = 1$, and so $G = K_1$. \square

We also characterize graphs G such that $\tilde{\gamma}_{2i}(G) + \tilde{\gamma}_{2i}(\bar{G}) = 2n - 1$.

PROPOSITION 19

If G is a connected graph, then $\tilde{\gamma}_{2i}(G) + \tilde{\gamma}_{2i}(\bar{G}) = 2n - 1$ if and only if $G = K_2$.

Proof. We have $\tilde{\gamma}_{2i}(K_2) + \tilde{\gamma}_{2i}(\bar{K}_2) = 2 + 1 = 3 = 2n - 1$. Now assume that for some graph G we have $\tilde{\gamma}_{2i}(G) + \tilde{\gamma}_{2i}(\bar{G}) = 2n - 1$. This implies that $\tilde{\gamma}_{2i}(G) = n$ or $\tilde{\gamma}_{2i}(\bar{G}) = n$. Without loss of generality, we assume that $\tilde{\gamma}_{2i}(G) = n$. By Proposition 10, we have $G = \bar{K}_n$ and so \bar{G} is a complete graph. Observation 2 implies that $\bar{G} = K_2$. \square

4. Cartesian product of graphs

In this section we investigate the outer-2-independent domination and outer-connected domination for Cartesian product of graphs. We also solve Vizing-type conjectures for those two variants of domination.

4.1 Outer-2-independent domination for Cartesian product of graphs

First we consider the Cartesian product of two cycles.

Theorem 20. *For any positive integers m and n , $\tilde{\gamma}_{2i}(C_m \square C_n) = \lceil mn/2 \rceil$.*

Proof. Proposition 7 implies that $\tilde{\gamma}_{2i}(C_m \square C_n) = mn - \alpha^2(C_m \square C_n)$. Let $V(C_m \square C_n) = \{v_{ij} : 1 \leq i \leq m \text{ and } 1 \leq j \leq n\}$, where v_{ij} is the vertex in row i and column j , and it is adjacent to vertices v_{i-1j} , v_{i+1j} , v_{ij-1} , v_{ij+1} (where $m + 1$ for rows means 1, and $n + 1$ for columns means 1). Let S be a maximum 2-independent set of $C_m \square C_n$. Let us observe that the graph $G[S]$ has at most one of edges $v_{ij}v_{ij+1}$, $v_{ij}v_{ij-1}$, $v_{ij}v_{i-1j}$, $v_{ij}v_{i+1j}$. Also, if two vertices v_{ij} and v_{ij+1} belong to the set S , then none of the vertices v_{i+1j} , v_{i+1j+1} , v_{i-1j} , v_{i-1j+1} , v_{ij-1} , v_{ij+2} belongs to S . First assume that m or n is even. Without loss of generality, we assume that $m = 2k$. For each column, at most k vertices can be in S , thus $|S| \leq n \cdot k = mn/2$. Let $S = \{v_{11}, v_{31}, \dots, v_{m-11}, v_{22}, v_{42}, \dots, v_{m2}, v_{13}, v_{33}, \dots, v_{m-13}, \dots, v_{2n}, v_{4n}, \dots, v_{mn}\}$. Let us observe that the set S is 2-independent. We have $|S| = mn/2$. This implies that $\alpha^2(S) = mn/2 = \lfloor mn/2 \rfloor$. Now assume that both m and n are odd. Let $m = 2k + 1$ and $n = 2l + 1$. Let us observe that for any two adjacent columns, at most $2k + 1$ vertices belong to the set S . Thus $|S| \leq \lfloor (2k + 1)(2l + 1)/2 \rfloor = \lfloor mn/2 \rfloor$. Let $S = \{v_{12}, v_{14}, \dots, v_{1n-1}, v_{21}, v_{23}, \dots, v_{2n}, v_{32}, v_{34}, \dots, v_{3n-1}, v_{m-11}, v_{m-13}, \dots, v_{m-1n}, v_{m2}, v_{m4}, \dots, v_{mn-1}\}$. Let us observe that the set S is 2-independent. We have $|S| = (mn - 1)/2 = \lfloor mn/2 \rfloor$. This implies that $\alpha^2(S) = \lfloor mn/2 \rfloor$. We now get $\tilde{\gamma}_{2i}(C_m \square C_n) = mn - \alpha^2(C_m \square C_n) = mn - \lfloor mn/2 \rfloor = \lceil mn/2 \rceil$. \square

Similarly we obtain the following results.

Theorem 21. For any positive integers m and n , $\tilde{\gamma}_{2i}(P_m \square C_n) = \lfloor mn/2 \rfloor$.

Theorem 22. For any positive integers m and n with m or n even we have $\tilde{\gamma}_{2i}(P_m \square P_n) = \lfloor mn/2 \rfloor$. Also, if $m, n \leq 3$, then $\tilde{\gamma}_{2i}(P_m \square P_n) = \lfloor mn/2 \rfloor$.

Theorem 23. Let $m \geq 3$ and $n \geq 3$ be positive odd integers. Then $\tilde{\gamma}_{2i}(P_m \square P_n) = \lfloor mn/2 \rfloor - \lfloor (n + 1)/6 \rfloor$.

Proof. Let $m = 2t + 1$. If $3k - 1 \leq n \leq 3k + 3$ for even positive integers k , then $\lfloor (n + 1)/6 \rfloor = k/2$. Thus without loss of generality we can assume that $n \in \{3k - 1, 3k + 1, 3k + 3\}$ for an even $k \geq 2$.

Let $n = 3k - 1$ and v_{ij} be the vertex in i -th row and j -th column of $P_m \square P_n$. The set $\{v_{ij} : i \text{ is odd, } 1 \leq i \leq 2t + 1 \text{ and } j \in \{1, 2, 4, 5, \dots, 3k - 2, 3k - 1\}\} \cup \{v_{ij} : i \text{ is even, } 2 \leq i \leq 2t \text{ and } j \in \{3, 6, \dots, 3k - 3\}\}$ is a maximum 2-independent set of $P_n \square P_m$. Therefore $\alpha^2(P_m \square P_n) = (t + 1) \cdot 2k + t(k - 1) = t(3k - 1) + 2k$ and $\tilde{\gamma}_{2i}(P_m \square P_n) = t(3k - 1) + k - 1$. It is easy to see that $t(3k - 1) + k - 1 = \lfloor (2t + 1)(3k - 1)/2 \rfloor - k/2$. Thus $\tilde{\gamma}_{2i}(P_m \square P_n) = \lfloor mn/2 \rfloor - \lfloor (n + 1)/6 \rfloor$.

Let $n = 3k + 1$. The set $\{v_{ij} : i \text{ is odd and } j \in \{1, 2, 4, 5, \dots, 3k - 2, 3k - 1, 3k + 1\}\} \cup \{v_{ij} : i \text{ is even and } j \in \{3, 6, \dots, 3k - 3, 3k\}\}$ is a maximum 2-independent set of $P_n \square P_m$. Similar calculation shows that $\tilde{\gamma}_{2i}(P_m \square P_n) = \lfloor mn/2 \rfloor - \lfloor (n + 1)/6 \rfloor$.

Let $n = 3k + 3$. The set $\{v_{ij} : i \text{ is odd and } j \in \{1, 2, 4, 5, \dots, 3k + 1, 3k + 2\}\} \cup \{v_{ij} : i \text{ is even and } j \in \{3, 6, \dots, 3k, 3k + 3\}\}$ is a maximum 2-independent set of $P_n \square P_m$. Also it is easily seen that $\tilde{\gamma}_{2i}(P_m \square P_n) = \lfloor mn/2 \rfloor - \lfloor (n + 1)/6 \rfloor$. \square

We now consider the Cartesian product of a path and a complete graph.

Theorem 24. Let m and $n \geq 4$ be positive integers. We have $\tilde{\gamma}_{2i}(P_m \square K_n) = m(n - 2)$.

Proof. Proposition 7 implies that $\tilde{\gamma}_{2i}(P_m \square K_n) = mn - \alpha^2(P_m \square K_n)$. Let $V(P_m \square K_n) = \{v_{ij} : 1 \leq i \leq m \text{ and } 1 \leq j \leq n\}$, where v_{ij} is the vertex in row i and column j . Let S be a maximum 2-independent set of the graph $P_m \square K_n$. Let us observe that S contains at most two vertices from every row. Thus $\alpha^2(P_m \square K_n) \leq 2m$. Now let S consist of two first vertices of every odd row and two last vertices of every even row. Let us observe that the set S is 2-independent. We have $|S| = 2m$. This implies that $\alpha^2(P_m \square K_n) = 2m$. We now get $\tilde{\gamma}_{2i}(P_m \square K_n) = mn - \alpha^2(P_m \square K_n) = mn - 2m = m(n - 2)$. \square

We now consider the Cartesian product of a cycle and a complete graph.

Theorem 25. *If m is an odd integer, then $\tilde{\gamma}_{2i}(C_m \square K_4) = 2m + 2$ and $\tilde{\gamma}_{2i}(C_m \square K_5) = 3m + 1$. Now let $m \geq 4$ and $n \geq 4$ be any integers. We have $\tilde{\gamma}_{2i}(C_m \square K_n) = m(n - 2)$ if m is even or $n \geq 6$.*

Proof. Proposition 7 implies that $\tilde{\gamma}_{2i}(C_m \square K_n) = mn - \alpha^2(C_m \square K_n)$. Let $V(C_m \square K_n) = \{v_{ij} : 1 \leq i \leq m \text{ and } 1 \leq j \leq n\}$, where v_{ij} is the vertex in row i and column j . Let S be a maximum 2-independent set of the graph $C_m \square K_n$. Let us observe that S has at most $2m$ vertices. Thus $\alpha^2(C_m \square K_n) \leq 2m$. If m is even, then let S consist of two first vertices of every odd column and two last vertices of every even column. Let us observe that the set S is 2-independent. Now assume that m is odd and n is at least six. Let S differ from the above defined S only in that from the last column it contains the third and the fourth vertex instead of the first two vertices. Let us observe that the set S is 2-independent. We have $|S| = 2m$. This implies that $\alpha^2(C_m \square K_n) = 2m$. We now get $\tilde{\gamma}_{2i}(C_m \square K_n) = mn - \alpha^2(C_m \square K_n) = mn - 2m = m(n - 2)$. \square

4.2 Vizing-type conjecture for outer-2-independent domination

We now prove that the Vizing-type conjecture for the outer-2-independent domination is true.

Theorem 26. *For any graphs G and H , $\tilde{\gamma}_{2i}(G \square H) \geq \tilde{\gamma}_{2i}(G) \cdot \tilde{\gamma}_{2i}(H)$.*

Proof. We consider $G \square H$ in a matrix form, where we have $|V(G)|$ rows H and $|V(H)|$ columns G . Proposition 7 implies that $\tilde{\gamma}_{2i}(G \square H) = |V(G)| \cdot |V(H)| - \alpha^2(G \square H)$. Let S be any minimum 2-independent set of the graph $G \square H$. Let us observe that the vertices of any column, which belong to the set S , form a 2-independent set of the graph G . Therefore $\alpha^2(G \square H) \leq |V(H)| \cdot \alpha^2(G)$. We now get

$$\begin{aligned} \tilde{\gamma}_{2i}(G \square H) &= |V(G)||V(H)| - \alpha^2(G \square H) \geq |V(G)||V(H)| - |V(H)|\alpha^2(G) \\ &\geq |V(G)||V(H)| - |V(H)|\alpha^2(G) + \alpha^2(H)(\alpha^2(G) - |V(G)|) \\ &= |V(G)||V(H)| - |V(H)| \cdot \alpha^2(G) + \alpha^2(H)\alpha^2(G) - \alpha^2(H)|V(G)| \\ &= (|V(G)| - \alpha^2(G))(|V(H)| - \alpha^2(H)) = \tilde{\gamma}_{2i}(G) \cdot \tilde{\gamma}_{2i}(H). \quad \square \end{aligned}$$

We have the following corollary from results of the previous subsection.

COROLLARY 27

For certain positive integers m and n , we have

- $\tilde{\gamma}_{2i}(C_m \square C_n) > \tilde{\gamma}_{2i}(C_m)\tilde{\gamma}_{2i}(C_n)$,
- $\tilde{\gamma}_{2i}(P_m \square P_n) > \tilde{\gamma}_{2i}(P_m)\tilde{\gamma}_{2i}(P_n)$,
- $\tilde{\gamma}_{2i}(P_m \square C_n) > \tilde{\gamma}_{2i}(P_m)\tilde{\gamma}_{2i}(C_n)$,
- $\tilde{\gamma}_{2i}(P_m \square K_n) > \tilde{\gamma}_{2i}(P_m)\tilde{\gamma}_{2i}(K_n)$,
- $\tilde{\gamma}_{2i}(C_m \square K_n) > \tilde{\gamma}_{2i}(C_m)\tilde{\gamma}_{2i}(K_n)$.

4.3 Vizing-type conjecture for outer-connected domination

We disprove the Vizing-type conjecture for outer-connected domination.

Observation 28 [1]. For integers $n \geq 3$, $\tilde{\gamma}_c(C_n) = n - 2$.

Counterexample. Let G be a graph $C_5 \square C_5$ with vertex set $\{v_{ij} : 1 \leq i, j \leq 5\}$. Let us observe that $\{v_{11}, v_{23}, v_{35}, v_{42}, v_{54}\}$ is a minimum outer-connected dominating set of G . Thus $\tilde{\gamma}_c(C_5 \square C_5) = 5$. By Observation 28, we have $\tilde{\gamma}_c(C_5) = 3$. We now get $\tilde{\gamma}_c(C_5 \square C_5) = 5 < 9 = \tilde{\gamma}_c(C_5) \cdot \tilde{\gamma}_c(C_5)$.

Acknowledgements

Thanks are due to the anonymous referee for comments that helped improve the paper. The research of the first author (MK) was supported by the Claude Leon Foundation, South Africa and by the Polish National Science Centre (Grant 2011/02/A/ST6/00201). The second author (DAM) was partially supported by a grant from IPM (No. 89050047).

References

- [1] Cyman J, The outer-connected domination number of a graph, *Australasian J. Combinatorics* **38** (2007) 35–46
- [2] Gravier S and Mollard M, On domination numbers of Cartesian product of paths, *Discrete Mathematics* **80** (1997) 247–250
- [3] Haynes T, Hedetniemi S and Slater P, Fundamentals of domination in graphs (1998) (New York: Marcel Dekker)
- [4] Mojdeh D, Firoozi P and Hasni R, On connected (γ, k) -critical graphs, *Australasian J. Combinatorics* **46** (2010) 25–35

COMMUNICATING EDITOR: B V Rajarama Bhat