

Bianchi Type-I bulk viscous fluid string dust magnetized cosmological model in general relativity

RAJ BALI and ANJALI

Department of Mathematics, University of Rajasthan, Jaipur 302 004, India

MS received 6 December 2003; accepted 8 April 2004

Abstract. Bianchi Type-I magnetized bulk viscous fluid string dust cosmological model is investigated. To get a determinate model, we have assumed the conditions $\sigma \propto \theta$ and $\zeta\theta = \text{constant}$ where σ is the shear, θ the expansion in the model and ζ the coefficient of bulk viscosity. The behaviour of the model in the presence and absence of magnetic field together with physical and geometrical aspects of the model are also discussed.

Keywords. Bianchi I; bulk viscous; magnetized string cosmological.

PACS Nos 04.20.-q; 04.20.Jb; 04.20.Cv

1. Introduction

The general relativistic treatment of strings was initiated by Letelier [1] and Stachel [2]. It is interesting to note that magnetic field present in galactic and inter galactic spaces plays a significant role at cosmological scale. Melvin [3] in the cosmological solution for dust and electromagnetic field suggested that during the evolution of the universe, the matter was in highly ionized state and is smoothly coupled with the field and consequently form a neutral matter as a result of universe expansion. Hence in string dust universe the presence of magnetic field is not unrealistic. Banerjee *et al* [4] investigated an axially symmetric Bianchi Type-I string dust cosmological model in the presence and absence of magnetic field using a supplementary condition $\alpha = a\beta$ where $\alpha = \alpha(t)$, $\beta = \beta(t)$ and a is a constant. The string cosmological models with magnetic field are also investigated by Chakraborty [5] and Tikekar and Patel [6]. Recently Bali and Upadhaya [7] investigated LRS Bianchi Type-I bulk viscous fluid string cosmological model in General Relativity. To get a determinate model, it has been assumed that $\sigma \propto \theta$ where σ is the shear and θ the expansion in the model. Bali and Upadhaya [8] have also investigated Bianchi Type-I magnetized string cosmological model in General Relativity.

In this paper, we have investigated Bianchi Type-I magnetized bulk viscous fluid string dust cosmological model in General Relativity. To get a determinate model, we have assumed the condition $\sigma \propto \theta$ and $\zeta\theta = \text{constant}$ where σ is the shear, θ the expansion in the model and ζ the coefficient of bulk viscosity. The behaviour of

the model in the presence and absence of magnetic field together with geometrical and physical aspects of the model are also discussed. We consider Bianchi Type-I metric in the form

$$ds^2 = -dt^2 + A^2 dx^2 + B^2 dy^2 + C^2 dz^2 \quad (1.1)$$

where A , B and C are functions of t alone.

The energy momentum T_i^j for string dust is given by

$$T_i^j = \varepsilon v_i v^j - \lambda x_i x^j - \zeta v_{;\ell}^{\ell} (g_i^j + v_i v^j) + E_i^j \quad (1.2)$$

together with

$$v^i v_i = -x^i x_i = -1 \quad (1.3)$$

and

$$v^i x_i = 0, \quad (1.4)$$

where $\varepsilon = \varepsilon_p + \lambda$ is the rest energy density for a cloud of strings. Here ε_p is the particle density, λ the string tension density, v^i the flow velocity vector, x^i the direction of strings and ζ the coefficient of bulk viscosity. Here E_{ij} is the electromagnetic field given by Lichnerowicz [9]

$$E_{ij} = \bar{\mu} \left[|h|^2 \left(v_i v_j + \frac{1}{2} g_{ij} \right) - h_i h_j \right], \quad (1.5)$$

where v_i is the flow vector satisfying

$$g_{ij} v^i v^j = -1, \quad (1.6)$$

$\bar{\mu}$ is the magnetic permeability and h_i the magnetic flux vector defined by

$$h_i = \frac{\sqrt{-g}}{2\bar{\mu}} \varepsilon_{ijk\ell} F^{k\ell} v^j, \quad (1.7)$$

where $F_{k\ell}$ is the electromagnetic field tensor and $\varepsilon_{ijk\ell}$ is the Levi Civita tensor density. We assume the coordinates to be comoving so that

$$v^1 = 0 = v^2 = v^3, \quad v^4 = 1. \quad (1.8)$$

The incident magnetic field is taken along x -axis so that

$$h_1 \neq 0, \quad h_2 = 0 = h_3 = h_4. \quad (1.9)$$

The first set of Maxwell's equation

$$F_{ij;k} + F_{jk;i} + F_{ki;j} = 0 \quad (1.10)$$

leads to

$$F_{23} = \text{constant} = H(\text{say}). \quad (1.11)$$

Bianchi Type-I cosmological model

Here $F_{14} = 0 = F_{24} = F_{34}$ due to assumption of infinite electrical conductivity. The only non-vanishing component of F_{ij} is F_{23} . Hence

$$h_1 = \frac{AH}{\bar{\mu}BC} \quad (1.12)$$

since

$$|h|^2 = h_\ell h^\ell = h_1 h^1 = g^{11}(h_1)^2 \quad (1.13)$$

and

$$|h|^2 = \frac{H^2}{\bar{\mu}^2 B^2 C^2}. \quad (1.14)$$

From eq. (1.5)

$$E_1^1 = -\frac{H^2}{2\bar{\mu}B^2C^2} = -E_2^2 = -E_3^3 = E_4^4. \quad (1.15)$$

Equation (1.2) leads to

$$T_1^1 = -\lambda - \frac{H^2}{2\bar{\mu}B^2C^2} - \zeta v_{;\ell}^\ell, \quad (1.16)$$

$$T_2^2 = \frac{H^2}{2\bar{\mu}B^2C^2} - \zeta v_{;\ell}^\ell, \quad (1.17)$$

$$T_3^3 = \frac{H^2}{2\bar{\mu}B^2C^2} - \zeta v_{;\ell}^\ell \quad (1.18)$$

$$T_4^4 = -\left(\varepsilon + \frac{H^2}{2\bar{\mu}B^2C^2}\right). \quad (1.19)$$

Using $(8\pi G/c^4) = 1$, the Einstein field equation

$$R_i^j - \frac{1}{2}Rg_i^j = -T_i^j \quad (1.20)$$

the line element (1.1) reduces to

$$\frac{B_{44}}{B} + \frac{C_{44}}{C} + \frac{B_4C_4}{BC} = \lambda + \frac{H^2}{2\bar{\mu}B^2C^2} + \zeta v_{;\ell}^\ell, \quad (1.21)$$

$$\frac{A_{44}}{A} + \frac{C_{44}}{C} + \frac{A_4C_4}{AC} = -\frac{H^2}{2\bar{\mu}B^2C^2} + \zeta v_{;\ell}^\ell, \quad (1.22)$$

$$\frac{A_{44}}{A} + \frac{B_{44}}{B} + \frac{A_4B_4}{AB} = -\frac{H^2}{2\bar{\mu}B^2C^2} + \zeta v_{;\ell}^\ell, \quad (1.23)$$

$$\frac{A_4 B_4}{AB} + \frac{B_4 C_4}{BC} + \frac{A_4 C_4}{AC} = \varepsilon + \frac{H^2}{2\bar{\mu} B^2 C^2}, \quad (1.24)$$

where the suffix 4 after A, B, C denotes ordinary differentiation with respect to t . Equations (1.21)–(1.24) are four equations in five unknowns A, B, C, λ and ε . For the complete determination of these equations, we assume two supplementary conditions: (i) The shear (σ) is proportional to the scalar of expansion θ which leads to

$$A = (BC)^n \quad (1.25)$$

and (ii)

$$\varepsilon = \lambda, \quad (1.26)$$

i.e. the rest energy density is equal to the string tension density.

From eqs (1.24) and (1.23), we have

$$\frac{A_{44}}{A} + \frac{B_{44}}{B} - \frac{B_4 C_4}{BC} - \frac{A_4 C_4}{AC} = -\frac{H^2}{\bar{\mu} B^2 C^2} - \varepsilon + \zeta v_{;\ell}^\ell. \quad (1.27)$$

Equations (1.27) and (1.21), together with condition (1.26) lead to

$$\frac{A_{44}}{A} + \frac{2B_{44}}{B} + \frac{C_{44}}{C} - \frac{A_4 C_4}{AC} = -\frac{H^2}{2\bar{\mu} B^2 C^2} + 2\zeta v_{;\ell}^\ell. \quad (1.28)$$

From eqs (1.25) and (1.28), we have

$$\begin{aligned} n(n-1)\frac{B_4^2}{B^2} + n(n-2)\frac{C_4^2}{C^2} + (2n^2-n)\frac{B_4 C_4}{BC} \\ + (n+2)\frac{B_{44}}{B} + (n+1)\frac{C_{44}}{C} \\ = -\frac{K}{B^2 C^2} + 2\zeta v_{;\ell}^\ell, \end{aligned} \quad (1.29)$$

where

$$K = \frac{H^2}{2\bar{\mu}}.$$

From eqs (1.22) and (1.23), we get

$$\frac{C_{44}}{C} - \frac{B_{44}}{B} = \frac{A_4}{A} \left(\frac{B_4}{B} - \frac{C_4}{C} \right). \quad (1.30)$$

Applying eq. (1.25) in (1.30), we get

$$\frac{(CB_4 - BC_4)_4}{(CB_4 - BC_4)} = -n \frac{(BC)_4}{BC}. \quad (1.31)$$

On integration, it leads to

Bianchi Type-I cosmological model

$$C^2 \left(\frac{B}{C} \right)_4 = \frac{L}{(BC)^n}, \quad (1.32)$$

where L is the integration constant.

Thus

$$B^2 = \mu\nu, \quad C^2 = \frac{\mu}{\nu}.$$

Equation (1.32) leads to

$$\frac{\nu_4}{\nu} = \frac{L}{\mu^{n+1}}. \quad (1.33)$$

Equation (1.29) leads to

$$\begin{aligned} & \left(\frac{2n+3}{2} \right) \frac{\mu_{44}}{\mu} + \left(\frac{4n^2-6n-3}{4} \right) \frac{\mu_4^2}{\mu^2} + \frac{1}{4} \frac{\nu_4^2}{\nu^2} + \frac{(n+1)}{2} \frac{\mu_4 \nu_4}{\mu\nu} + \frac{1}{2} \frac{\nu_{44}}{\nu} \\ & = -\frac{K}{\mu^2} + 2\zeta v_{;\ell}^\ell, \end{aligned} \quad (1.34)$$

where

$$BC = \mu, \quad \frac{B}{C} = \nu.$$

Equations (1.33) and (1.34) lead to

$$\left(\frac{2n+3}{2} \right) \frac{\mu_{44}}{\mu} + \left(\frac{4n^2-6n-3}{4} \right) \frac{\mu_4^2}{\mu^2} + \frac{3}{4} \frac{L^2}{\mu^{2n+2}} = -\frac{K}{\mu^2} + 2\zeta v_{;\ell}^\ell. \quad (1.35)$$

In special case, if $L = 0$, eq. (1.35) leads to

$$\left(\frac{2n+3}{2} \right) \frac{\mu_{44}}{\mu} + \left(\frac{4n^2-6n-3}{4} \right) \frac{\mu_4^2}{\mu^2} = -\frac{K}{\mu^2} + 2\zeta v_{;\ell}^\ell. \quad (1.36)$$

Applying the condition

$$\zeta\theta = \text{constant}, \quad (1.37)$$

eq. (1.36) leads to

$$\left(\frac{2n+3}{2} \right) \frac{\mu_{44}}{\mu} + \left(\frac{4n^2-6n-3}{4} \right) \frac{\mu_4^2}{\mu^2} - \beta = -\frac{K}{\mu^2} \quad (1.38)$$

which leads to

$$\mu\mu_{44} + \ell\mu_4^2 - \beta\mu^2 = -K, \quad (1.39)$$

where

Raj Bali and Anjali

$$\ell = \frac{4n^2 - 6n - 3}{2(2n + 3)}, \quad \beta = \frac{4\zeta v_{;\ell}^\ell}{(2n + 3)}$$

which leads to

$$2ff' + \frac{2\ell}{\mu}f^2 = 2\beta\mu - \frac{2K}{\mu} \quad (1.40)$$

where

$$\mu_4 = f(\mu) \quad \text{and} \quad f' = \frac{df}{d\mu}.$$

From eq. (1.40), we have

$$\frac{d}{d\mu}(f^2) + \frac{2\ell}{\mu}f^2 = 2\beta\mu - \frac{2K}{\mu}. \quad (1.41)$$

Equation (1.41) leads to

$$f^2 = \frac{\beta}{\ell + 1}\mu^2 - \frac{K}{\ell} + \frac{P}{\mu^{2\ell}}, \quad (1.42)$$

where P is the constant of integration.

To find the solution, we take $P = 0$. Equation (1.42) leads to

$$\mu = \sqrt{\frac{K\alpha}{\beta\ell}} \cosh\left(\sqrt{\frac{\beta}{\alpha}}t + N\right), \quad (1.43)$$

where $\ell + 1 = \alpha$ and N is the integration constant.

Since $L = 0$, from eq. (1.33)

$$\nu = b \text{ (constant)}. \quad (1.44)$$

Therefore metric (1.1) reduces to

$$\begin{aligned} ds^2 = & -dt^2 + \left(\frac{K\alpha}{\ell\beta}\right)^n \cosh^{2n}\left(\sqrt{\frac{\beta}{\alpha}}t + N\right) dx^2 \\ & + b\sqrt{\frac{K\alpha}{\ell\beta}} \cosh\left(\sqrt{\frac{\beta}{\alpha}}t + N\right) dy^2 \\ & + \frac{1}{b}\sqrt{\frac{K\alpha}{\ell\beta}} \cosh\left(\sqrt{\frac{\beta}{\alpha}}t + N\right) dz^2. \end{aligned} \quad (1.45)$$

On applying transformation

$$\begin{aligned} \alpha &= e^\beta - 1, \\ \ell &= a \sin K, \\ t &= \sin \frac{K\tau}{K}, \end{aligned}$$

Bianchi Type-I cosmological model

metric (1.45) reduces to

$$\begin{aligned}
 ds^2 = & -\cos^2 K\tau d\tau^2 + \left(\frac{K(e^\beta - 1)}{\beta a \sin K}\right)^n \\
 & \times \cosh^{2n} \left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right) dX^2 \\
 & + \left(\frac{K(e^\beta - 1)}{\beta a \sin K}\right)^{1/2} \cosh \left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right) dY^2 \\
 & + \left(\frac{K(e^\beta - 1)}{\beta a \sin K}\right)^{1/2} \cosh \left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right) dZ^2. \quad (1.46)
 \end{aligned}$$

In the absence of magnetic field, i.e. $K = 0$, metric (1.46) reduces to

$$\begin{aligned}
 ds^2 = & -d\tau^2 + \left(\frac{(e^\beta - 1)}{a\beta}\right)^n \cosh^{2n} \left(\sqrt{\frac{\beta}{e^\beta - 1}} \tau + N\right) dX^2 \\
 & + \left(\frac{(e^\beta - 1)}{a\beta}\right)^{1/2} \cosh \left(\sqrt{\frac{\beta}{e^\beta - 1}} \tau + N\right) dY^2 \\
 & + \left(\frac{(e^\beta - 1)}{a\beta}\right)^{1/2} \cosh \left(\sqrt{\frac{\beta}{e^\beta - 1}} \tau + N\right) dZ^2. \quad (1.47)
 \end{aligned}$$

In the absence of viscosity, i.e., $\beta = 0$, the metric (1.46) reduces to

$$\begin{aligned}
 ds^2 = & -\cos^2 K\tau d\tau^2 + \left(\frac{K}{a \sin K}\right)^n \cosh^{2n} \left(\frac{\sin K\tau}{K} + N\right) dX^2 \\
 & + \left(\frac{K}{a \sin K}\right)^{1/2} \cosh \left(\frac{\sin K\tau}{K} + N\right) dY^2 \\
 & + \left(\frac{K}{a \sin K}\right)^{1/2} \cosh \left(\frac{\sin K\tau}{K} + N\right) dZ^2. \quad (1.48)
 \end{aligned}$$

2. Some physical and geometrical features

The density (ε) for model (1.46) is given by

$$\begin{aligned}
 \lambda = \varepsilon = & \left(\frac{4n + 1}{4}\right) \sqrt{\frac{\beta}{e^\beta - 1}} \tanh \left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right) \\
 & - \frac{\beta a \sin K}{(e^\beta - 1)} \operatorname{sech}^2 \left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right). \quad (2.1)
 \end{aligned}$$

The scalar of expansion (θ) and the expression of shear tensor (σ_i^j) for the space-time (1.46) are given by

$$\theta = (n + 1)\sqrt{\frac{\beta}{e^\beta - 1}} \tanh\left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right), \quad (2.2)$$

$$\sigma_1^1 = \left(\frac{2n - 1}{3}\right) \sqrt{\frac{\beta}{e^\beta - 1}} \tanh\left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right), \quad (2.3)$$

$$\sigma_2^2 = \sigma_3^3 = \left(\frac{1 - 2n}{6}\right) \sqrt{\frac{\beta}{e^\beta - 1}} \tanh\left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right). \quad (2.4)$$

Thus

$$\sigma^2 = \frac{1}{2}(\sigma_{ij}\sigma^{ij}),$$

$$\sigma^2 = \frac{(2n - 1)^2}{12} \left(\frac{\beta}{e^\beta - 1}\right) \tanh^2\left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right). \quad (2.5)$$

Spatial volume

$$R^3 = \left(\frac{K(e^\beta - 1)}{\beta a \sin K}\right)^{(n+2)/2} \cosh^{n+1}\left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right). \quad (2.6)$$

3. Discussion

The expansion in the model (1.46) increases as $\left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right)$ increases, i.e. it represents expanding universe. From (2.1), we find that $\varepsilon > 0$. For $\tau = 0$ we find ε as a finite quantity. The spatial volume tends to infinity as $\left(\sqrt{\frac{\beta}{e^\beta - 1}} \frac{\sin K\tau}{K} + N\right)$ tends to infinity. Since $\lim_{\tau \rightarrow \infty} \frac{\sigma}{\theta} \neq 0$, the model does not approach isotropy for large values of τ . For $n = \frac{1}{2}$, model (1.46) represents an isotropic universe.

In the absence of magnetic field, i.e., when $K = 0$ then the energy density (ε), string tension (λ), the expansion θ , the components of shear tensor and the spatial volume (R^3) in the presence of bulk viscosity are given by

$$\varepsilon = \lambda = \frac{(4n + 1)}{4} \left(\frac{\beta}{e^\beta - 1}\right) \tanh\left(\sqrt{\frac{\beta}{e^\beta - 1}} \tau + N\right), \quad (3.1)$$

$$\varepsilon = (n + 1)\sqrt{\frac{\beta}{e^\beta - 1}} \tanh\left(\sqrt{\frac{\beta}{e^\beta - 1}} \tau + N\right), \quad (3.2)$$

$$\sigma_1^1 = \left(\frac{2n - 1}{3}\right) \sqrt{\frac{\beta}{e^\beta - 1}} \tanh\left(\sqrt{\frac{\beta}{e^\beta - 1}} \tau + N\right), \quad (3.3)$$

Bianchi Type-I cosmological model

$$\sigma_2^2 = \sigma_3^2 = \left(\frac{1-2n}{6}\right) \sqrt{\frac{\beta}{e^\beta - 1}} \tanh\left(\sqrt{\frac{\beta}{e^\beta - 1}}\tau + N\right). \quad (3.4)$$

Thus

$$\sigma^2 = \frac{(2n-1)^2}{12} \left(\frac{\beta}{e^\beta - 1}\right) \tanh^2\left(\sqrt{\frac{\beta}{e^\beta - 1}}\tau + N\right) \quad (3.5)$$

and

$$R^3 = \left(\frac{(e^\beta - 1)}{a\beta}\right)^{(2+n)/2} \cosh^{n+1}\left(\sqrt{\frac{\beta}{e^\beta - 1}}\tau + N\right). \quad (3.6)$$

In the absence of viscosity, i.e., $\beta = 0$ then the above quantities in the presence of magnetic field are given by

$$\varepsilon = \left(\frac{4n+1}{4}\right) \tanh\left(\frac{\sin K\tau}{K} + N\right) - a \sin K \operatorname{sech}^2\left(\frac{\sin K\tau}{K} + N\right), \quad (3.7)$$

$$\theta = (n+1) \tanh\left(\frac{\sin K\tau}{K} + N\right), \quad (3.8)$$

$$\sigma_1^2 = \left(\frac{2n-1}{3}\right) \tanh\left(\frac{\sin K\tau}{K} + N\right), \quad (3.9)$$

$$\sigma_2^2 = \sigma_3^2 = \frac{1}{3} \left(\frac{1-2n}{2}\right) \tanh\left(\frac{\sin K\tau}{K} + N\right). \quad (3.10)$$

Thus

$$\sigma^2 = \frac{(2n-1)^2}{12} \tanh^2\left(\frac{\sin K\tau}{K} + N\right) \quad (3.11)$$

and

$$R^3 = \left(\frac{K}{a \sin K}\right)^{(2+n)/2} \cosh^{n+1}\left(\frac{\sin K\tau}{K} + N\right) \quad (3.12)$$

when $K \rightarrow 0$ and $\beta \rightarrow 0$ then these quantities are given by

$$\varepsilon = \lambda = \left(\frac{4n+1}{4}\right) \tanh(\tau + N), \quad (3.13)$$

$$\theta = (n+1) \tanh(\tau + N), \quad (3.14)$$

$$\sigma_1^2 = \left(\frac{2n-1}{3}\right) \tanh(\tau + N), \quad (3.15)$$

Raj Bali and Anjali

$$\sigma_2^2 = \sigma_3^3 = \left(\frac{1-2n}{6} \right) \tanh(\tau + N), \quad (3.16)$$

$$\sigma^2 = \frac{(2n-1)^2}{12} \tanh^2(\tau + N) \quad (3.17)$$

and

$$R^3 = a^{2/2+n} \cosh^{n+1}(\tau + N). \quad (3.18)$$

In the absence of magnetic field and bulk viscosity, the model represents an expanding universe and $\varepsilon > 0$. The spatial volume increases as τ increases and it is infinity when $\tau = \infty$. Since $\lim_{\tau \rightarrow \infty} (\sigma/\theta) \neq 0$, the model does not approach isotropy for large values of τ .

References

- [1] P S Letelier, *Phys. Rev.* **D20**, 1294 (1979); *Phys. Rev.* **D28**, 10, 2414 (1983)
- [2] J Stachel, *Phys. Rev.* **D21**, 2171 (1980)
- [3] M A Melvin, *Ann. N.Y. Acad. Sci.* **262**, 253 (1975)
- [4] A Banerjee, A K Sanyal and S Chakraborty, *Pramana - J. Phys.* **42(6)**, 483 (1990)
- [5] S Chakraborty, *Indian J. Pure Appl. Phys.* **29**, 31 (1991)
- [6] R Tikekar and L K Patel, *Gen. Relativ. Gravit.* **24**, 397 (1992)
- [7] R Bali and R D Upadhaya, *Astrophys. and Space Science* **288**, 287 (2003)
- [8] R Bali and R D Upadhaya, *Ganita Sandesh - J. Raj. Ganit Parishad* **16**, 69 (2002)
- [9] A Lichnerowicz, *Relativistic hydrodynamics and magnetohydrodynamics* (Benjamin, New York, 1967) p. 13