

Scalar top study: Detector optimization

C MILSTÉNE¹ and A SOPCZAK^{2,*}

¹Fermi National Laboratory, Batavia, IL-60510, USA

²Department of Physics, Lancaster University, LA1 4YW, UK

E-mail: caroline@fnal.gov; andre.sopczak@cern.ch

Abstract. A vertex detector concept of the linear collider flavour identification (LCFI) collaboration, which studies pixel detectors for heavy quark flavour identification, has been implemented in simulations for *c*-quark tagging in scalar top studies. The production and decay of scalar top quarks (stops) is particularly interesting for the development of the vertex detector as only two *c*-quarks and missing energy (from undetected neutralinos) are produced for light stops. Previous studies investigated the vertex detector design in scenarios with large mass differences between stop and neutralino, corresponding to large visible energy in the detector. In this study we investigate the tagging performance dependence on the vertex detector design in a scenario with small visible energy for the international linear collider (ILC).

Keywords. Supersymmetry; scalar top quarks; CCD detector; vertex detector; linear collider flavour identification; detector concept; linear collider; beam delivery system; radiation damage; multiple scattering.

PACS Nos 12.20.Pb; 14.80.ty; 29.00.00; 14.65.DW; 13.66.JN

1. Introduction

The development of a vertex detector for a linear collider is an important and challenging enterprise. A key aspect is the distance of the innermost layer to the interaction point, which is related to radiation hardness and beam background. Another key aspect is the number of radiation lengths the particles go through, since it determines the multiple scattering which affects the vertex reconstruction.

The optimization of the vertex detector tagging performance is of great importance for studies of physics processes. While mostly at previous and current accelerators (e.g. SLC, LEP, Tevatron) *b*-quark tagging has revolutionized many searches and measurements, *c*-quark tagging will be very important at a future linear collider, for example, in studies of supersymmetric dark matter [1]. Therefore, *c*-quark tagging could be a benchmark for vertex detector developments. The scalar top

*Speaker

Table 1. Number of background events.

Thickness	Layers	12% Signal efficiency	25% Signal efficiency
Single	5 (4)	68 (82)	2300 (2681)
Double	5 (4)	69 (92)	2332 (2765)

production and decay process, and the implemented vertex detector geometry are shown in ref. [2].

The analysis for a large mass difference with the SPS-5 parameter point (ISAJET) $m_{\tilde{t}_1} = 220.7$ GeV, $m_{\tilde{\chi}_1^0} = 120$ GeV and $\cos\theta_{\tilde{t}} = 0.5377$ was previously performed [3]. For 25% (12%) efficiency 3800 (1800) signal events and 5400 (170) background events without c -quark tagging were obtained, while the background was reduced to 2300 (68) events with c -quark tagging.

The vertex detector radiation length was varied between single thickness (TESLA TDR) and double thickness. In addition, the number of vertex detector layers was varied between five layers (innermost layer at 1.5 cm as in the TESLA TDR) and four layers (innermost layer at 2.6 cm). For SPS-5 parameters, the following number of background events remain as shown in table 1 [3].

A significantly larger number of background events was expected if the first layer of the vertex detector is removed. The distance of the first layer to the interaction point is also an important aspect from the accelerator physics (beam delivery) perspective. The interplay between the beam delivery and vertex detector design in regard to critical tolerances like hardware damage of the first layer and occupancy (unable to use the data of the first layer) due to beam background goes beyond the scope of this study and will be addressed in the future.

For large visible energy (large mass difference) no significant increase in the expected background was observed for doubling the thickness of the vertex detector layers [3,4]. In this study the effect of the vertex detector design on the events with smaller visible energy in the detector is addressed.

2. Signal and background simulations

The production of simulated light stops at a 500 GeV linear collider is analyzed using high luminosity $\mathcal{L} = 500 \text{ fb}^{-1}$. The signature for stop pair production at an e^+e^- collider is two charm jets and large missing energy:

$$e^+e^- \rightarrow \tilde{t}_1 \bar{\tilde{t}}_1 \rightarrow c \tilde{\chi}_1^0 \bar{c} \tilde{\chi}_1^0. \tag{1}$$

For small $\Delta m = m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0}$, the jets are relatively soft and separation from backgrounds is very challenging. Backgrounds arising from various standard model processes can have cross-sections that are several orders of magnitude larger than the signal. Thus, it is necessary to study this process with a realistic detector simulation. Signal and background events are generated with PYTHIA 6.129 [5], including a scalar top signal generation [6] previously used in ref. [7]. The detector simulation is based on the fast simulation SIMDET [8], describing a typical ILC

detector. Good agreement in comparisons with SGV [9] detector simulations was obtained [3,4].

Cross-sections for the signal process and the relevant backgrounds have been computed with a code used in ref. [10] and by GRACE 2.0 [11], with cross-checks to COMHEP 4.4 [12]. A minimal transverse momentum cut, $p_t > 5$ GeV, is applied for the two-photon background, to avoid the infra-red divergence. Details of the event selection are given in ref. [1].

The c -tagging with the LCFI detector is based on vertex identification and a neural network application [13]. The vertex identification considered three cases for each jet independently:

- (a) *Only a primary vertex.* In this case, the two tracks with the largest separation in the r - ϕ plane are considered and for these tracks the neural network variables include the impact parameter and its significance (impact parameter divided by uncertainty) both in the r - ϕ plane and in the z -direction, their momenta, and the joined probability in r - ϕ plane and z -direction.
- (b) *One secondary vertex.* In addition to the previous variables, the decay length significance of the secondary vertex, the multiplicity and momenta of all associated tracks, and the P_t corrected mass of the secondary vertex (corrected for neutral hadrons and neutrinos), the P_t of the decay products perpendicular to the flight direction between primary and secondary vertex, and the joint probability in r - ϕ and z -direction.
- (c) *More than one secondary vertex.* Two secondary vertices are considered, where the tracks are assigned to the vertex closest to the primary vertex, and the neural network input variables are defined as in case (b).

The neural network is tuned to 255,000 simulated signals and 240,000 $W\epsilon\nu$ background events. The signal events are a combination of all simulated signal events for the scalar top mass range between 120 and 220 GeV and for $\Delta m = 5, 10$ and 15 GeV.

After a preselection which substantially reduces the background while keeping about 70% of the signal, six sequential cuts are applied: number of jets, missing energy, acollinearity, thrust angle, transverse momentum, and the jet-jet invariant mass and c -tagging [1]. The background consists of the following processes: W^+W^- , ZZ , $W\epsilon\nu$, eeZ , $q\bar{q}(q \neq t)$, $t\bar{t}$, and two-photon. After all the cuts, the total background of about 5680 events is dominated by about 5044 $W\epsilon\nu$ events [1]. A scalar top signal of 120 GeV has been simulated with a neutralino mass of 110 GeV. The selection efficiency is about 20% [14] and 11,500 signal events are expected for a standard LCFI vertex detector configuration as given in the TESLA TDR.

3. Varying the vertex detector design

This study of the vertex detector design is based on 50,000 simulated 120 GeV signals and 210,000 $W\epsilon\nu$ background events for each detector design. After preselection 29,842 signals and 53,314 $W\epsilon\nu$ events are selected, corresponding to 34,318 and 779,450 events per 500 fb⁻¹, respectively. This preselection signal efficiency of

59.7% does not depend on the vertex detector design. Four detector designs are compared:

- VX₁₂: The TESLA TDR design with 5 layers and a single layer (0.064% X_0 radiation length per layer).
- VX₂₂: Four layers (the innermost layer removed). This scenario could for example occur if the vertex detector is exposed to a large dose of machine background from the accelerator. The optimization of the radius of the innermost layer is an important aspect in the design of a vertex detector for a linear collider.
- VX₃₂: Five layers and double material thickness (0.128% X_0 radiation length per layer). As the rigidity of the sensitive elements and the support structure is another important aspect in the detector design, the material budget has to be taken into account.
- VX₄₂: Four layers (the innermost layer removed) and double thickness (0.128% X_0 radiation length per layer).

The c -tagging efficiency per event is normalized to the number of signal events after the preselection and requiring two jets. At least one c -tagged jet is required and the efficiency is given in figure 1 and table 2 as a function of the purity, where purity is defined as the ratio of the number of simulated signal events after c -tagging to all the c -tagged events assuming the same luminosity for signal and background. The different purities are obtained by varying the cut on the c -tagging neural network variable. The effect of the detector design variation increases with increasing purity (harder c -tagging neural network cut). For the second set of points in the plot with purities about 18% and c -tagging efficiencies between 85% and 90%, the variation of the signal efficiency and the number of $W e \nu$ background events are given in table 2 after all selection cuts.

4. Results

These results for small visible energy ($\Delta m = 10$ GeV) lead to the same observation as for large visible energy (SPS-5 scenario with $\Delta m = 100.7$ GeV). The radius of the innermost layer of the vertex detector has a large effect on the c -quark tagging performance. Curves VX₁₂ and VX₂₂ in figure 1 show the performance for radii 15 and 26 mm, respectively. There is no significant effect on the c -quark tagging performance from doubling the material budget (e.g. curves VX₁₂ and VX₃₂ of

Table 2. Expected signal efficiency and number of $W e \nu$ background events for the four detector designs. The statistical uncertainties are also given.

Thickness	Layers	Signal efficiency (in %)	$W e \nu / 210$ k	$W e \nu / 500$ fb ⁻¹
Single	5 (4)	20.46 (19.67)	364 (369)	5322 (5396)
Double	5 (4)	20.32 (19.52)	366 (385)	5352 (5630)
		± 0.18	± 19	± 280

Scalar top study




Figure 1. Scalar top c -tagging efficiency and purity with $W\ell\nu$ background for different detector designs. The VX_{12} curve is for a detector design with five layers (innermost at 15 mm) and single density, curve VX_{22} is for a detector design with four layers (innermost at 26 mm). Curves VX_{32} and VX_{42} are for double density (0.128% X_0 radiation length per layer) with four and five layers, respectively.

figure 1). The increase of multiple scattering is not significant between single and double thicknesses (0.128% X_0 radiation length per layer).

In order to quantitatively estimate the multiple scattering effect, the number of tracks per signal event and the visible energy have been determined. The minimum visible energy per event is about 10 GeV and the maximum number of tracks is about 20, therefore the minimum track energy is about 0.5 GeV. The analytical calculation of the multiple scattering angle is given by $\theta \approx 13.6/P \cdot \sqrt{x/X_0}$, where the track momentum P is given in MeV. The displacement at the interaction point is $d \approx R\theta$, where R is the radius of the innermost layer of the vertex detector. For $P = 500$ MeV, $x/X_0 = 0.128\%$ and $R = 15$ mm, $d = 15 \mu\text{m}$. This small value compared to the flight distance of charm mesons explains the insignificant effect on the c -quark tagging from the multiple scattering increase by doubling the vertex detector layer thickness.

5. Conclusions

In conclusion, studies with a small visible energy signal lead to the same results as in the previous study for large visible energy regarding the vertex detector design. A strong dependence on the detector performance on the radius of the innermost vertex detector layer has been observed, while doubling the material thickness has no significant effect on the c -quark tagging performance in scalar top studies at the ILC. The optimization of the radius of the innermost vertex detector layer will have to take into account the accelerator background which depends on the machine optics and the collimation system.

Acknowledgements

The author AS would like to thank the organizers of the workshop for making this presentation possible.

References

- [1] M Carena, A Finch, A Freitas, C Milstène, H Nowak and A Sopczak, *Phys. Rev.* **D72**, 115008 (2005)
- [2] C Milstène and A Sopczak, Econf C0508141:ALCPG1431, 2005
- [3] A Finch, H Nowak and A Sopczak, CCD vertex detector charm-tagging performance in studies of scalar top quark decays, *Proc. of the Int. Conference on Linear Colliders (LCWS 04)*, Paris, France, 19–24 April 2004
- [4] A Sopczak, M Carena, A Finch, A Freitas, C Milstène and H Nowak, Scalar top quark studies with various visible energies, presented at the *13th Int. Conference on Supersymmetry and Unification of Fundamental Interactions (SUSY'05)*, Durham, UK, 18–23 July 2005
- [5] T Sjöstrand *et al*, *Comput. Phys. Commun.* **135**, 238 (2001)
- [6] A Sopczak, in PHYSICS AT LEP2, Event Generators for Discovery Physics, CERN Yellow Report CERN 96-01
- [7] A Finch, H Nowak and A Sopczak, contributed paper EPS370, *Int. Conference on High-Energy Physics (HEP 2003)*, Aachen, Germany, 17–23 July 2003 (LC Note LC-PHSM-2003-075)
- [8] M Pohl and H J Schreiber, hep-ex/0206009
- [9] M Berggren, Simulation Grand Vitesse (SGV), <http://berggren.home.cern.ch/berggren/sgv.html>
- [10] A Freitas, D J Miller and P M Zerwas, *Euro. Phys. J.* **C21**, 361 (2001)
A Freitas, A von Manteuffel and P M Zerwas, *Euro. Phys. J.* **C34**, 487 (2004)
- [11] F Yuasa *et al*, *Prog. Theor. Phys. Suppl.* **138**, 18 (2000)
- [12] CompHEP Collaboration: E Boos *et al*, *Nucl. Instrum. Methods* **A534**, 250 (2004)
- [13] T Kuhl, Hadronic branching ratio of a SM-like Higgs boson at a future linear collider, *Proc. of the Int. Conference on Linear Colliders (LCWS 04)*, Paris, France, 19–24 April 2004
- [14] The re-tuning of the c -tagging neural network increased the selection efficiency from 19% [1] to 20% and the $W e \nu$ background from 5044 [1] to 5322 ± 280 events