

A search for charged massive long-lived particles at D0

SUDESHNA BANERJEE, for the D0 Collaboration

Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India

E-mail: sudeshna@tifr.res.in

Abstract. Results are presented on a search for charged massive long-lived particles (CMLLPs), based on 5.2 fb^{-1} of integrated luminosity collected with the D0 detector at the Fermilab Tevatron $p\bar{p}$ collider. This search selects events in which one or more particles are reconstructed as muons but their speed and ionization energy loss (dE/dx) are different from muons produced in beam–beam collisions. CMLLPs have been predicted by several theories of physics beyond the Standard Model. In this search we exclude pair-produced long-lived gaugino-like charginos with masses below 251 GeV and higgsino-like charginos with masses below 230 GeV at 95% CL, as well as long-lived stop quarks with masses below 265 GeV. We place cross-section limits on long-lived staus as 0.04 to 0.006 pb for stau masses in the range 100 to 300 GeV.

Keywords. Charged massive long-lived particle; D0.

PACS Nos 13.85.Rm; 14.80.Ly

Historically, the strange, long-lived kaons started a new era in particle physics. Today some extensions of Standard Model and some SUSY models suggest the existence of another kind of massive long-lived particles (MLLP) [1]. Existence of these particles could solve some of the puzzles that have no answers yet, namely, dark matter in the universe. Lithium abundance, which is not explained by the model of Big Bang nucleosynthesis (BBN) might be explained by the existence of a MLLP that decays during or after the time of BBN [2]. These massive long-lived particles could have colour and/or electric charge. They appear in many supersymmetric (SUSY) models as bound states of squarks or gluinos (R -hadrons) or as sleptons or charginos. Current results are from a search of charged massive long-lived particles (CMLLP) performed with 5.2 fb^{-1} of data collected by the D0 experiment at the Tevatron $p\bar{p}$ collider at $\sqrt{s} = 1.96 \text{ TeV}$.

1. Models

We have used SUSY models to study the CMLLPs. However, CMLLPs can occur in models other than SUSY and our search will have some sensitivity for other types of

CMLLPs which have similar production kinematics. We know from cosmology that in SUSY models, the lightest supersymmetric particle (LSP) is stable and must be neutral. Next-to-lightest supersymmetric particles (NLSP) could be long-lived due to weak couplings to the LSPs or due to mass degeneracy with the LSPs. Such NLSPs could be CMLLP candidates. Four different types of CMLLPs are considered in this analysis. In the gauge-mediated SUSY breaking (GMSB) models, a tau slepton (stau) NLSP can be a CMLLP if its decay to the gravitino LSP is suppressed. A chargino can be a CMLLP candidate in those models where it is the NLSP and its mass difference with the LSP of the model is small. Two extreme scenarios are explored, where the chargino is mostly a gaugino or it is mostly a higgsino. The top squark (stop) can be a CMLLP candidate if it is the lightest coloured particle. In some hidden valley theories stop is an LSP and hence long-lived [3].

2. The D0 detector and CMLLP signatures

The D0 detector is a multipurpose detector consisting of three primary systems: a central tracking system, calorimeters and a muon spectrometer [4]. The parts relevant for this search are the muon system for the time-of-flight (TOF) measurement and the silicon microstrip tracker (SMT) for the dE/dx measurement. The muon system has wire chambers for muon tracking and scintillation counters for triggering on muons. There are three layers in the muon system with a 1.8 Tesla toroid between the first and the second layers. The SMT is the innermost part of the tracking system and has a six-barrel longitudinal structure, where each barrel consists of a set of nine layers in the central region which gives an accurate momentum and dE/dx measurement.

The properties of CMLLPs which can be used to detect them are, charge, due to which they leave a track in the tracker, long lifetime, due to which they do not decay inside the detector, and large mass (compared to muons), due to which they have low speed. From these properties one can conclude that they act like muons but because they are much heavier, their TOF to the muon system and dE/dx in the SMT are expected to be much larger than those of muons produced in $p\bar{p}$ collisions. For this search, events with one or more muons have been chosen and the highest p_T muon is required to satisfy the criteria of being a CMLLP candidate.

3. Signal, background, and data samples

To obtain signal samples, CMLLP pairs are generated using PYTHIA(6.409) [5]. Cascade decays are not included to keep the analysis model-independent. Staus, stops, and charginos (gaugino-like and higgsino-like) are generated at masses of 100, 150, 200, 250, and 300 GeV. Stop quarks are hadronized by linking an external routine with PYTHIA [6]. These samples are then processed through a GEANT3-based simulation of the D0 detector. The background contribution comes from mismeasured muons (wrong speed (β) and dE/dx), the biggest source of which is $W \rightarrow \mu\nu$ events.

This search is based on 5.2 fb^{-1} of data taken during June 2006 to June 2009. Events are triggered by a single high $p_T (> 20 \text{ GeV})$ muon. These events are then checked further using the following selection criteria. There should be at least one isolated muon of good quality in the event with $p_T > 60 \text{ GeV}$, $|\eta| < 1.6$, and $\beta < 1$. The muon track in the muon chamber should have a matching partner in the SMT and the speeds of the muon measured in different layers of the muon system should be consistent with each other. Since background events are mostly $W \rightarrow \mu\nu$ decays, the shape of the background is determined using a W -rich dataset selected by requiring $M_T < 200 \text{ GeV}$ [$M_T = \sqrt{(E_T + \cancel{E}_T)^2 - (p_x + \cancel{E}_x)^2 - (p_y + \cancel{E}_y)^2}$]. The background normalization is done using a signal-free dataset chosen with $\beta > 1$.

4. Analysis technique

It is observed that the requirement of $p_T > 60 \text{ GeV}$ for the highest p_T muon in the event reduces the background from W decays substantially. Additionally the two variables, β and dE/dx , help to further distinguish signal from background events. This can be seen in figure 1. The following unique feature of stop hadrons is treated separately. Stop quarks hadronize into neutral and charged mesons and baryons [7]. These hadrons flip charge due to nuclear interactions while passing through matter. In order to be detected in the D0 detector they have to be charged, at least at the entry point to the SMT and at the entry and exit points in the muon system. This feature reduces the efficiency of stop detection. In this analysis at least one of the stop or antistop hadrons needs to be charged in order to be detected. The probability for that is 38%. This is used as an additional efficiency factor after imposing the other selection criteria.

The boosted decision trees (BDT), a multivariate technique, is used for final signal selection. The BDT is trained on signal and background distributions of β , dE/dx , and several related variables. The BDTs are applied to signal, background, and data distributions to get a final variable (BDT output) distribution. Figure 2 shows the BDT distributions for 300 GeV gaugino-like chargino signal, background, and data. A CLS




Figure 1. β and dE/dx distributions of the highest p_T muon in the event for data, background, and gaugino-like charginos. Each distribution is normalized to have the same number of events.




Figure 2. BDT outputs for data, background, and signal (gaugino-like chargino of mass 300 GeV) are shown. The signal histogram is normalized to the expected number of events.

method is used to get limits at 95% CL on production cross-sections of CMLLPs [8]. Inputs to the limit setting procedure are the BDT distributions and systematic errors.

5. Results and conclusions

We searched for pair produced charged massive long-lived particles in 5.2 fb^{-1} of data from the D0 experiment. We obtained a lower mass limit for stop quarks of 265 GeV. This limit is enhanced to 281 GeV if it is assumed that stop hadrons do not flip charge in a nuclear interaction. The lower mass limit for gaugino-like chargino is 251 GeV and that for higgsino-like chargino is 230 GeV. Cross-section limits for staus are obtained as 0.04 to 0.006 pb for stau masses between 100 and 300 GeV. The mass limits on the charginos are currently the best in the world. A paper based on this search has now been submitted for publication [9].

References

- [1] M Byrne, C Kolda and P Regan, *Phys. Rev.* **D66**, 075007 (2002)
- [2] Particle Data Group: K Nakamura *et al*, *J. Phys.* **G37**, 075021 (2010), Section 20.5
- [3] J Gunion and S Mrenna, *Phys. Rev.* **D62**, 015002 (2000)
C Chen, M Drees and J Gunion, *Phys. Rev.* **D55**, 330 (1997)
G F Giudice and A Rattazzi, *Phys. Rep.* **322**, 419 (1999)
M Strassler, [arXiv:hep-ph/0607160](https://arxiv.org/abs/hep-ph/0607160)
- [4] D0 Collaboration: V M Abazov *et al*, *Nucl. Instrum. Methods Phys. Res. Sect.* **A565**, 463 (2006)
- [5] T Sjostrand, L Lonnblad, S Mrenna and P Skands, Physics and manual, hep-ph/0308153
- [6] The code for generating stop quarks is at <http://projects.hepforge.org/pythia6/examples/main78.f>
- [7] M Fairbairn *et al*, *Stable massive particles at colliders*, [arXiv:hep-ph/0611040v2](https://arxiv.org/abs/hep-ph/0611040v2), Table 3, p. 40
- [8] W Fisher, FERMILAB Report No. TM-2386-E (2007)
- [9] V M Abazov *et al*, [arXiv:1110.3302](https://arxiv.org/abs/1110.3302), 14 October 2011