

Oscillating solitons in nonlinear optics

XIAO-GANG LIN¹, WEN-JUN LIU^{2,*} and MING LEI²

¹Key Laboratory of Optoelectronic Technology & Systems (Chongqing University),
Ministry of Education, Chongqing, China

²School of Science, Beijing University of Posts and Telecommunications,
P.O. Box 122, Beijing 100876, China

*Corresponding author. E-mail: jungliu@bupt.edu.cn

MS received 7 March 2014; revised 28 November 2014; accepted 15 December 2014

DOI: 10.1007/s12043-015-1020-x; ePublication: 11 July 2015

Abstract. Oscillating solitons are obtained in nonlinear optics. Analytical study of the variable-coefficient nonlinear Schrödinger equation, which is used to describe the soliton propagation in those systems, is carried out using the Hirota's bilinear method. The bilinear forms and analytic soliton solutions are derived, and the relevant properties and features of oscillating solitons are illustrated. Oscillating solitons are controlled by the reciprocal of the group velocity and Kerr nonlinearity. Results of this paper will be valuable to the study of dispersion-managed optical communication system and mode-locked fibre lasers.

Keywords. Solitons; nonlinear optics; analytic soliton solutions; Hirota's bilinear method.

PACS Nos 42.65.-k; 05.45.Yv

1. Introduction

The concept of solitons is one of the fundamental unifying ideas in modern physics and mathematics [1]. The soliton theory can be applied to diverse branches of physics such as plasmas physics, fluid dynamics, nonlinear optics, Bose–Einstein condensates (BECs), and nuclear physics [2–9]. The study of solitons in those physical systems reveals some exciting problems from both fundamental and application points of view [10–20].

In nonlinear optics, solitons are the focus of intense research interest due to their potential applications in telecommunication and ultrafast signal processing systems [21]. Solitons can be regarded as the balance between group velocity dispersion (GVD) and nonlinear effects [22]. They have various applications in pulse amplification, optical switch and pulse compression [23], and some studies on solitons have been carried out both theoretically and experimentally [24–28]. General analytic soliton solutions have been investigated [24], and nonlinear optics models giving rise to the appearance of solitons in a narrow sense have been considered [25]. Gazeau [26] was concerned with the

evolution of solitons driven by random polarization mode dispersion. Under some parametric conditions, Choudhuri and Porsezian [27] have solved the higher-order nonlinear Schrödinger (NLS) equation with non-Kerr nonlinearity, and the interaction dynamics of solitons was reconsidered in [28].

In BECs, solitons can be viewed as fundamental nonlinear excitations [29]. Recently, with various methods, bright, dark and gap matter-wave solitons in BECs have been created [30]. These solitons have attracted much theoretical and experimental interests [31–34]. Some matter-wave solitons in a system of three-component Gross–Pitaevskii equation arising from the context of spinor BECs with time-modulated external potential and scattering lengths are presented in [31]. A family of non-autonomous soliton solutions of BECs with the time-dependent scattering length in an expulsive parabolic potential was obtained in [32]. Becker *et al* [33] studied the dynamics of apparent soliton stripes in elongated BECs experimentally and theoretically. Kanna *et al* [34] studied the dynamics of non-autonomous solitons in two- and three-component BECs.

In this paper, we shall study the properties and features of oscillating solitons in non-linear optics, which can be described by the following variable-coefficient NLS (vcNLS) equation [21]:

$$i \frac{\partial u}{\partial z} + i \beta_1(z) \frac{\partial u}{\partial t} + \beta_2(z) \frac{\partial^2 u}{\partial t^2} + \gamma(z) |u|^2 u = 0, \quad (1)$$

where $u(z, t)$ is the temporal envelope of solitons, z is the longitudinal coordinate and t is the time in the moving coordinate system. $\beta_1(z)$ is the reciprocal of the group velocity, $\beta_2(z)$ represents the GVD coefficient and $\gamma(z)$ is the nonlinearity coefficient. Equation (1) can reduce to the standard vcNLS equation while $\beta_1(z) = 0$, and some studies on the standard vcNLS equation has been made.

However, eq. (1) has not been investigated due to the existence of $\beta_1(z)$. With the help of the Hirota’s bilinear method, analytic soliton solutions for eq. (1) will be obtained. The relevant properties and features of oscillating solitons will be illustrated. The influence of $\beta_1(z)$ will be analysed for the first time.

This paper is structured as follows. In §2, analytic soliton solutions for eq. (1) will be presented. In §3, the properties and features of oscillating solitons will be discussed, and the influence of $\beta_1(z)$ will be analysed. Finally, our conclusions will be presented in §4.

2. Analytic soliton solutions for eq. (1)

At first, the dependent variable transformation can be introduced as [35–38]

$$u(z, t) = \frac{g(z, t)}{f(z, t)}, \quad (2)$$

where $g(z, t)$ is a complex differentiable function and $f(z, t)$ is a real one. With symbolic computation, the bilinear forms for eq. (1) are obtained as

$$i D_z g \cdot f + i \beta_1(z) D_t g \cdot f + \beta_2(z) D_t^2 g \cdot f = 0, \quad (3)$$

$$\beta_2(z) D_t^2 f \cdot f - \gamma(z) g g^* = 0, \quad (4)$$

where the asterisk denotes the complex conjugate. D_z and D_t [39] are the Hirota's bilinear operators, and are defined by

$$D_z^m D_t^n (a \cdot b) = \left(\frac{\partial}{\partial z} - \frac{\partial}{\partial z'} \right)^m \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^n a(z, t) b(z', t') \Big|_{z'=z, t'=t}. \quad (5)$$

With the following power series expansions for $g(z, t)$ and $f(z, t)$:

$$g(z, t) = \varepsilon g_1(z, t) + \varepsilon^3 g_3(z, t) + \varepsilon^5 g_5(z, t) + \dots, \quad (6)$$

$$f(z, t) = 1 + \varepsilon^2 f_2(z, t) + \varepsilon^4 f_4(z, t) + \varepsilon^6 f_6(z, t) + \dots, \quad (7)$$

where ε is a formal expansion parameter, bilinear forms (3) and (4) can be solved. Substituting expressions (6) and (7) into bilinear forms (3) and (4) and equating coefficients of the same powers of ε to zero, yield recursion relations for $g_n(z, t)$ s and $f_n(z, t)$ s. Then, the analytic soliton solutions for eq. (1) can be derived.

To get the analytic soliton solutions for eq. (1), we assume

$$g(z, t) = g_1(z, t), \quad f(z, t) = 1 + f_2(z, t), \quad (8)$$

where

$$g_1(z, t) = e^{\theta_1}, \quad \theta_1 = [a_{11}(z) + i a_{12}(z)] z + (b_{11} + i b_{12})t + k_{11} + i k_{12}, \quad (9)$$

with b_{11}, b_{12}, k_{11} and k_{12} are real constants. $a_{11}(z)$ and $a_{12}(z)$ are the differentiable functions to be determined. With $g_1(z, t)$, and collecting the coefficient of ε in eq. (3), we obtain the constraints on $a_{11}(z)$ and $a_{12}(z)$ as

$$a_{11}(z) = \frac{1}{z} \int [-b_{11}\beta_1(z) - 2b_{11}b_{12}\beta_2(z)] dz,$$

$$a_{12}(z) = \frac{1}{z} \int [b_{11}^2\beta_2(z) - b_{12}\beta_1(z) - b_{12}^2\beta_2(z)] dz.$$

Substituting $g_1(z, t)$ into eq. (4), and collecting the coefficient of ε^2 , we get

$$f_2(z, t) = M e^{\theta_1 + \theta_1^*} \quad (10)$$

where

$$M = \frac{\gamma(z)}{8b_{11}^2\beta_2(z)}, \quad \beta_2(z) = c \gamma(z),$$

and c is an arbitrary constant.

Without loss of generality, we set $\varepsilon = 1$, and can write the analytic soliton solutions as

$$u(z, t) = \frac{g(z, t)}{f(z, t)} = \frac{g_1(z, t)}{1 + f_2(z, t)}$$

$$= \frac{1}{\sqrt{M}} e^{i[a_{12}(z)z + b_{12}t + k_{12}]} \operatorname{sech} \left[a_{11}(z)z + b_{11}t + k_{11} + \frac{1}{2} \ln M \right]. \quad (11)$$

3. Discussion

In figure 1, oscillating solitons are exhibited in nonlinear optics. In this case, nonlinearity is assumed to be constant; $\gamma(z) = 0.01$ in figure 1. When $\beta_1(z) = 2 \exp(-0.5z) + \sin(2z)$ as in figure 1a, the solitons are stable at first, and then exhibit periodic soliton propagation. We can adjust the soliton propagation status in comparison with figure 1a. By changing $\beta_1(z)$ the stable solitons can be changed to oscillating solitons when $\beta_1(z) = 2 \exp(-0.5z^2) + \sin(2z)$ (see figure 1b).

If nonlinearity $\gamma(z)$ is also taken as a function in figure 1b, the periodic and oscillation state of solitons can be changed. When $\gamma(z) = \sin(z)$, the periodicity of soliton increases, and the oscillation of solitons is enhanced (figure 2a). Increasing the coefficient of $\gamma(z)$ results in the decrease of the periodic and oscillation state of solitons, which can be seen in figure 2b with $\gamma(z) = 1.5 \sin(2z)$.

By changing the values of $\beta_1(z)$ and $\gamma(z)$, we get different propagation status of solitons in figure 3. For $\beta_1(z)$ and $\gamma(z)$, if we transform the function type, the change of the phase shift is opposite. Moreover, when $\beta_1(z)$ and $\gamma(z)$ are linear combinations of




Figure 1. Solitons in nonlinear optics. The parameters are $c = 1, k_{11} = -1, k_{12} = 2, b_{11} = 1, b_{12} = -1, \gamma(z) = 0.01$ with (a) $\beta_1(z) = 2 \exp(-0.5z) + \sin(2z)$, (b) $\beta_1(z) = 2 \exp(-0.5z^2) + \sin(2z)$.




Figure 2. Solitons in nonlinear optics with the same parameters as given in figure 1, but with (a) $\beta_1(z) = 2 \exp(-0.5z^2) + \sin(2z), \gamma(z) = \sin(z)$, (b) $\beta_1(z) = 2 \exp(-0.5z^2) + \sin(2z), \gamma(z) = 1.5 \sin(2z)$.




Figure 3. Solitons in nonlinear optics with the same parameters as given in figure 1, but with (a) $\beta_1(z) = 1.5 \sin(2z)$, $\gamma(z) = \exp(-0.5z^2)$, (b) $\beta_1(z) = 1.5 \sin(2z) + \cos(z)$, $\gamma(z) = 1.5 \cos(2z) + \sin(z)$.

trigonometric functions with different periods, the periodic propagation status of solitons is disrupted as shown in figure 3b. Soliton oscillations are sometimes weakened, and sometimes exacerbated. Thus, the soliton propagation status can be controlled with $\beta_1(z)$ and $\gamma(z)$ in nonlinear optics.

4. Conclusions

In this paper, oscillating solitons in nonlinear optics were obtained. The vNLS equation (see eq. (1)), which can be used to describe the soliton propagation, was investigated analytically. The bilinear forms (3) and (4) were derived, and the analytic soliton solutions (11) were obtained. According to solutions (11), the oscillating solitons were exhibited with different values of $\beta_1(z)$ and $\gamma(z)$ (see figures 1–3), and the influences of $\beta_1(z)$ and $\gamma(z)$ were analysed. The results have shown that the status of the soliton propagation can be controlled with $\beta_1(z)$ and $\gamma(z)$ in nonlinear optics. Results of this paper might be of potential use in the design of the dispersion-managed optical communication system and mode-locked fibre lasers.

Acknowledgements

The authors express their sincere thanks to the Editors and referee for their valuable comments. This work has been supported by the National Natural Science Foundation of China under Grant No. 61205064, by Fund of State Key Laboratory of Information Photonics and Optical Communications (Beijing University of Posts and Telecommunications), and by the Visiting Scholar Funds of the Key Laboratory of Optoelectronic Technology and Systems under Grant No. 0902011812401_5, Chongqing University.

References

- [1] S K Turitsyn, B G Bale and M P Fedoruk, *Phys. Rep.* **521**, 135 (2012)
- [2] E Infeld, *Nonlinear waves, solitons and chaos*, 2nd edn (Cambridge University Press, Cambridge, UK, 2000)

- [3] P K Shukla and A A Mamun, *New J. Phys.* **5**, 17.1 (2003)
- [4] Y Kodama, *J. Phys. A* **43**, 434004 (2010)
- [5] R Yang, R Hao, L Li, Z Li and G Zhou, *Opt. Commun.* **242**, 285 (2004)
- [6] S Burger, K Bongs, S Dettmer, W Ertmer, K Sengstock, A Sanpera, G V Shlyapnikov and M Lewenstein, *Phys. Rev. Lett.* **83**, 5198 (1999)
- [7] L Khaykovich, F Schreck, G Ferrari, T Bourdel, J Cubizolles, L D Carr, Y Castin and C Salomon, *Science* **296**, 1290 (2002)
- [8] K E Strecker, G B Partridge, A G Truscott and R G Hulet, *Nature* **417**, 150 (2002)
- [9] E R Arriola, W Broniowski and B Golli, *Phys. Rev. D* **76**, 014008 (2007)
- [10] H Triki, F Azzouzi and P Grelu, *Opt. Commun.* **309**, 71 (2013)
- [11] R Guo, H Q Hao and L L Zhang, *Nonlinear Dynam.* **74**, 701 (2013)
- [12] R Guo and H Q Hao, *Commun. Nonlinear Sci. Numer. Simulat.* **18**, 2426 (2013)
- [13] C Q Dai and H P Zhu, *J. Opt. Soc. Am. B* **30**, 3291 (2013)
- [14] C Q Dai and H P Zhu, *Ann. Phys. (NY)* **341**, 142 (2014)
- [15] B Tang, D J Li and Y Tang, *Can. J. Phys.* **91**, 788 (2013)
- [16] J D Moores, *Opt. Lett.* **21**, 555 (1996)
- [17] V N Serkin and A Hasegawa, *IEEE J. Sel. Top. Quant. Electron.* **8**, 418 (2002)
- [18] V I Kruglov, A C Peacock and J D Harvey, *Phys. Rev. Lett.* **90**, 113902 (2003)
- [19] T S Raju, P K Panigrahi and K Porsezian, *Phys. Rev. E* **71**, 026608 (2005)
- [20] R Atre, P K Panigrahi and G S Agarwal, *Phys. Rev. E* **73**, 056611 (2006)
- [21] G P Agrawal, *Nonlinear fiber optics*, 4th edn (Academic Press, San Diego, 2007)
- [22] Z G Chen, M Segev and D N Christodoulides, *Rep. Prog. Phys.* **75**, 086401 (2012)
- [23] M Saha and A K Sarma, *Commun. Nonlinear Sci. Numer. Simulat.* **18**, 2420 (2013)
- [24] H Triki and T R Taha, *Math. Comput. Simulat.* **82**, 1333 (2012)
- [25] A I Maimistov, *Quant. Electron.* **40**, 756 (2010)
- [26] M Gazeau, *J. Opt. Soc. Am. B* **30**, 2443 (2013)
- [27] A Choudhuri and K Porsezian, *Phys. Rev. A* **88**, 033808 (2013)
- [28] R Radha, P S Vinayagam and K Porsezian, *Phys. Rev. E* **88**, 032903 (2013)
- [29] R Carretero-González, D J Frantzeskakis and P G Kevrekidis, *Nonlinearity* **21**, R139 (2008)
- [30] Z J Li, W H Hai and Y Deng, *Chin. Phys. B* **22**, 090505 (2013)
- [31] D S Wang, Y R Shi, K W Chow, Z X Yu and X G Li, *Eur. Phys. J. D* **67**, 242 (2013)
- [32] Q Y Li, Z D Li, L Li and G S Fu, *Opt. Commun.* **283**, 3361 (2010)
- [33] C Becker, K Sengstock, P Schmelcher, P G Kevrekidis, R Carretero-González, *New J. Phys.* **15**, 113028 (2013)
- [34] T Kanna, R B Mareeswaran, F Tsitoura, H E Nistazakis and D J Frantzeskakis, *J. Phys. A* **46**, 475201 (2013)
- [35] W J Liu, B Tian, H Q Zhang, T Xu and H Li, *Phys. Rev. A* **79**, 063810 (2009)
- [36] W J Liu, B Tian, H Q Zhang, L L Li and Y S Xue, *Phys. Rev. E* **77**, 066605 (2008)
- [37] W J Liu, B Tian and H Q Zhang, *Phys. Rev. E* **78**, 066613 (2008)
- [38] W J Liu, B Tian, T Xu, K Sun and Y Jiang, *Ann. Phys. (NY)* **325**, 1633 (2010)
- [39] R Hirota, *Phys. Rev. Lett.* **27**, 1192 (1971)