

ON A RELATION BETWEEN TWO CONJECTURES OF THE THEORY OF NUMBERS.

BY S. CHOWLA.

(Government College, Lahore.)

Received November 11, 1936.

§1. Let $r_{k,s}(n)$ denote the number of representations of n as a sum of s k th powers of integers ≥ 0 . Since $r_{k,2}(n) = O(n^\epsilon)$ for every positive ϵ , we have

$$r_{k,k}(n) = O\left(n^{1 - \frac{2}{k} + \epsilon}\right)$$

and no better result is known. Denote by (A_s) the unproved result that

$$(A_s) \quad r_{k,k}(n) = O\left(n^{\frac{s}{k}}\right)$$

where $k \geq 5$ and s is a positive integer $\leq k - 3$. Denote by (B_s) the conjecture that

$$(B_s) \quad \sum_{m=1}^{k-s} x_m^k = \sum_{m=1}^{k-s} y_m^k \text{ has a non-trivial solution in positive integers } x_1, x_2, \\ \text{etc., } y_1, y_2, \text{ etc.}$$

Then we have the

*Theorem.** If (A_s) is false then (B_s) is true, and conversely.

In other words at least one of (two desirable results) (A_s) and (B_s) is true.

Proof. If (B_s) is false, then

$$r_{k,k-s}(n) = O(1)$$

whence

$$r_{k,k-s+1}(n) = O\left(n^{\frac{1}{k}}\right)$$

$$r_{k,k-s+2}(n) = O\left(n^{\frac{2}{k}}\right)$$

etc.

$$r_{k,k}(n) = O\left(n^{\frac{s}{k}}\right)$$

* The proof shows that the result is trivial; my object is to draw attention to the conjectures themselves.

i.e., if (B_s) is false then (A_s) is true. Conversely, if (A_s) is false, it follows that (B_s) must be true.

§2. Note the special case when $s = 3$ ($k \geq 6$). at least one of the following conjectures is true :

$$(A) \quad a_1^k + a_2^k + a_3^k = b_1^k + b_2^k + b_3^k$$

has a non-trivial solution in positive integers, $a_1, a_2, a_3, b_1, b_2, b_3$.

$$(B) \quad r_{k,k}(n) = O\left(n^{\frac{k-3}{k}}\right)$$

It seems to the writer that (A) is *not likely* to be proved in the near future. On the other hand, it is possible that (B), which asserts so much less than "Hypothesis K" of Hardy and Littlewood, is not a difficult result. We observe also that (A) *may be false*, for solutions of (A) are not known even for $k = 5$.